

**UNICORN**

**PERMETHRIN DUST**

**For Livestock and Pets**

**ACTIVE INGREDIENT:**

Permethrin (3-phenoxyphenyl) methyl (±) cis, trans-3-(2,2-dichloroethenyl)-2,2- dimethylcyclopropanecarboxylate *.....	0.25%
--	-------

<b>INERT INGREDIENTS:.....</b>	<b>99.75%</b>
--------------------------------	---------------

* Cis/trans ratio:	Min. 35% (±) cis
	Max. 65% (±) trans

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**See Side/Back Panel for Additional Precautions**

**NET CONTENTS: 6 OUNCES**

**MANUFACTURED BY:**

Unicorn Laboratories  
1002 118th Avenue North  
St. Petersburg, Florida 33716

**EPA REG. NO. 28293-126(5/19/92)(E)**

**EPA EST. NO. 28292-FL-1**

**ACCEPTED**

MAR 22 1994

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 28293-126.

503

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed, inhaled. Avoid breathing dust. Avoid contact with eyes and skin. Wash thoroughly with soap and water after handling. Avoid contamination of feed and foodstuffs.

**Statement of Practical Treatment**

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**Physical and Chemical Hazards**

Do not use or store near heat or open flame.

**Environmental Hazards**

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**BEEF AND DAIRY CATTLE AND HORSES:**

Can be used in dust bags, shaker can and mechanical dust applicator. **Horn Flies, Lice, Face Flies:** Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through which the animals must pass daily for water, feed, or minerals. Bags may also be placed in loafing sheds or in front of mineral feeders. For dairy cows, bags may be suspended in the exit through which the cows leave the milking barn. The bags should hang 4 to 6 inches below the back line of the cattle. For reduction of Face Flies, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted.

**Horn Flies, Lice Direct Application:** Apply 2 oz. of dust per animal by shaker can over the head, neck, shoulders and back and tailhead. Repeat as necessary.

**HORSES:** Dust horse for Hornflies with shaker can or dust glove along back, neck and legs.

**SWINE:** Lice on Swine Direct Application: Apply only 1 ounce per head as a uniform coat to the head, shoulder and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary but not more often than once every 10 days. In severe infestation, both animals and the bedding may be treated. Do not ship animals for slaughter within 5 days of treatment.

**POULTRY:** To control Northern Fowl Mites, apply at a rate of 1 lb per 100 birds. Ensure thorough treatment of vent area.

**PETS:** To control Fleas, Ticks and Lice on Cats and Dogs, dust animals thoroughly, working the dust into the hair down to the skin. Pay close attention to legs and feet. Use 1/2 oz. on animals 20 pounds and under and 1 oz. on pets over 20 pounds. Avoid getting dust in pet's eyes. Repeat treatment as needed. Dust sleeping area including bedding and cracks and crevices.

**STORAGE AND DISPOSAL**

**STORAGE:** Store in cool, dry place. Do not contaminate feed or foodstuffs by storage or disposal.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.