

128293-126

04-07-2010

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Ms. Rebecca M. Horton
Phaeton Corporation
P.O. Box 290
Madison, GA 30650

APR 7 2010

Subject: Label Notification(s) for Pesticide Registration 2007-4 & PR Notice 2008-1

Dear: Ms. Horton:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated for March 9, 2010

EPA Registration Number: 28293-126 Unicorn Permethrin Dust

The Registration Division (RD) has conducted a review of this request for applicability Under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please re reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on non-refillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact Melody Banks on 703 305-5413.

Sincerely,

A handwritten signature in black ink, appearing to read "Richard Gebken".

Richard Gebken
Product Manager 11
Insecticide Branch
Registration Division (7505P)



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 28293-126	2. EPA Product Manager Suarez	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Phaeton Corporation / Unicorn Permethrin Dust	PM# Team 13	
5. Name and Address of Applicant (Include ZIP Code) Phaeton Corporation P.O. Box 290 Madison, GA 30650 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

APR 7 - 2010

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION of label changes per June 4, 2009 letter from Environmental Protection Agency and PR Notices 2008-1, 2007-4, 98-10 and 83-3.

This notification is consistent with the provisions of Agency letter dated June 4, 2009, PR Notices 2008-1, 2007-4, 98-10 and 83-3 and the requirements of EPA regulations at 40 CFR 156.10, 156.140, 156.144, 156.146 and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. Phaeton Corporation understands that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA, and further understands that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146 and 156.156, this product may be in violation of FIFRA and Phaeton Corporation may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input checked="" type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container

3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container 1, 2, 5 LB	5. Location of Label Directions <input checked="" type="checkbox"/> on label/attached booklet
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued Stenciled <input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Rebecca M. Horton	Title Consultant/Agent for PHAETON CORPORATION	Telephone No. (Include Area Code) 540-375-8823
---------------------------	---	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received (Stamped)

2. Signature <i>Rebecca M. Horton</i>	3. Title Consultant/Agent for PHAETON CORPORATION	6. Date Application Received (Stamped)
4. Typed Name Rebecca M. Horton	5. Date March 9, 2010	

Registrations
by Design, Inc.™

CERTIFIED MAIL #7009 0080 0002 3611 3686
March 9, 2010

██████████
██████████
██████████
phone 540.375.8826
fax 540.375.8827

Document Processing Desk (NOTIF)
Office of Pesticide Programs/Registration Division (7504P)
U.S. Environmental Protection Agency
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, DC 20460

Re: Notification: EPA Reg. No. 28293-126, Unicorn Permethrin Dust

Attn: Mark Suarez
PM Team 13

On behalf of Phaeton Corporation of Madison, Georgia, enclosed is notification of the following label changes for the above pesticide product:

- 1) revised Storage/Disposal text per PR Notices 83-3 and 2007-4;
- 2) revised Environmental Hazards text per PR Notice 2008-1;
- 3) revised Environmental Hazards and general labeling for pyrethroid non-agricultural outdoor products notification -- per June 4, 2009 EPA letter (R.Gebken)

The following documents are enclosed:


- 1) EPA Form 8570-1, Application for Pesticide – Notification (2 copies)
- 2) Copy of the revised label with areas of revision highlighted in blue.

Best regards,

Rebecca M. Horton
Consultant/Agent for
PHAETON CORPORATION
e-mail: hortonb@ntelos.net

/bmh

Enclosures



Mailing address:
P.O. Box 1019, Salem, VA 24153

Federal & State Pesticide/Fertilizer
Registration Services plus
typesetting/design of product labeling

UNICORN PERMETHRIN DUST

[] Indicates alternate or optional wording

For Use On: [Can be used on]

- Livestock
- Pets

- Home Ornamentals
- Home Gardens

[Ready to Use]

- Kills [over] 60 [60+] [types of] insects
[on Vegetables, Ornamentals, Fruits]

NOTIFICATION

APR 7 - 2010

[Kills Bugs (Apple Aphid, Armyworm, Artichoke Plume Moth, asparagus Beetle, Aster Leafhopper, Beet Armyworm, Cabbage Aphid, Cabbage Looper, Codling Moth, Colorado Potato Beetle, Corn Earworm, Corn Rootworm (Adult), Cucumber Beetle (Adult), Cutworm, Cutworm Complex, Diamondback Moth, European Corn Borer, Fall Armyworm, Filbertworm, Flea Beetle, Fruit Fly, Granulate Cutworm, Green Fruitworm, Hornworm, Imported Cabbageworm, Japanese Beetle, Leafhopper, Leafminer, Lepidopterous Larvae, Lesser Peachtree Borer, Lygus Bug, Melon Worm, Mirid Bug, Navel Orangeworm, Obliquebanded Leafroller, Omnivorous Leafroller, Onion Maggot (Adult), Onion Thrips, Oriental Fruit Moth, Peach Twig Borer, Pear Pyslia, Pepper Weevil, Pickle Worm, Plant Bug (including Lygus and Stink Bug), Plum Curcullo, Potato Aphid, Potato Flea Beetle, Potato Leafhopper, Potato Pysllid, Potato Tuberworm, Redbanded Leafroller, Rindworm, Rose Chafer, Southern Armyworm, Spotted Tentiform Leafminer, Squash Bug, Squash Vine Borer, Stink Bug, Tarnished Plant Bug, Thrips, Tomato Fruitworm, Tomato Pinworm, Vegetable Leafminer, Walnut Husk Fly, White Apple Leafhopper)]

[Kills Beetles, Caterpillars & Bugs] [Kills Japanese Beetles, Aphids, Horn Worms and other listed pests]

- Controls Fleas & Ticks on Dogs & Cats [Use on Dogs & Cats to Kill Fleas, Tick and Lice]

- Effective Long Lasting Control [Use on Vegetables, Roses, Flowers & Fruit Trees]

[For Home Gardens – Vegetables, Fruit Trees and Ornamental Plants]

[Use on Home Gardens Fruits, Roses, Vegetables & Ornamental Plants]

[Use on Fruit Trees, Flowers, Fruits & Vegetables]

[Kills Aphids, Armyworms, Beetles, Caterpillars, Japanese Beetles, Plus others listed! See Back]

[Product Facts: Kills Insects • Beetles (including Japanese & Cucumber Beetles), Caterpillars (including Armyworm, Beet Armyworm, Cabbage Looper, Corn Earworm, Cutworm, Diamondback Moth, Fall Armyworm, Hornworm, Imported Cabbageworm, Lepidopterous Larvae, Lesser Peachtree Borer, Oriental Fruit Moth, Peach Twig Borer, Southern Armyworm, Tomato Fruitworm, Tomato Pinworm), Leafhoppers & other Garden Insects (and other insects as listed) • Plus Fleas & Ticks

Where to Use • In your Garden on Vegetables, Fruits & Ornamentals (including Roses) • On Dogs & Cats]

ACTIVE INGREDIENT:

Permethrin*.....0.25%

INERT [OTHER] INGREDIENTS:.....99.75%

Total: 100.00%

*Cis/trans ratio: Min 35% (±) cis, Max 65% (±) trans

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side/Back Panel [Booklet] for Additional Precautions

NET WT: _____

Manufactured by
PHAETON CORPORATION
 P.O. Box 290, Madison, GA 30650

- READ ENTIRE LABEL BEFORE EACH USE
- FOR PET USE – USE ONLY ON CATS OR DOGS

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION: Humans: Harmful if swallowed, inhaled. Avoid breathing dust. Avoid contact with eyes and skin. Wash thoroughly with soap and water after handling. Avoid contamination of feed and foodstuffs.
Animals: Harmful if swallowed, inhaled. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin Or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. [You may also contact 1-800-222-1222 {or specific company number} for emergency medical treatment information.]	

Physical and Chemical Hazards

Do not use or store near heat or open flame.

Environmental Hazards

This product is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEEF AND DAIRY CATTLE AND HORSES:

Can be used in dust bags, shaker can and mechanical dust applicator. **Horn Flies, Lice, Face Flies:** Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through with the animals must pass daily for water, feed, or minerals. Bags may also be placed in loafing sheds or in front of mineral feeders. For dairy cows, bags may be suspended in the exit through with the cows leave the milking barn. The bags should hang 4 to 6 inches below the back line of the cattle. For reduction of Face Flies, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted. **Horn Flies, Lice Direct Application:** Apply 2 oz. of dust per animal by shaker can over the head, neck, shoulders and back and tailhead. Repeat as necessary.

HORSES: Dust horse for Hornflies with shaker can or dust glove along back, neck and legs.

SWINE: Lice on Swine Direct Application: Apply only 1 ounce per head as a uniform coat to the head, shoulder and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary but not more often than once every 10 days. In severe infestation, both animals and the bedding may be treated. Do not ship animals for slaughter within 5 days of treatment.

POULTRY: To control Northern Fowl Mites, apply at a rate of 1 lb per 100 birds. Ensure thorough treatment of vent area.

PETS: To control [kills] Fleas, Ticks and Lice [use] on Cats and Dogs, dust animals thoroughly, working the dust into the hair down to the skin. Pay close attention to legs and feet. Use ½ oz. on animals 20 pounds and under and 1 oz. on pets over 20 pounds. Avoid getting dust in pet's eyes. Reapply every two (2) weeks as needed. Do not use on cats or dogs under twelve (12) weeks of age. Dust sleeping area including bedding and cracks and crevices.

HOME GARDENS: For Use on Home Gardens. Not for Use on Plants being grown for Sale or other Commercial Use. Do not water the treated area to the point of run-off. Do not make applications during rain

[HOW TO APPLY] Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust guy, puffer or other bulbous duster, rotary duster or other appropriate duster as available. [Wearing household latex or rubber gloves, open shaker side of cap and shake gently to apply. Wearing household latex or rubber gloves, pull red cap open and squeeze container to apply. (directions to be used when packaged in an applicator container – the statement used depends upon type of container packages.)] Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. (Many insects hide on the underside of the leaf). **Apply when air is calm to avoid drift and contact with eyes and skin.** Controls a wide range of insect pests.

[WHEN TO APPLY] Use at first signs of insect infestation, preferable before damage from feeding occurs. **Note the pre-harvest (time between use and harvest) interval with each crop listed.**

Editor's note: alternate directions below are for a specific applicator container:

[HOW TO APPLY]

1. Apply when air is calm to avoid drift and contact with eyes and skin.
2. Wearing household latex or rubber gloves, unscrew cap and remove seal from bottle top.
3. Tightly screw cap back on bottle and flip top to open.
4. From a distance of 12 inches point bottle towards plant to be treated.
5. Point bottle downward in a 45° angle, and with one hand, apply light constant pressure to the center of the bottle while shaking sideways to produce a light dusting.
6. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Many insects hide on the underside of the leaf.
7. To continue applying, release pressure from bottle before squeezing again.

Directions for Application		
Crop	Target pest (Insects Controlled)	Remarks [Applications Per Season (max)] [Days to Wait to Harvest][as separate columns]
Cole Crops		
Broccoli	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Brussell Sprouts	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Cabbage	Cabbage Looper Imported Cabbageworm	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.

	Diamondback Moth Cabbage Aphid	
Cauliflower	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Chinese Broccoli* *Not for use in California	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Chinese Cabbage (tight-heading varieties only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.
Collards (For use in AR, AZ, GA, IL, NC, OK, SC & TX only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Cucurbit Vegetables		
Cucurbit Vegetables*	Leafminer Squash Bug Cabbage Looper Cucumber Beetle (adult) Cutworm Melon Worm Pickle Worm Plant Bug (including Lygus and Stink Bug) Rindworm Squash Vine Borer	Dust lightly. Apply up to 8 times per season. May be applied until harvest.
*Cucurbit Vegetables are: Chayote(fruit), Chinese Waxgourd, citron melon, cucumber(gherkin), gourds(edible includes hyotan, cucuzza, hechima, Chinese Okra), Momordica spp.(includes balsam apple, balsam pear, bitter melon, Chinese cucumber) muskmelons(includes true cantaloupe, casaba, creshaw, golden porshaw, honeydew melons, honey bells, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, pumpkin, squash (summer such as crookneck, scallop, straightneck, vegetable marrow, zucchini), squash(winter such as butternut, calabaze, hubbard, acorn, spaghetti), watermelon(includes hybrids).		
Fruiting Vegetables		
Eggplant	Colorado Potato Beetle Vegetable Leafminer Cabbage Looper Flea Beetle	Dust lightly. Apply up to 10 times per season. May be applied until 3 days before harvest.
Peppers	Pepper Weevil Vegetable Leafminer Cabbage Looper Flea Beetle European Corn Borer	Dust lightly. Apply up to 8 times per season. May be applied until 3 days before harvest.

Tomatoes	Tomato Fruitworm Cabbage Looper Tomato Pinworm Vegetable Leafminer Beet Armyworm Colorado potato Beetle Granulate Cutworm Hornworm	Dust lightly. Apply up to 6 times per season. May be applied until day of harvest.
Leafy Vegetables		
Leafy Vegetables* (Except brassica)	Vegetable Leafminer Cabbage Looper Cutworm Complex Fall Armyworm Beet Armyworm Corn Earworm Leafhopper European Corn Borer	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.
*Leafy vegetables are: Amaranth, arugula, cardoon, celery, chinese celery, celtuce, chervit, corn salad, chrysanthemum, edible-leaved chrysanthemum, garland, cress(garden, yellow rocket and winter), dandelion, dock(sorrel), endive, fennel, lettuce (head and leaf), orach, parsley, purslane(winter and garden), radicchio, rhubarb, spinach, spinach vine(Malabar spinach and Indian spinach), spinach(New Zealand), Swiss Chard.		
Root, Tuber, and Bulb Vegetables		
Onions (Dray bulb and Garlic)	Onion Thrips Armyworm Cutworm Leafminer Onion Maggot(Adult) Stink Bug	Dust lightly. Apply up to 6 times per season. May be applied until 1 day before harvest.
Potatoes	Colorado Potato Beetle Potato Leafhopper Aster Leafhopper Potato Flea Beetle Potato Aphid Potato Tuberworm Potato Psyllid Cabbage Looper	Dust lightly. Apply up to 8 times per season. May be applied until 14 days before harvest.
Turnips (FL, GA, IL, IN, OK, SC, TX, WA only) (TX, OK, IL, IN only) (SC, GA, FL, WA only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. May be applied until 1 day before harvest. Apply up to 8 times per season Apply up to 4 times per season
Cereal Grains		
Sweet Corn (FL only)	Corn Earworm European Corn Borer Fall Armyworm Southern Armyworm Corn Rootworm(Adult) Cutworm Aster leafhopper Flea Beetle	Dust lightly. Apply up to 6 times per season. May be applied until 1 day before harvest. Apply up to 8 times per season
Other Crops		

Artichokes	Artichoke Plume Moth	Dust lightly. Apply up to 5 times per season. May be applied until day of harvest.
Asparagus	Cutworm Complex Asparagus Beetle	Dust lightly. Apply up to 4 times per season. May be applied until 1 day before harvest.
Ornamentals		
Asparagus Fern	Japanese Beetle Asparagus Beetle Lygus Bug	Dust lightly. Apply to the fern stage of asparagus plant after spear harvest when adult stage of the Asparagus beetle, Lygus bugs and the adult stage of the Japanese Beetle are present.
Roses	Beet Armyworm Cabbage Looper Omnivorous Leafroller	Dust lightly.
Tree Crops		
Dust lightly, using a dust gun or rotary duster [an appropriate duster as available].		
Almonds	Navel Orangeworm Peach Twig Borer	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.
Apples	Apple Aphid Redbanded Leafroller Obliquebanded Leafhopper Plum Curculio White Apple Leafhopper Spotted Tentiform Leafminer Tarnished Plant Bug	Dust lightly. Do not apply after petal fall.
Avocados	Thrips Leafhopper Mirid Bug Lepidopterous Larvae	Dust lightly. Apply up to 6 times per season. May be applied until 7 days before harvest.
Cherries	Lesser Peachtree Borer Rose Chafer Green Fruitworm Redbanded Leafroller Plum Curculio Tarnished Plant Bug	Dust lightly. Apply when insects appear. In the East: do not exceed 6 applications per season, with no more than 4 applications after petal fall. In the West: do not apply more than 4 applications per season, with no more than 3 applications after petal fall.
Filberts	Filbertworm Obliquebanded Leafroller	Dust lightly. Apply up to 4 times per season. May be applied until 14 days before harvest.
Papaya (For use in Florida only)	Fruit Fly	Dust lightly. Apply up to 6 times per season. May be applied until 7 days before harvest.
Peaches	Peach Twig Borer Oriental Fruit Moth Green Fruitworm Lesser Peachtree Borer Plum Curculio Tarnished Plant Bug Rose Chafer	Dust lightly. Apply up to 5 times per season. May be applied until 14 days before harvest.
Pears (Dormant)	Pear Psylla	Dust lightly. Apply up to 2 times per season. May be applied during the dormant season through the delayed dormant growth periods only.
Pears (Pre-Bloom)	Pear Psylla Codling Moth Green Fruitworm	Dust lightly. Apply up to 2 times per season. Maybe applied from dormant through bud burst.
Pistachios	Peach Twig Borer Navel Orangeworm	Dust lightly. Apply up to 2 times per season. May be applied until 10% hull split.
Walnuts	Navel Orangeworm Codling Moth	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.

12
13

	Walnut Husk Fly	
--	-----------------	--

[Editor's Notes: Alternate Chart for Vegetables/Melons and Fruit/Nut Trees and Insects Controlled narrative]

VEGETABLES & MELONS [Crop - Dust lightly]	Applications Per Season (Max)	Days to Wait To Harvest [Pre-harvest Interval]
[Cole Crops-] Broccoli, Brussels sprouts & cauliflower	8	1 [day]
[Cole Crops-] Cabbage & Chinese cabbage (tight-heading varieties only)	5	1 [day]
[Cucurbit Vegetables including] Cucumbers, gourds, [honeydew] melons, [pumpkins, cantaloupe, zucchini,] squash [and watermelon (including hybrids)]	8	0 [May be applied until harvest]
Eggplant	10	3 [days]
Peppers (bell)[including green peppers]	8	3 [days]
Tomatoes	6	0 [May be applied until harvest]
[Leafy Vegetables (except brassica) including] Amaranth, arugula celery, lettuce (head and leaf), parsley, radicchio, rhubarb, spinach & Swiss chard	5	1 [day]
Onions (dry bulb & garlic)	6	1 [day]
Potatoes	8	14 [days]
Sweet corn	6 (FL only 8) [8 in Florida]	1 [day]
Artichokes	5	0 [May be applied until harvest]
Asparagus	4	1 [day]

FRUIT & NUT TREES [Tree Crops - Dust lightly using a dust gun or rotary duster]	APPLICATIONS PER SEASON (MAX)	DAYS TO WAIT TO HARVEST/COMMENTS [Pre-harvest Interval/Remarks]
Apples		Do not apply after petal fall.
Cherries (in the East)	6	No more than 4 applications after petal fall.
Cherries (in the West)	4	No more than 3 applications after petal fall.
Peaches	5	14 [days]
Pears (Dormant)	2	May be applied during the dormant through the delayed dormant growth periods only.
Pears (Pre-bloom)	2	May be applied from dormant through bud burst.
Walnuts	8	1 [day]

[INSECTS CONTROLLED]

On Fruit & Nut Trees: Apply aphid, caterpillars, codling moth, filbertworm, fruit fly, green fruitworm, lepidopterous larvae, leafhoppers, leafrollers, lesser peachtree borer, mired bug, navel orangeworm, oriental

fruit moth, peach twig borer, pear psylla, plum curculio, redbanded leafroller, rose chafer, spotted tentiform leafminer, tarnished plant bug, thrips, white apple leafhopper and walnut husk fly.

On Vegetables & Melons: Aphids, armyworms, artichoke plume moth, asparagus beetle, Aster leafhopper, cabbage aphid, cabbage looper, Colorado potato beetle, corn earworm, corn rootworm (adult), cutworm complex, cucumber beetle (adult), cutworms, diamondback moth, European corn borer, fall armyworm, flea beetles, granulate cutworm, hornworms, imported cabbageworm, leafhoppers, leafminers, melon worm, onion maggot (adult), onion thrips, pepper weevil, pickle worm, plant bugs (including lygus, squash and stink bugs), potato psyllid, potato aphid, potato flea beetle, potato leafhopper, potato tuberworm, rindworm, squash vine borer, tomato fruitworm and tomato pinworm, vegetable leafminer, southern armyworm.

On Ornamentals (including Roses): Asparagus beetle, beet armyworm, cabbage looper, Japanese beetle, lygus bug and omnivorous leafroller]

Alternate or optional wording below in brackets “[]” to be used before Directions for Use or on front panel.

- [100% Satisfaction {company logo} Guaranteed]
- [Shaker Top Applicator]
- [This product delivers effective long lasting control of Aphids, Armyworms, Beetles, Caterpillars, Japanese Beetles plus other listed pests.]
- [Protecting your garden from pests has never been easier.]
- [New Easy To Use Applicator]
- [Use on Ornamentals, Fruits & Vegetables]

READ THE LABEL FIRST!

QUICK FACTS	
<ul style="list-style-type: none"> · Kills Over 60 Types of Insects · Effective Long Lasting Control · Use on Fruit Trees, Flowers, Fruits and Vegetables · Use on Dogs and Cats to Kill Fleas, Ticks and Lice 	
KILLS	Aphids, Armyworms, Beetles, Japanese Beetles, Cabbage Worms, Caterpillars, Horn Worms, Fruit Flies, Fleas, Ticks and other listed pests
WHERE TO USE	Use on Home Garden Fruits, Roses, Flowers and Vegetables
(telephone, computer icons)	For questions or comments, call toll-free (specific customer number) or visit us on-line at (specific customer website)

PRODUCT FACTS	
WHERE TO USE	On ornamentals including roses, vegetables, melons, fruit and nut trees
INSECTS CONTROLLED	Aphids, beetles, caterpillars, cutworms, leafhoppers, and other listed garden insects
WHEN TO USE	Use at first signs of insect infestation, preferably before damage from feeding occurs.

[Product Usage: Home Gardens – Kills – 60+ Insects including Apple Aphids, Armyworms, Cabbage Loopers, Colorado Potato Beetles, Cutworms, Leafminers, Japanese Beetles and other insects as listed.

Use On – Vegetables, Fruit and Ornamental Plants

Usage – Dust Lightly

Pet Use – Kills – Fleas, Ticks, Lice

Use On – Dogs and Cats

Usage – Dust Animals Thoroughly]

12
13

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: If empty – Nonrefillable container. Do not reuse or refill this container.

[When packaged in plastic containers] Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[When packaged in paper and plastic bags] Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

STORAGE AND DISPOSAL

[final printed labeling for residential use only]

PESTICIDE STORAGE: Store in cool, dry place. Do not contaminate feed or foodstuffs by storage or disposal.

PESTICIDE DISPOSAL: If partly filled – Call your local solid waste agency for disposal instructions.

Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: If empty – Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

[NOTICE: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.] [If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.]

[Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.]

13
13

Option Marketing Statements/Graphics that may be featured on basic and/or supplemental registration labeling:

Graphic Images of listed insect pests (e.g., beetle, caterpillar, aphid, etc.), applicator packaging and/or listed sites (e.g., flower, tomatoes, etc.)

As applicable to booklet label/packaging:

OPEN Resealable Label for Use Directions & Precautions

Español ver página 6

Questions, Comments, Medical Information

1-800-225-2883 www.ortho.com

Specially formulated for residential use

Made in USA

Illustration of use of applicator packaging; e.g.:

The Ortho Guarantee: If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund price.