

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDE PROGRAMS
REGISTRATION DIVISION (75-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.
28293-155

DATE OF ISSUANCE
AUG - 2 1991

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

TERM OF ISSUANCE
Until Reregistration

NAME OF PESTICIDE PRODUCT
Unicorn Crawling Insect Spray E.C.

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

UNICORN LABORATORIES
10211 21st Avenue North
St. Petersburg, FL 33716

161 / 5399163
18 /

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Comments on this registration should be directed to the Registration Division, Office of Pesticide Programs, U.S. Environmental Protection Agency, Washington, DC 20460.

The registrant is advised that the registration of this pesticide is subject to the provisions of the Act, including the requirement that the registrant maintain a file of all data submitted to the Agency in support of the registration and make such data available to the Agency upon request.

The registrant is advised that the registration of this pesticide is subject to the provisions of the Act, including the requirement that the registrant maintain a file of all data submitted to the Agency in support of the registration and make such data available to the Agency upon request.

Comments on this registration should be directed to the Registration Division, Office of Pesticide Programs, U.S. Environmental Protection Agency, Washington, DC 20460.

Change the "UNICORN" heading to "Insect Control."

Change the "UNICORN" heading to "Insect Control."

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE
6/1/91

d. Add "feed" to the limitation regarding use in food handling establishments, i.e.

- . . . in food/feed areas . . .
- . . . food/feed handling establishments.
- . . . food/feed is commercially prepared.
- . . . Non food/feed areas are . . .
- . . . for use in non-food/feed areas indoors . . .

e. Under directions add, "Do not apply to classrooms when in use."

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LARocca
 Product Manager (13)
 Insecticide-Rodenticide Branch
 Registration Division (H7505C)

Enclosures

377

UNICORN

CRAWLING INSECT SPAY E.C.

**An Insecticide Offering Residual Kill of Crawling Insects for
Up to 4 Weeks**

RECOMMENDED FOR USE BY COMMERCIAL APPLICATORS

ACTIVE INGREDIENT:

*Permethrin (3-phenoxyphenyl)methyl (+) Cis, trans-
3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate.....13.30% w/w

INERT INGREDIENTS:.....86.70%

100.00% w/w

*Cis/trans isomers ration: min. 35% (+) cis and max. 65% (+) trans

KEEP OUT OF REACH OF CHILDREN

WARNING

**See Side Panel for Additional Precautionary Statements and
Statement of Practical Treatment**

MANUFACTURED FOR:

Unicorn Laboratories
1002 118th Avenue North
St. Petersburg, FL 33716

EPA REG. NO. 28293-

EPA EST. NO. 28293-FL-1

In EPA Letter Dated:

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
amended for the pesticide,
registered under EPA Reg. No.

28293-155

487

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye irritation. Do not get in eyes. Avoid contact with skin or clothing. Harmful if swallowed. Avoid breathing vapor. Wash thoroughly after handling.

For space spraying indoors wear a NIOSH or MSHA approved respiratory protection device.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Do not induce vomiting. This product contains aromatic petroleum solvent. Aspiration may be a hazard.

IF IN EYES: Flush with plenty of water. Get medical attention.

IF ON SKIN: Wash with soap and plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove person to fresh air. Apply artificial respiration if indicated.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not contaminate water by disposing of equipment washwaters. Keep out of lakes, streams or ponds, tidal marshes or estuaries. Apply this product only as specific on this label.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame. Flashpoint minimum of 105°F (41°C).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT USE THIS PRODUCT IN OR ON ELECTRICAL EQUIPMENT DUE TO THE POSSIBILITY OF SHOCK HAZARD.

Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where food may be prepared or held.

Non-food areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

For use in non-food areas indoors and outside premises of hotels, motels, homes, preschools, nurseries, day camps, greenhouses, lathouses, seed houses, food handling establishments, packaging and processing plants, commercial, industrial, office, institutional and storage buildings, dairies, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, and hospitals. **In Hospitals:** Patients should be removed prior to treatment. Also for use in campers, mobile homes, buses, automobiles, trucks, and boxcars. Also for use in dog kennels and horse stables. **In Horse Stables:** Not for application if horses will be used for food.

One gallon of finished spray is sufficient to treat 1580 linear feet (482 m) with a 6 inch (15.2 cm) spray band.

For contact and residual sprays, mix with water in the following manner:

<u>Finished Spray</u>	<u>(Fluid Ounces)</u>	<u>Water</u>
0.25%	2-2/3 (79 ml)	1 gallon (3.79 L)
0.25%	2/3 (20 ml)	1 quart (0.95 L)
0.50%	5-1/2 (163 ml)	1 gallon (3.79 L)
0.50%	1-1/3 (39 ml)	1 quart (0.95 L)

For heavy infestations, use the 0.50% dilution. For lighter infestations and for maintenance purposes, use the 0.25% dilution.

PEST INDOORS: For residual and contact control of cockroaches (waterbugs), crickets, silverfish (firebrats), clover mites, millipedes, centipedes, earwigs, sowbugs (pillbugs), spiders, brown dog ticks, lone star ticks, ants, fire ants, clothes moths, carpet beetles, fleas, boxelder bugs, scorpions, spider beetles, bed bugs, grasshoppers, house flies, mosquitoes, gnats, fruit flies, cluster flies, flying moths, confused flour beetles, saw-toothed grain beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible.

For Bedbugs: Apply as a spot treatment to cracks and crevices, around baseboards, floorboards, bedboards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor bedbugs, concentrating on tufts, folds and seams. Repeat as necessary. Allow all treated articles to dry thoroughly before use.

Grain Insects: To control confused flour beetles, saw-toothed grain beetles, spider beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths, carpet beetles: Remove infested grain, cereal or flour and destroy. Avoid contamination of other food and foodstuffs in the areas by removing before spraying. Spray directly on exposed stages and into cracks and crevices where grain insects may be hiding. In homes, dispose of all infested products. The shelves and drawers should be cleaned and thoroughly sprayed so as to contact all pests, and covered with shelf paper after allowing shelves to dry. Any utensils that are contacted with spray should be washed before using. Repeat as necessary.

Spiders: Spray directly on these pests and use repeated applications to the webbing and all cracks and crevices which harbor spiders.

Ants: To kill ants, spray trails, nests and points of entry within buildings and homes. Spray directly on ants where possible. Repeat as necessary.

Wasps, Hornets, Yellow Jacks, Bees: Application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible. Repeat as necessary.

Fleas, Brown Dog and Lone Star Ticks on Dogs, Cats and Premises: Spray animals along back, under forelegs and on other areas, except scrotum, face and anus. Part hair to contact skin and insure good coverage. Hold sprayer about one foot (30 cm) away while spraying. Apply until hair is thoroughly moist. Do not spray directly in eyes, mouth or nose of animal. To prevent reinfestation, thoroughly spray the animals' bedding and other resting areas until the surface is slightly moist contacting as many insects as possible. Treat the floor areas, cracks and crevices in the floor and along baseboards at the rate of 2 seconds per linear foot (30 cm) at a distance of 12 to 18 inches (30-45 cm). Also spray the entire inside surface of dog houses.

House Flies, Mosquitoes, Cluster Flies, Gnats, Fruit Flies, Flying Moths: Close off all doors and windows. Point sprayer upward. Direct spray mist to all parts of the room, especially windows and other light sources which may attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Do not remain in treated room. Ventilate the room when treatment is completed.

Carpet Beetles and Clothes Moths: To treat carpets hold container 14-18 inches (36-46 cm) from carpet. Apply spray evenly, covering the entire surface until slightly moist. Spray underside of carpet when possible. Avoid wetting or soaking. **To Treat Garments, Blankets, Carpets and Other Fabrics Before Storage:** Dry clean or wash before treatment and after storage. Apply spray evenly, covering the material on both sides until slightly moist. For best results, place treated materials in plastic bags after drying, and store in a dark place. Also spray floor and baseboards, closets, cracks, crevices, infested areas of shelving and/or wherever these insects are seen or suspected.

Pests on Outside Surfaces of Buildings: Use this product to control ants, clover mites, crickets, earwigs, millipedes and sowbugs (pillbugs) on the outside surfaces of buildings. To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet (0.5 to 1 m), where pests are active and may find entrance. Also apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps, and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

House Flies, Mosquitoes, Cluster Flies, Gnats, Flying Moths Outdoors: For outdoor use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, and other places where these insects alight or congregate. Repeat as necessary.

Ants, Lone Star Ticks, Crickets, Fleas, and Earwigs Outdoors: For treatment of localized infestations of these insects in outdoor areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as infestations warrant and as reinfestations occur.

TO CONTROL FIRE ANTS: Dilute 2 tablespoons (30 ml) of this product in 1 gallon (3.79 L) of water. Apply 1 gallon (3.79 L) of solution as gentle rain to each fire ant mound using a sprinkler can. Thoroughly wet the mound and surrounding area to a 4 foot diameter (1.22 m). For best results, apply in cool weather, 65-80°F (18.5-27°C), or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity, and treat all colonies which may not as yet have constructed a mound. Larger quantities of solution can be prepared for multiple mound treatments. Equipment to deliver the diluted product as a gentle rain at the rate of 1 gallon (3.79 L) per mound is still required. Do not use equipment that produces a pressurized spray as pressurized sprays may disturb fire ants and cause migration, reducing product effectiveness.

WARRANTY STATEMENT

Our recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (75-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.

2003-115

DATE OF ISSUANCE

AUG - 2 1991

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

TERM OF ISSUANCE

NAME OF PESTICIDE PRODUCT

UNICORN Crawling Insect Killer, etc.

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Unicorn Laboratories
1002 Fifth Avenue North
St. Petersburg, FL 33710

161 / 5399163
18 /

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered, reregistered, or suspended under FIFRA section 3(c)(2)(A) provided that you:

1. Do not use any other material required to register or reregister your product under FIFRA section 3(c)(2)(B) or FIFRA section 4, and do not use any other material required to register or reregister your product under FIFRA section 4.

2. Use the following changes listed below before you release the product for shipment:

- Use the phrase "EPA Registration No. 2003-115."
- Change the "Pest Indoors" heading to "Indoor Pest Control."
- Insert the "Outdoor Pest Control" heading before the paragraph that begins "Pests on Outside Surfaces."

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

8/1/91

d. Add "feed" to the limitation regarding use in food handling establishments, i.e.

- . . . in food/feed areas . . .
- . . . food/feed handling establishments.
- . . . food/feed is commercially prepared.
- . . . Non food/feed areas are . . .
- . . . for use in non-food/feed areas indoors . . .

e. Under directions add, "Do not apply to classrooms when in use."

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (13)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosures

RECEIVED
MAY 1 1964
U.S. DEPARTMENT OF AGRICULTURE
WASHINGTON, D.C.

UNICORN

CRAWLING INSECT SPAY E.C.

An Insecticide Offering Residual Kill of Crawling Insects for
Up to 4 Weeks

RECOMMENDED FOR USE BY COMMERCIAL APPLICATORS

ACTIVE INGREDIENT:

*Permethrin (3-phenoxyphenyl)methyl (±) Cis, trans-
3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate.....13.30% w/w

INERT INGREDIENTS:.....86.70%
100.00% w/w

*Cis/trans isomers ration: min. 35% (±) cis and max. 65% (±) trans


KEEP OUT OF REACH OF CHILDREN

WARNING

See Side Panel for Additional Precautionary Statements and
Statement of Practical Treatment

MANUFACTURED FOR:

Unicorn Laboratories
1002 118th Avenue North
St. Petersburg, FL 33716



EPA REG. NO. 28293-

EPA EST. NO. 28293-FL-1

with 2000000000
in EPA Letter Dated:

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide
registered under EPA Reg. No.

28293-155

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye irritation. Do not get in eyes. Avoid contact with skin or clothing. Harmful if swallowed. Avoid breathing vapor. Wash thoroughly after handling.

For space spraying indoors wear a NIOSH or MSHA approved respiratory protection device.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Do not induce vomiting. This product contains aromatic petroleum solvent. Aspiration may be a hazard.

IF IN EYES: Flush with plenty of water. Get medical attention.

IF ON SKIN: Wash with soap and plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove person to fresh air. Apply artificial respiration if indicated.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not contaminate water by disposing of equipment washwaters. Keep out of lakes, streams or ponds, tidal marshes or estuaries. Apply this product only as specific on this label.

PHYSICAL AND CHEMICAL HAZARDS


Do not use or store near heat or open flame. Flashpoint minimum of 105°F (41°C).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT USE THIS PRODUCT IN OR ON ELECTRICAL EQUIPMENT DUE TO THE POSSIBILITY OF SHOCK HAZARD.

Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where food may be prepared or held.



Non-food areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

For use in non-food areas indoors and outside premises of hotels, motels, homes, preschools, nurseries, day camps, greenhouses, lathhouses, seed houses, food handling establishments, packaging and processing plants, commercial, industrial, office, institutional and storage buildings, dairies, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, and hospitals. **In Hospitals:** Patients should be removed prior to treatment. Also for use in campers, mobile homes, buses, automobiles, trucks, and trailers. Also for use in dog kennels and horse stables. **In Horse Stables:** Not for application if horses will be used for food.

One gallon of finished spray is sufficient to treat 1580 square feet (482 m) with a 6 inch (15.2 cm) spray band.

For contact and residual sprays, mix with water in the following manner:

<u>Finished Spray</u>	<u>(Fluid Ounces)</u>	<u>Water</u>
0.25%	2-2/3 (79 ml)	1 gallon (3.79 L)
0.25%	2/3 (20 ml)	1 quart (0.95 L)
0.50%	5-1/2 (163 ml)	1 gallon (3.79 L)
0.50%	1-1/3 (39 ml)	1 quart (0.95 L)

For heavy infestations, use the 0.50% dilution. For lighter infestations and for maintenance purposes, use the 0.25% dilution.

PEST INDOORS: For residual and contact control of cockroaches (waterbugs), crickets, silverfish (firebrats), clover mites, millipedes, centipedes, earwigs, sowbugs (pillbugs), spiders, brown dog ticks, lone star ticks, ants, fire ants, clothes moths, carpet beetles, fleas, boxelder bugs, scorpions, spider beetles, bed bugs, grasshoppers, house flies, mosquitoes, gnats, fruit flies, cluster flies, flying moths, confused flour beetles, saw-toothed grain beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible.

For Bedbugs: Apply as a spot treatment to cracks and crevices, around baseboards, floorboards, bedboards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor bedbugs, concentrating on tufts, folds and seams. Repeat as necessary. Allow all treated articles to dry thoroughly before use.

Grain Insects: To control confused flour beetles, saw-toothed grain beetles, spider beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths, carpet beetles: Remove infested grain, cereal or flour and destroy. Avoid contamination of other food and foodstuffs in the areas by removing before spraying. Spray directly on exposed stages and into cracks and crevices where grain insects may be hiding. In homes, dispose of all infested products. The shelves and drawers should be cleaned and thoroughly sprayed so as to contact all pests, and covered with shelf paper after allowing shelves to dry. Any utensils that are contacted with spray should be washed before using. Repeat as necessary.

Spiders: Spray directly on these pests and use repeated applications to the webbing and all cracks and crevices which harbor spiders.

Ants: To kill ants, spray trails, nests and points of entry within buildings and homes. Spray directly on ants where possible. Repeat as necessary.

Wasps, Hornets, Yellow Jacks, Bees: Application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible. Repeat as necessary.

Fleas, Brown Dog and Lone Star Ticks on Dogs, Cats and Premises: Spray animals along back, under forelegs and on other areas, except scrotum, face and anus. Part hair to contact skin and insure good coverage. Hold sprayer about one foot (30 cm) away while sprayer. Apply until hair is thoroughly moist. Do not spray directly in eyes, mouth or nose of animal. To prevent reinfestation, thoroughly spray the animals' bedding and other resting areas until the surface is slightly moist contacting as many insects as possible. Treat the floor areas, cracks and crevices in the floor and along baseboards at the rate of 2 seconds per linear foot (30 cm) at a distance of 12 to 18 inches (30-45 cm). Also spray the entire inside surface of dog houses.

House Flies, Mosquitoes, Cluster Flies, Gnats, Fruit Flies, Flying Moths: Close off all doors and windows. Point sprayer upward. Direct spray mist to all parts of the room, especially windows and other light sources which may attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Do not remain in treated room. Ventilate the room when treatment is completed.

Carpet Beetles and Clothes Moths: To treat carpets hold container 14-18 inches (36-46 cm) from carpet. Apply spray evenly, covering the entire surface until slightly moist. Spray underside of carpet when possible. Avoid wetting or soaking. **To Treat Garments, Blankets, Carpets and Other Fabrics Before Storage:** Dry clean or wash before treatment and after storage. Apply spray evenly, covering the material on both sides until slightly moist. For best results, place treated materials in plastic bags after drying, and store in a dark place. Also spray floor and baseboards, closets, cracks, crevices, infested areas of shelving and/or wherever these insects are seen or suspected.

Pests on Outside Surfaces of Buildings: Use this product to control ants, clover mites, crickets, earwigs, millipedes and sowbugs (pillbugs) on the outside surfaces of buildings. To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet (0.5 to 1 m), where pests are active and may find entrance. Also apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps, and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

House Flies, Mosquitoes, Cluster Flies, Gnats, Flying Moths Outdoors: For outdoor use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, and other places where these insects alight or congregate. Repeat as necessary.

Ants, Lone Star Ticks, Crickets, Fleas, and Earwigs Outdoors: For treatment of localized infestations of these insects in outdoor areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as infestations warrant and as reinfestations occur.

TO CONTROL FIRE ANTS: Dilute 2 tablespoons (30 ml) of this product in 1 gallon (3.79 L) of water. Apply 1 gallon (3.79 L) of solution as gentle rain to each fire ant mound using a sprinkler can. Thoroughly wet the mound and surrounding area to a 4 foot diameter (1.22 m). For best results, apply in cool weather, 65-80°F (18.5-27°C), or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity, and treat all colonies which may not as yet have constructed a mound. Larger quantities of solution can be prepared for multiple mound treatments. Equipment to deliver the diluted product as a gentle rain at the rate of 1 gallon (3.79 L) per mound is still required. Do not use equipment that produces a pressurized spray as pressurized sprays may disturb fire ants and cause migration, reducing product effectiveness.

WARRANTY STATEMENT

Our recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

