

# UNICORN GENERAL PURPOSE INSECTICIDE

- Clear - Stable -  
Leaves No Oily or Objectionable Residues

Effective, Fast Acting Insecticide

For Use in Homes, Institutions, Restaurants, Food Processing Plants, Dairies, Livestock  
and Outdoors for Mosquito Control.

For Control of Fleas and Brown Dog Ticks on Premises and on Pets

Apply with Conventional Mechanical or Compressed Air Equipment

**ACTIVE INGREDIENTS:**

Pyrethrins.....	0.1%
Piperonyl Butoxide, Technical*.....	1.0%


<b>INERT INGREDIENTS:</b> .....	<u>98.9%</u>
<b>Total:</b>	100.0%

\* Equivalent to min. 0.8% (butylcarbityl) (6-propylpiperonyl)  
ether and 0.2% related compounds.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See Side/Back Panel for Additional Precautionary Statements  
DO NOT TRANSPORT OR STORE UNDER 32° F.



CONTENTS:

**MANUFACTURED BY:**  
Unicorn Laboratories  
12385 Automobile Blvd.  
Clearwater, Florida 33762  
(727) 578-4545

- READ ENTIRE LABEL BEFORE EACH USE.
- FOR PET USE - USE ONLY ON DOGS OR CATS.

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets, birds and cover fish aquaria before space spraying or surface application. Sensitivities may occur after using **ANY** pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

**Environmental Hazards**

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

<b>FIRST AID</b>	
<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If Inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If On Skin or Clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If In Eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**INDOOR APPLICATION**

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All

food processing surfaces should be covered during treatment or thoroughly cleaned with potable water before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

**Surface Spraying:** To control cockroaches, palmetto bugs, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites and cadelles. Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish spray bookcases. For ants spray trails, nests and points of entry. If surface application only is to be used spray floors and other surfaces applying a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles, spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control bedbugs, spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks, thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed.

To control clothes moths, remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage, directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

**SPACE SPRAYING:** Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths: Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3 - 6 ounces per 1000 cubic feet of space.

**FOGGING:** To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

**ON LIVESTOCK:** To protect cattle and horses from horn flies, mosquitoes, and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies and deer

flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection. To control blood-sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

**ON PETS:** Start spraying at the tail, moving the dispenser rapidly and making sure that the animals entire body is covered including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wet thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Reapply every week as necessary. Do not use on kittens or puppies under twelve (12) weeks of age. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication.

**OUTDOOR APPLICATION**

**ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, (low marshy places, stagnant pools or ponds and shore lines). Apply at a rate of 25 pints per acre (0.1 ounce per 1000 square feet). Using truck-mounted conventional misting equipment such as Rotomist® models 51C, 91 or 100 HT traveling at 5 m.p.h. to treat a 300 foot swath calibrate equipment to deliver 0.9 gallon per minute.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE AND SPILL PROCEDURES:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

**PESTICIDE DISPOSAL:** Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

**CONTAINER DISPOSAL: OVER ONE GALLON** -Triple rinse (or equivalent) then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

**CONTAINERS 1 GALLON AND UNDER:** Do not reuse empty container. Rinse thoroughly, securely wrap in several layers of newspaper and discard in trash.

**Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.**