

UNICORN GENERAL PURPOSE AQUEOUS INSECTICIDE II

- ◆ For Domestic Use both Indoors and Outdoors
- ◆ A Ready-To-Use Stable Emulsion Insecticide
- ◆ Provides 180 Day (6 Months) (non-porous surfaces) Control of Roaches, including the German and American Cockroach.
 - ◆ For Control of Fleas, Brown Dog Ticks and Lice on Premises and on Dogs
- ◆ For Control of Ticks which may carry Lyme disease.
- ◆ Controls Fleas and Ticks on Dogs for 35 Days. Aids in Control of Ticks on Dogs for an Additional 2 Weeks.
 - ◆ For Control of Pests on Livestock
- ◆ Water Base ◆ READY TO USE

Use Indoors - Spray hiding places, cracks and crevices.

Use Outdoors- Treat around windows, doors, porches, screens, patios and garages.

Product Usage: Kills nuisance insects (as listed on the label) in and around the home, fleas and ticks on dogs.

Use on: In cracks and crevices where insects hide, around windows and doors, where insects enter the home.

Amount to Use: Ready to use, no mixing required.

ACTIVE INGREDIENTS:

Permethrin* -----0.50%

INERT INGREDIENTS:-----99.50%

Total: 100.00%

*Cis/trans ratio: Min. 35% (±) cis, Max. 65% (±) trans

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side/Back Panel for Additional Precautionary Statements.

ACCEPTED
NOV 26 2004
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. 28293-179

NET CONTENTS:

MANUFACTURED BY:
 Unicorn Laboratories
 12385 Automobile Blvd.
 Clearwater, Florida 33762
 (727) 578-4545

- READ ENTIRE LABEL BEFORE EACH USE.
- FOR PET USE - USE ONLY ON DOGS.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications.

Environmental Hazards

This product is extremely toxic to fish and other aquatic organisms. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Affix trigger pump or other type spray head attachment to container for application. Do not transfer to another container.

SHAKE WELL BEFORE USING.

INDOOR APPLICATION:

For spot application only. Do not use as a space spray indoors. Do not apply to classrooms while in use. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed.

SURFACE SPRAYING: To Control Cockroaches (180 day/6 month control) (non-porous surfaces), Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Darkling Beetles, Meal Worms, Grain Mites and Cadelles:

Direct the spray into hiding places, cracks and crevices, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

To Control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas.

To Control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Allow all treated articles to dry thoroughly before use.

To Control Fleas (adults and larvae) and Ticks, including Ticks which may carry Lyme disease (adults and larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Allow to dry thoroughly before allowing pets to contact treated areas.

To Control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages, directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To Kill Lice and Louse Eggs: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

ON LIVESTOCK: To Protect Cattle (beef and dairy), Goats, Sheep, Hogs and Horses from horn flies, face flies, house flies, mosquitoes and gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To Control Stable Flies, Horse Flies and Deer Flies: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To Control Blood-sucking Lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To Control Poultry Lice: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To Control Bedbugs and Mites in Poultry Houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS: To Control Fleas, Ticks and Lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes and face. Avoid contact with genitalia. Repeat every two (2) weeks as necessary. Sensitivities may occur after using **ANY** pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Do not use on dogs under twelve (12) weeks of age. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication.

OUTDOOR APPLICATION

To Control Ticks, Including Ticks Which May Carry Lyme Disease (Perimeter Treatment):

Use this product as a residual treatment to control Ticks including the Deer Tick (*Ixodes dammini*), Lone Star Tick (*Amblyomma americanum*), Brown Dog Tick (*Rhipicephalus sanguineus*), American Dog Tick (*Dermacentor variabilis*) and Western Black-legged Tick (*Ixodes pacificus*).

For each area listed below, apply until surface is wet (approximately 1 gal/750 - 1000 ft²).

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces where these pests may enter.

Treat a band of vegetation and leaf litter 6 to 10 feet wide around the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6 - 10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

FOR CONTROL OF ORNAMENTAL GARDEN INSECTS:

Not for use on plants being grown for sale or other commercial or research purposes. Not for use in commercial greenhouses, nurseries, on sod farms or grass grown for seed.

For use on Roses, Dahlias, Asters and other Ornamentals to kill Japanese beetles hit by spray as well as whiteflies, aphids, exposed thrips, spiders, red mites and leafminers. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against whiteflies, aphids, spider and red mites, exposed thrips, lace bugs, Japanese beetles, pavement ants, clover mites and armyworms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area away from heat or open flame. Store above 32°F.

DISPOSAL: If empty – Do not reuse this container. Place in trash or offer for recycling if available. If partly filled – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.