

**UNICORN BACKUP
POUR-ON INSECTICIDE**

**Pour-On for Beef, Lactating and Non-Lactating
Dairy Cattle and Sheep**

Controls Lice and Flies on Cattle; Keds on Sheep

ACTIVE INGREDIENT

Permethrin (3-phenoxyphenyl) methyl
(±) cis, trans-3-(2,2-dichloroethenyl)
2,2-dimethylcyclopropanecarboxylate*..... 1.0%

INERT INGREDIENTS**..... 99.0%

100.0%

* Cis/trans ratio: Min. 35% (±) cis
Max. 65% (±) trans

** Contains petroleum distillates

Keep Out of Reach of Children

CAUTION

See Side Panel for Additional Precautionary Statements


NET WEIGHT: ONE U.S. GALLON

Manufactured by:

Unicorn Laboratories
1000 118th Avenue North
St. Petersburg, Florida 33716

EPA Reg. No. 28293-182(E:11/12/92)

EPA Est. No. 28293-FL-1



28293-182

2 8 2

))

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION: Avoid contact with eyes.

Statement of Practical Treatment

IF IN EYES: Immediately flush eyes with plenty of water. Get medical attention if discomfort persists.

IF SWALLOWED: Call a physician immediately. Do not induce vomiting unless advised by a physician.

NOTE TO PHYSICIAN: Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally.

Environmental Hazards

This pesticide is extremely toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not add directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

Physical and Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READY TO USE** - No dilution necessary.

APPLICATION INSTRUCTIONS: Pour along back line and down face. Apply 1/2 fl. oz. per 100 lbs. body weight of animal, up to a maximum of 5 fl. oz. for any one animal to control Lice, Horn Flies and Face Flies. Aids in control of Horse Flies, Stable Flies, Mosquitoes, Black Flies and Ticks on lactating and non-lactating dairy cattle, beef cattle, calves and sheep. For Keds on sheep, pour along back line over neck, shoulders and rump. Apply 1/4 fl. oz. per 50 lbs body weight of animal, up to a maximum of 3 fl. oz. for any one animal.

Repeat treatment as needed, but not more than once every 2 weeks. For optimum lice control, a double treatment at 14 day intervals is recommended.

NOTE: This product is not effective for grub control on cattle.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep container sealed when not in use. Do not store near food or feed.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **Container One Gallon and Under:** Do not reuse empty container. Rinse thoroughly, wrap in several layers of newspaper and place in trash.