	MMENTAL PROTECTION AGENCY	28293-206	DATE OF ISSUANCE
RECHE	OF PESTICIDES PROGRAMS TRATION DIVISION (75-767) IASHINGTON, DC 20480	TERM OF ISSUANCE	JUN 1 8 1991
NOTICE OF P		Conditional NAME OF PESTICIDE PRODUC	
(Under the Federal Insecticide, Fungicide,		Unicorn House	e Plant Spray
	lenticide Act, as amended) OF REGISTRANT (Include ZIP code)		
	corn Laboratories	7	
	2 118th Avenue, North Petersburg, FL 33716	· •	
L		١	
submitted to and ac	labefing formula differing in substance from the period by the Registration Division prior to the above U.S. EPA registration numbers.	to use of the label in commerc	
	rmation furnished by the registrant, the alcide, Fungicide, and Rodenticide Act.	bove named pesticide is hereb	y Registered/Reregistered under
is copy of the label:	ing accepted in connection with this Regi	stration/Reregistration is retu	rned herewith.
health and the envir icide in accordance	o way to be construed as an indorsement on onment, the Administrator, on his motion, with the Act. The acceptance of any nam strued as giving the registrant a right to	, may at any time suspend or c ie in connection with the regis	ancel the registration of a pest- tration of a product under this
This FIFRA sec	s product is conditiona c. 3(c)(7)(A) provided	lly registered in that you:	accordance with
1. reregist; when the submit su	Submit and/or cite all ration of your product a Agency requires all reach data.	under FIFRA sec.	3(c)(5) and $4(a)$
2.	Make the labeling charthe product for shipm	nges listed below ent:	before releasing
	a. Add the phrase, "	EPA Registration	No. 28293-206".
	b. Add the following space spray" se	statement to the ection of the pro	"Indoors as a duct label:
	to treatment or t	thoroughly washed ng compound follo	be covered prior with an wed by a potable
	water rinse prior	r to use.	
ATTACHMENT IS	water rinse prior	next page)	
ATTACHMENT IS	APPLICABLE (1		JUN 1 8 1991

Ŋ

))

c. Revise the sentence under the "To control mosquito larvae" section to read as follows:

Apply as a course wetting spray to the border of shallow waters of wetlands, swamps, and marshes where mosquito larvae are known to breed.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

Richard F. Mountfort
Product Manager 10
Insecticide-Rodenticide Pranch
Registration Division (H7505C)

UNICORN

HOUSE PLANT SPRAY

Kills Gypsy Noth Caterpillars and Japanese Beetles. An Effective Insecticide For Use on Vegetable and Ornamental Pests.

A Concentrate Pyrenone(R) Micro-Emulsion, Clear Stable.

Apply with Conventional Mechanical Compressed Air Equipment or

Non-Aerosol Pump.

ACTIVE INGREDIENTS:

Pyrethrins* *Piperonyl Butoxide, Technical	0.02% 0.20%	/
INERT INGREDIENTS:	99.78	/
	100.00%	
* Equivalent to min. 0.16% (butylcarbityl) (6-propylical ether and 0.04% related compounds. **COMME OF THE PARTIES OF THE PAR	Seronyl)	~
in EPA Levier 18	al Inserticide.	3
Keep Out of Reach of Children (100 der of children	todernineshing the or the AREG. No der EPA 200	,o.
CAUTION Solicitude of the control of	93	

See Side/Back Panel for Additional Precautionary Statements

NET CONTENTS: 12 OUNCES

MANUFACTURED BY:

Unicorn Laboratories 1002 118th Avenue North St. Petersburg, Florida 33716

EPA REG. NO. 28293-EM

EPA EST. NO. 28293-FL-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists.

Remove pets, birds and cover fish aquaria before spraying.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly into lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

Physical or Chemical Hazards

DO NOT TRANSPORT OR STORE UNDER 32° F. If freezing occurs, thaw and agitate vigorously.

Statement of Practical Treatment

IF SWALLOWED:

Contact a physician or Poison Control Center. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF EYES OR IN SKIN: Flush affected areas with plenty of water.

Contact a physician if irritation persists.

IF INHALED:

(

Remove affected person to fresh air. Apply artificial respiration if indicated.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO OPERATE: SHAKE BEFORE USING. Hold sprayer 18. inches from foliage and spray with circular motion. Direct spray to that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

TO OPERATE: SHAKE BEFORE USING. Turn nozzle to "spray" or "steam" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct the spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage.

BEST AVAILABLE COPY

TO OPERATE: SHAKE BEFORE USING. Remove shipping cap and save. Insert sprayer hose into bottle and tighten securely. Prime sprayer by holding at cap level of bottle and pumping trigger. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 21 inches from foliage or surfaces being treated. Direct spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage. Remove sprayer and replace shipping cap between uses. Spray water through sprayer after each use. Sprayer should be rinsed thoroughly and pumped empty for storage.

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

ŧ(

G

Use on the following crops: Asparagus, Beans, Broccoli, Cauliflower, Cabbage, Brussels Sprouts, Celery, Cranberries, Eggplant, Lettuce, Mustard Greens, Kale, Collards, Turnips, Peppers, Potatoes, Radishes, Spinach and Tomatoes To Control: Asparagus Beetles, Aphids, Leafhoppers, Whiteflies, Mexican Bean Beetles, 12-Spotted Cucumber Beetles, Cabbage Loopers, Cross-Striped Cabbageworms, Diamondback Moth Larvae, Flea Beetles, Harlequin Bugs, Imported Cabbageworms, Stink Bugs, Green Peach Aphids, Leftiers, Fireworms, Blister Beetles, Colorado Potato Beetles and Webworms.

Use on the following ornamentals: African Violets, Azaleas, Begonias, Camellias, Dahlias, Dogwood, Roses, Wander Jew, Asters, Carnations, Chrysanthemums, Geraniums, Gladiolus and Marigolds, To Control: Aphids, Flea Beetles, Leafhoppers and Whiteflies.

To Kill Gypsy Moth Caterpillars and Adults, Adult Japanese Beetles and Eastern and Forest Tent Caterpillars on contact:

On Trees and Shrubs: Treat trees and shrubs as soon as insacts emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.



Indoors as a Space Spray to control House Flies. Fruit Flies.
Mosquitoes. Gnats and Small Flying Moths: Apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed rood and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

Outdoors. Adult Mosquito Control: Apply as a fine line mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at a rate of 5 to 6 quarts per acre (4 ounces per 1000 square feet).

<u>To Control Mosquito Larvae:</u> Apply at the rate of 7.75 gallons per acre or .18 gallons per 1000 square feet. Apply as a coarse wetting spray to the border of or on shallow waters of wetlands, swamps and marshes where mosquito larvae are known to breed.

✓ STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperature. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fillers' earth, etc. Dispose of with chemical waste.

<u>PESTICIDE DISPOSAL</u>: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

<u>CONTAINER DISPOSAL:</u> Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

CONTAINERS ONE GALLON AND SMALLER: Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

Buyer assumes all risks of use, storage or handling or thir material not in strict accordance with directions given herewith.

