

28293-231

OCT 14 1993

Until Reregistration

Unicorn Insecticide Aqueous  
Pressurized Spray

160/5439027  
18/1

Unicorn Laboratories  
1000 118th Avenue North  
St. Petersburg, FL 33716

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the labeling changes listed below before you release the product for shipment:

- a. Add the phrase, "EPA Registration No. "
- b. Under the Hazards to Humans and Domestic Animals, change the statement "Harmful for swallowed" to read "Harmful is absorbed through the skin."

3. Submit five copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

**BEST AVAILABLE COPY**

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca  
Product Manager 13  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

## UNICORN

## INSECTICIDE AQUEOUS PRESSURIZED SPRAY

A Combination of Insecticides Offering  
Quick Knockdown and Residual Kill  
of Crawling Insects for Up to 4 Weeks

Can be used in non-food areas of food handling establishments such as garbage rooms, lavatories, floor drains (to sewers), entries, vestibules and hallways, offices, locker rooms, machine rooms, basements, boiler rooms, garages, utility rooms, mop closets and storage rooms (after canning or bottling).

## ACTIVE INGREDIENTS:

Permethrin* (3-phenoxyphenyl) methyl (±) cis, trans-3-(2,2-dichloroethenyl)-2-2- dimethylcyclopropanecarboxylate.....	0.15% w/w
d-Trans Allethrin.....	0.25%

INERT INGREDIENTS:.....	99.60%
	100.00% w/w

\* Cis/trans isomers ratio: min. 35% (±) cis  
max. 65% (±) trans

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.  
(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

KEEP OUT OF REACH OF CHILDREN

## CAUTION

See Side/Back Panel for Additional Precautionary Statements.

## NET CONTENTS:

Manufactured by:  
Unicorn Laboratories  
1000 118th Avenue North  
St. Petersburg, FL 33716

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

EPA REG. NO. 28293-

OCT 14 1993

EPA EST. NO. 28293-FL-1

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

28293-231

BEST AVAILABLE COPY

4 of 8

06-2

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans and Domestic Animals**

**CAUTION:** Avoid contact with eyes, skin or clothing. Harmful if swallowed. Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held.

In the home, cover all food handling surfaces and cover or remove all food and cooking utensils or wash thoroughly after treatment.

- Do not apply this product in hospital patient rooms while occupied or in any room while occupied by the elderly or infirm.
- Do not apply to classrooms when in use.
- Do not apply to institutions (including libraries, sports facilities, etc.) in the immediate area when occupants are present.

**Statement of Practical Treatment**

**IF IN THE EYES OR ON SKIN:** Flush with plenty of water. Get medical attention if irritation persists.

**Environmental Hazards**

This product is extremely toxic to fish. Do not apply directly to any body of water. Apply this product only as specified on this label.

**Physical and Chemical Hazards**

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54.5°C) may cause bursting. Do not apply this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. Do not use on surfaces that can be harmed or stained by water.

**STORAGE AND DISPOSAL**

**STORAGE:** Do not store near heat or open flame. Keep in locked storage area. Do not store this water-based formulation at temperatures below 32°F (0°C).

**CONTAINER DISPOSAL:** Replace cap and wrap container in several layers of newspaper. Discard in trash.

## DIRECTIONS FOR USE

SHAKE WELL BEFORE EACH USE.

Remove protective cap. Aim spray opening away from person. Push button to spray.

FOR USE IN NON-FOOD AREAS INDOORS AND OUTSIDE PREMISES OF HOTELS, MOTELS, HOMES, PRESCHOOLS, NURSERIES, DAY CAMPS, GREENHOUSES, LATHHOUSES, SEED HOUSES, FOOD HANDLING ESTABLISHMENTS, PACKAGING AND PROCESSING PLANTS, COMMERCIAL, INDUSTRIAL, OFFICE, INSTITUTIONAL AND STORAGE BUILDINGS, DAIRIES, MILK STORAGE ROOMS, EGG PROCESSING PLANTS, SUPERMARKETS, RESTAURANTS, BOTTLING PLANTS, CANNERIES, FLOUR AND FEED MILLS, BAKERIES, AND HOSPITALS. IN HOSPITALS, PATIENTS SHOULD BE REMOVED PRIOR TO TREATMENT. ALSO FOR USE IN CAMPERS, MOBILE HOMES, BUSES, AUTOMOBILES, TRUCKS, AND BOXCARS. ALSO FOR USE IN DOG KENNELS AND HORSE STABLES. IN HORSE STABLES, NOT FOR APPLICATION IF HORSES WILL BE USED FOR FOOD.

### PESTS INDOORS

For residual and contact control of cockroaches (waterbugs), crickets, silverfish (firebrats), clover mites, millipedes, centipedes, earwigs, sowbugs (pillbugs), spiders, ants, clothes moths, carpet beetles, boxelder bugs, scorpions, spider beetles, grasshoppers, house flies, mosquitoes, gnats, fruit flies, cluster flies, flying moths, confused flour beetles, saw-toothed grain beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths - Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible.

**GRAIN INSECTS:** For residual use, to control confused flour beetles, saw-toothed grain beetles, spider beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths, carpet beetles - Avoid contamination of other food and foodstuffs in the areas by removing before spraying. Spray directly on exposed stages and into cracks and crevices where grain insects may be hiding. Dispose of all infested products. The shelves and drawers should be cleaned and thoroughly sprayed so as to contact all pests, and covered with shelf paper after allowing shelves to dry. Any utensils that are contacted with spray should be washed before using. Repeat as necessary.

**SPIDERS:** Spray directly on these pests and use repeated applications to the webbing and all cracks and crevices which harbor spiders.

**ANTS:** To kill ants, spray trails, nests and points of entry within buildings and homes. Spray directly on ants where possible. Repeat as necessary.

**WASPS, HORNETS, YELLOW JACKETS, BEES:** Application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible. Repeat as necessary.

**HOUSE FLIES, MOSQUITOES, CLUSTER FLIES, GNATS, FRUIT FLIES, FLYING MOTHS:** Close off all doors and windows. Point nozzle upward. Direct spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

**CARPET BEETLES AND CLOTHES MOTHS:** To Treat Carpets - Hold container 14 - 18 inches (36-46 cm) from carpet. Apply spray evenly, covering the entire surface until slightly moist. Spray underside of carpet when possible. Avoid wetting or soaking. To Treat Garments, Blankets, Carpets and Other Fabrics Before Storage - Dry clean or wash before treatment and after storage. Apply spray evenly, covering the material on both sides until slightly moist. For best results, place treated materials in plastic bags after drying, and store in a dark place. Also spray floor and baseboards, closets, cracks, crevices, infested areas of shelving and/or wherever these insects are seen or suspected.

**CONTACT APPLICATION ONLY:** (Not for Residual Control of Fleas, Ticks and Bedbugs)

**FLEAS, BROWN DOG, DEER, BEAR, BLACK LEGGED - WHICH VECTOR LYME DISEASE - AND LONE STAR TICKS ON DOGS, CATS AND PREMISES:** Spray animals along back, under forelegs and on other areas, except scrotum and anus. Part hair to contact skin and insure good coverage. Hold nozzle about one foot (30 cm) away while spraying. Apply until hair is thoroughly moist. Do not spray directly in eyes, mouth or nose of animal. To prevent reinfestation, thoroughly spray the animals' bedding and other resting areas until the surface is slightly moist contacting as many insects as possible. Treat the floor areas, cracks and crevices in the floor and along baseboards at the rate of 2 seconds per linear foot (30 cm) at a distance of 12 to 18 inches (30-45 cm). Also spray the entire inside surface of dog houses. Do not use this product on old, sick or debilitated dogs.

**FOR BEDBUGS:** Apply as a spot treatment to cracks and crevices, around baseboards, floorboards, bedboards and walls. First thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor bedbugs, concentrating on tufts, folds and seams. Repeat as necessary. For persistent infestations, consult professional applicator.

**PESTS OUT-OF-DOORS**

**PESTS ON OUTSIDE SURFACES OF BUILDINGS:** Use this product to control ants, clover mites, crickets, earwigs, millipedes and sowbugs (pillbugs) on the outside surfaces of buildings. To help prevent infestation of buildings treat the building foundation to a height of 2 to 3 feet (0.5 to 1 m), where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps, and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

**HOUSE FLIES, MOSQUITOES, CLUSTER FLIES, GNATS, FLYING MOTHS:** For outdoor use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, and other places where these insects alight or congregate. Repeat as necessary.

**ANTS, LONE STAR TICKS, DEER, BEAR, BLACK LEGGED TICKS (VECTOR LYME DISEASE), CRICKETS, FLEAS, AND EARWIGS:** For treatment of localized infestations of these insects in outdoor areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as infestations warrant and as reinfestations occur.

**WASPS, HORNETS, YELLOW JACKETS, BEES:** Application should be made in late evening when insects are at rest. Aim spray at nest openings in ground, bushes, and in cracks and crevices which may harbor nests, saturating nest openings and contacting as many insects as possible.

**FOR USE ON PLANTS**

**CONTACT AND RESIDUAL SPRAY -** Apply spray to plants covering all foliage. Avoid heavy wetting.

**INDOORS:** To protect plants in greenhouses and lathhouses, and for indoor landscaping - Spray directly on insects to kill leafrollers, armyworms, cutworms, cabbage worms on chrysanthemums; fungus gnats on velvet plants. Spray as directed above.

Spray as directed above to control aphids, armyworms, corn earworms, fall cankerworms, inchworms, Japanese beetles, leafminers, leafrollers, and whiteflies on the following plants:

Ageratum	English ivy	Piggy-back plant	.....
Ardisia	Fuchsia	Poinsettia	.....
Aster	Gladiolus	Portulaca	.....
Azalea	Grape ivy	Pothos	.....
Baby's breath	Hypoestes	Prayer Plant	.....
Begonia	Ivy	Purple Passion	.....
Birds-nest fern	Juniper	Rabbits foot fern	.....
Bleeding heart	Lily	Rose	.....
Carnation	Marigold	Schefflera	.....
Coleus	Orchid	Snapdragon	.....
Crown of thorns	Palms	Snake Plant	.....
Cockscomb	Pansy	Statice	.....
Cyclamen	Piperomia	Velvet Plant	.....
Dracaena	Petunia	Verbena	.....
Dumbcane	Philodendron	Zinnia	.....

**BEST AVAILABLE COPY**

If application is to be made indoors, remove infested plants from living areas prior to application. Spray plants in an area not likely to be occupied by children or pets (particularly fish). When leaves dry, plants can be returned to original location.

Use with CAUTION on ferns, palms and fuchsia. As with all plants listed above, especially carnations, chrysanthemums and roses, there exists a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury.

Repeat applications may be necessary to control citrus black flies on non-bearing citrus; adult Japanese beetles on roses.

**OUTDOORS:** Spray as described above in the late afternoon or evening when the temperature ranges from 50°F to 75°F and when there is little or no wind. Spray at the first indication of insect pests. On those plants that will flower during the season, the spray schedules should be completed before blossom formation. Repeat as necessary; use intervals of 4 to 8 days. Application can be made up to the day of collection.

Spray as directed above to control aphids, cicadas, corn earworms, fall cankerworms, inchworms, leafminers, leafrollers, mealybugs, spider mites, exposed thrips and whiteflies on the following plants:

Ageratum	Gold bells	Pea shrub
Aster	Hypoestes	Petunia
Azalea	Ivy	Philodendron
Begonia	Juniper	Poinsettia
Coleus	Lilac	Rhododendron
Common ninebark	Marigold	Rose
Common snowberry	Mock Orange	Snapdragon
Exacum	Nannyberry	Taxus
Euonymous	Orchid	Wax plants
English ivy	Palm	Weeping fig
Gladiolus	Pansy	Zinnia

**TO KILL TENT CATERPILLARS:** Application should be made when caterpillars are young and tents are first noticed (usually late spring). For best results, apply in the late afternoon or evening when caterpillars have returned to their tents. Aim spray directly at tents, and saturate on all sides. Repeat as necessary.

#### WARRANTY STATEMENT

Our recommendations for use of this product are based upon test believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage resulting from its misuse as such, or in combination with other materials.