

Pm

19

28293-233

page 1 of 5

## UNICORN

### SEVIN<sup>(R)</sup> brand 6.3% Granular Carbaryl Insecticide

For Outdoor Home Insecticide Use  
For Home Insecticide Use

#### ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) ..... 6.3% by wt.

INERT INGREDIENTS: ..... 93.7% by wt.

Total: ..... 100.00%

Sevin is a registered trademark of Rhône-Poulenc for carbaryl insecticides


**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**See Side/Back Panel for Additional Precautionary Statements.**

**PRECAUCION AL CONSUMIDOR:** Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**(TO THE USER:** If you cannot read English, do not use this product until the label has been fully explained to you.)



NET CONTENTS:

**Manufactured by:**

Unicorn Laboratories  
13535 Feather Sound Drive, Suite 400  
Clearwater, FL 34622-5587

96 NOV 12 P12:47

REC'D EPA/CPP/DPD1

275

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed or absorbed through the skin. Avoid breathing dust. Do not take internally. Avoid contact with eyes, skin or clothing. Do not eat, drink or use tobacco while working with this product. Keep out of reach of children and domestic animals.

#### USER SAFETY REQUIREMENTS

- ◆ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- ◆ Change clothing as soon as possible after use.
- ◆ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- ◆ Do not allow children or pets to contact treated area during application and as directed, water lawn after application and allow to dry before permitting children or pets on treated area.

**ANTIDOTE STATEMENT:** ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN below.

#### Statement of Practical

**GENERAL:** Contact a physician immediately in all cases of suspected poisoning. Transport patient to a physician or hospital immediately and **SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.** If poisoning is suspected in animals, contact a veterinarian.

**IF SWALLOWED:** Never given anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger.

**IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Wash thoroughly with soap and water.

**IF INHALED:** Move from contaminated atmosphere and call a physician.

**NOTE TO PHYSICIAN:** Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Atropine is antidotal. Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are **NOT** recommended.

### **Environmental Hazards**

This product is extremely toxic to aquatic and estuarine invertebrates and toxic to birds and other wildlife.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

### **STORAGE AND DISPOSAL**

**STORAGE:** Store unused product in original container only, in cool, dry area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100°F.

**PESTICIDE DISPOSAL:** Securely wrap container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse empty bag. Discard bag in trash.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

**Wear household latex or rubber gloves. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with treated surfaces. Do not repeat application more than once a week.**

### **LAWNS, RECREATIONAL, AND ORNAMENTAL TURF AREAS**

For control of turfgrass pests, apply granules uniformly to the turf grass. For optimum control, follow specific directions for proper watering or irrigation after application. Do not allow public use of treated area during application.

INSECT	Pound of This Product Per 1000 Sq.Ft.	SPECIFIC DIRECTIONS
Ants, Armyworm, Centipedes Chiggers, Chinch bugs, Crickets, Cutworms, Darkling ground beetles, Earwigs, Essex skipper, European chafer, Fall armyworm, Flery skipper, Fleas, Grasshoppers, Green June beetle, Green June beetle grubs, Lawn moths (Sod webworm), Leafhoppers, Lucerne moth larvae, Millipedes, Sowbugs, Spittlebugs, Springtails, Ticks, Yellowstriped armyworm.	2.2 to 3.0	For general insect control, apply granules uniformly when grass is dry. For most effective control, lawn should be watered lightly after application to remove chemical from the granules.

4 7 5

White grubs (such as: Japanese beetle grubs, Chafer beetle grubs, *Phyllophaga* spp. grubs, and Annual white grubs), Bluegrass billbug larvae, European crane fly larvae

3.0

For white grub control applications should be made when grubs are feeding near the soil surface, usually during late March through May or July to early September, or as recommended by local agricultural extension agents. For most effective control, turf should be watered thoroughly after treatment to remove the chemical from the granules.

## NUISANCE PEST CONTROL IN AND AROUND FLOWER BEDS AND ORNAMENTAL PLANTINGS AND AROUND BUILDINGS

For control of ants, armyworms, ticks, cutworms, centipedes, crickets, earwigs, firebrats and silverfish, fleas, millipedes, sowbugs, scorpions and spiders; apply granules uniformly in a 6 foot band around the building. Apply granules at the rate of 5.8 ounces per 120 sq. ft. (6 x 20 foot band) to shrub and flower beds, foundations, ornamental plantings, and lawn or soil areas immediately adjacent to the building. For most effective control, treated area should be lightly watered after application to remove the chemical from the granules. Do not repeat application more than once a week.

## FRUIT AND VEGETABLE CROPS

(Not for Use in California)

For control of Ants, Centipedes, Crickets, Cutworms, Earwigs, Grasshoppers, Millipedes, Sowbugs (Pillbugs), Springtails, and Ticks, apply 0.75 lb. per 1000 ft. of this product evenly over the soil surface. Water lightly after application to release the chemical from the granules. Applications may be made at any time before, during or after the growing season. If applying during the growing season, observe indicated pre-harvest intervals (PHI). Do not repeat application more than once a week.

0 day PHI - Beans, Blueberries, Carrots, Corn, Cucumber, Eggplant, Melons, Okra, Peanuts, Pepper, Potato, Pumpkins, Squash, Sweet Potato, Tomato

1 day PHI - Asparagus, Strawberries

3 day PHI - Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Cowpeas, Garden Beet Root, Head Lettuce, Kohlrabi, Parsnips, Peas, Radishes, Rutabagas, Salsify Roots, Turnip Roots

7 day PHI - Blackberries, Dewberries (including boysenberries and loganberries), Grapes, Raspberries

14 day PHI - Celery, Chinese Cabbage, Collards, Dandelion, Endive (Escarole), Garden Beet Tops, Hanover salad, Kale, Leaf Lettuce, Mustard Greens, Parsley, Salsify Tops, Spinach, Swiss Chard, Turnip Tops

5 7 5

### IMPORTED FIRE ANT CONTROL

For use as a mound treatment to control imported fire ants, apply 1.5 to 3.0 ounces by weight over and around each fire ant mound. Irrigate mound thoroughly immediately after treatment to remove the chemical from the granules and flush it down into the mound. A minimum of 1 gallon of water should be applied to each mound, but do not use more than 2 gallons of water as flooding of the mound may result. Apply water gently to avoid disturbing ants and causing migration before chemical can be effective. For best results, apply in the early morning or late evening hours when ants are least active. Treat new mounds as they appear.

### CONTROL OF TICKS WHICH TRANSMIT LYME DISEASE

For control of juvenile and adult Ixodes spp. ticks (Deer tick, Bear tick, and Black legged tick) and Amblyomma spp. ticks (Lone star tick). Apply granules at the rate of 2.5 to 3.0 pounds of formulated product per 1000 square feet.

Begin applications in late spring or early summer (May/June). Uniformly apply granules to entire lawn, perimeter wooded areas, and property boundaries including shrub beds, ornamental plantings, and wooded areas where exposure to ticks may occur. For optimum results, irrigate treated areas after application to remove chemical from granules.

### NOTICE OF DISCLAIMER

BUYER SHOULD BE SOLELY RESPONSIBLE FOR ANY AND ALL INJURY, LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS, OR CAUTIONS.