



U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 MMH St., S.W. Washington, D.C. 20460

NOTICE OF PESTICIDE:

\_x Registration Reregistration

(under FIFRA, as amended)

EPA Reg. Number: 28293-

237

Date of Issuance:

JAN 06 135:

Term of Issuance: Until Reregistration

Name of Pesticide Product:

Unicorn Sevin Brand Carbaryl Insecticide 5% Dust

Name and Address of Registrant (include ZIP Code):

Unicorn Laboratories 1000 118th Avenue North St. Petersburg, FL 33716

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
- 2. Add the phrase "EPA Registration No. 28293-237" to your label before you release the product for shipment.
- 3. Add appropriate preharvest intervals for eggplant, pepper, and tomato. (Illay he explicate up to horsest).

Signature of Approving Official:

DENNIS H. EDWARDS

Date:

Dennis H. Edwards, Jr. Product Manager 19 Insecticide-Rodenticide Branch Registration Division JAN 0 3 1994

E?A Form 8570-6

page 2 EPA Reg. No. 28293-237

4. Submit five copies of the revised labeling for this product before you release this product for shipment.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR part 156, subpart K, section 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

DENINIS H. EDWARDS

Dennis H. Edwards, Jr. Product Manager 19 Insecticide-RodenticideBranch Registration Division (H7505C)

Enclosure

3.36

### UNICORN

# SEVIN<sup>(R)</sup> brand carbaryl insecticide 5% DUST

## KILLS GARDEN PESTS, PLEAS & TICKS

IN THE GARDEN: Armyworms, Cutworms, Squash Bugs, Tomato Hornworms, certain other Beetles and other Garden Pests, as listed.

ON DOGS AND CATS: Fleas and Ticks on Animals and other outdoor living areas.

ACTIVE INGREDIENT:  Carbaryl (1-naphthyl N-methylcarbamate)	5%
INERT INGREDIENTS:	95%
SEVIN <sup>(R)</sup> is the registered trademark of Rhône Poulenc for carbaryl insecticide.	

### KEEP OUT OF REACH OF CHILDREN

### CAUTION

See Back/Side Panel for Additional Precautionary Statements.

Net Weight:

Manufactured by:

Unicorn Laboratories 1000 118th Avenue North St. Petersburg, FL 33716

ACCLETED
with COMMENTS
in EPA Letter Deted

JAN - 6 1994

Under the Foderal Insecticide. Fungicide, and Redenticide Act to succeeded, for the posticide registered under EPA Reg. No. 28243 - 237

EPA REG. NO. 28293-

EPA EST. NO. 28293-FL-1

4.56

# PRECAUTIONARY STATEMENTS Hazards to Humi 's and Domestic Animals

CAUTION: Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly af 'r handling. Avoid contamination of food, feed, feeding troughs and water receptacles.

Statement of Practical Treatment

If Swallowed: Give water or milk to drink and telephone for medical advice. Consult medical personnel before inducing vomiting. If medical advice cannot be obtained, then take person and product container to the nearest medical emergency treatment center or hospital.

If In Eyes: In case of eye contact, wash eyes immediately with fresh water for at least 15 minutes. Get medical attention if irritation persists.

If On Skin: Wash with plenty of soap and water.

If Inhaled: Remove person from exposure area.

Note to Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

Environmental Hazards

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Apply dust at the first signs of insects. Repeat as necessary. For best results, dust in early morning or late evening when temperature is low and air is still. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces. May be applied up to harvest unless otherwise noted.

BEANS: Mexican Bean Beetle - 1/4 lb. per 1,000 sq. ft.
Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean
Cutworm, Velvetbean Caterpillar, Flea Beetles, Japanese Beetle 1/2 lb per 1,000 sq. ft. Corn Earworm, Armyworm, Tarnished Plant
Bug, Stink Bugs - 3/4 lb. per 1,000 sq. ft. May be applied up to
harvest.



5.46

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI: Plea Beetles, Harlequin Bug - 1/2 lb. per 1,000 sq. ft. Armyworms, Imported Cabbageworm, Young Cabbage Loopers, Salt Marsh Caterpillars, Corn Earworm - 1 lb. per 1,000 sq. ft. Do not apply within 3 days of harvest.

CORN: Corn Earworm (Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations). Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetle, Leafhoppers, Flea Beetles - 1 lb. per 1,000 sq. ft. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. May be applied up to harvest.

CUCUMBER, MELONS, PUMPKIN, SQUASH: Pickleworm, Melonworm - 1/2 lb. per 1,000 sq. ft. Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs - 1/2 lb. per 1,000 sq. ft. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use this product on watermelons in Florida. May be used up to harvest.

LETTUCE, CARROTS: Flea Beetles, Harlequin Bug, Leafhoppers - 1/2 lb. per 1,000 sq. ft. Sixspotted Leafhopper - 3/4 lb. in a 5 to 7 day schedule. Armyworms, Cabbage Caterpillars, Corn Earworm, Spittlebugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs - 1 lb. per 1,000 sq. ft. For best results in the control of Corn Earworm after lettuce heads begin to form use 1 lb. in a 5 to 7 day schedule. May be used up to harvest on carrots. Do not use on head lettuce within 3 days of harvest and to leaf lettuce within 14 days.

EGGPLANT, PEPPERS: Colorado Potato Beetles, Flea Beetles - 1/2 lb. per 1,000 sq. ft. Tomato Fruitworm, Tomato Hornworm, European Corn Borer, Lace Bugs - 1 lb. per 1,000 sq. ft.

STRAWBERRIES: Meadow Spittlebug, Strawberry Leafroller, Strawberry Weevil - 1 lb. per 1,000 sq. ft. Do not apply within 1 day of harvest.

TOMATO: Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer - 1 1/2 lbs. per 1,000 sq. ft.

POULTRY: (Chickens, Partridges, Pheasants, Pigeons): Northern Fowl Mite, Lice and as a supplement to premise treatments.for Chicken Mite, Fleas ~ 1 lb. per 100 birds. Apply thoroughly by means of a shaker can or hand duster. Repeat in 4 weeks if necessary. Do not use on Poultry or game birds within 7 days of slaughter.



IN OR AROUND POULTRY HOUSES: Chicken Mite, Fleas, Bed Bugs - 1 lb. per 40 sq. ft. of litter or floor space and to roosts and adjacent areas by means of a shaker can, hand duster or power duster. Force dust into cracks and crevices. Repeat as necessary. Do not treat nest litter. Avoid direct contamination of eggs. Do not use in premise treatment within 7 days of slaughter.

DOGS, CATS: Ticks, Fleas - Dust lightly over the animal and rub thoroughly into the skin. Apply also to the feet between the toes and on the legs. Apply at weekly intervals if needed. Avoid treatment of kittens and puppies under 4 weeks of age. NOTE: Do not allow treated pets back on carpet or rugs in the house as this product may stain or discolor rugs and wall-to-wall carpets. Dust lightly in and around kennels and dog houses by means of a shaker can, hand or power duster. Force dust into cracks and crevices. Repeat as necessary, but not more often than once weekly. DO NOT USE ON PREGNANT DOGS.

DO NOT USE IN THE HOME: DO NOT APPLY TO RUGS AND CARPETS, AS STAINING MAY OCCUR.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

## STORAGE AND DISPOSAL

**STORAGE:** Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

**DISPOSAL:** Product - Partially filled bag may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. Container - Do not reuse bag. Discard bag in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

EPA REG. NO. 28293-

EPA EST. NO. 28293-FL-1