UNICORN SYNERGIZED POUR-ON INSECTICIDE

Kills lice and flies on cattle, and keds on sheep For Use on Cattle, Sheep and Their Premises

ACTIVE INGREDIENTS

Permethrin*	1.0%
Piperonyl Butoxide Technical**	1.0%
INERT INGREDIENTS***	98.0%
Total:	100.0%

* Cis/trans ratio: Min. 35% (±) cis

Max. 65% (±) trans

** Equivalent to Min. 0.8% (butylcarbityl)(6-propylpiperonyl) ether and 0.2% related compounds.

*** Contains Petroleum Distillate

Keep Out of Reach of Children CAUTION

See Side Panel for Additional Precautionary Statements

NET CONTENTS: ONE U.S. GALLON (3.785 Liters)

Manufactured by:

Unicorn Laboratories 12385 Automobile Blvd. Clearwater, Florida 33762

Dogg 1

EPA EST. NO. 62478-FI.-1A 44616-MO-1B Subscript used is first letter of run code on container.

ACCEPTED

JAN 23 2002

Under the Federal Insecticide.
Punglicide, and Redomicilide Act, at amenden, for the posticide registered under

EPA REG. NO. 28293-259 (PRN 2001-1)

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes and clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling.

	FIRST AID
If Swallowed:	 Call poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
If in Eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Have the product containe for treatment.	er or label with you when calling a poison control center or doctor, or going
	NOTE TO PHYSICIAN
Contains petroleum distilla	te – vomiting may cause aspiration pneumonia.

Environmental Hazards

This product is extremely toxic to fish and other aquatic invertebrates. Do not add directly to water, or to areas where surface waters are present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep container sealed when not in use. Do not store near food or feed.

PESTICIDE Wastes resulting from the use of this product may be disposed of on site or at

DISPOSAL: an approved waste disposal facility.

EPA EST. NO. 62478-FL-1A 44616-MO-1B Subscript used is first letter of run code on container. CONTAINER DISPOSAL:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READY TO USE - No dilution necessary.

APPLY TO	TARGET INSECTS	APPLICATION INSTRUCTIONS
Lactating and Non-Lactating	Lice, Horn Flies, Face Flies.	Dosage: Apply ½ fl. oz (15
Dairy Cattle, Beef Cattle	Aids in control of Horse Flies,	cc) per 100 lbs body wt. of
and Calves	Stable Flies, House Flies,	animal up to a maximum of
•	Mosquitoes and Black Flies.	5 fl. oz. for any one animal.
		Pour-On: Pour correct dose along back and down face. Ready-To-Use Spray: Use undiluted in a mist sprayer to apply correct dose. Apply directly to neck, face, back, legs and
		ears. Back Rubber Use: Mix one pint per gallon #2 diesel or mineral oil. Keep rubbing device charged. Results improved by daily forced use.
Sheep	Sheep Keds, Lice	Pour along back. Apply ¼ fl. oz. (7.5 cc) per 50 lbs body wt. of animal, up to a maximum of 3 fl. oz. for any one animal. For optimum control, all animals in the flock should be treated after shearing.
Premises - in and around Horse, Beef, Dairy, Swine, Sheep and Poultry premises, Animal Hospital Pens and Kennels and "outside" Meat Processing Premises.	House Flies, Stable Flies, Face Flies, Gnats, Mosquitoes, Black Flies, Fleas, Little House Flies (Fannia spp.). Aids in control of cockroaches, ants, spiders, crickets.	For use as a ready-to-use spot spray or premise spray, use undiluted in a mist sprayer. Apply directly to surface to leave a residual insecticidal coating, paying particular attention to areas where insects crawl or alight. One gallon will treat approximately 7,300 square feet.

For cattle and sheep, repeat treatment as needed, but no more often than once every two weeks. For optimum lice control, two treatments at a 14-day interval are recommended.

Special Notes: This product is not effective in controlling cattle grubs. This is an oil-base, ready- to-use product, that may leave an oily appearance on the hair coat of some animals. This product should be used in an integrated pest management system which may involve repeat treatments and the use of other pest control practices.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.