



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:

28293-
262

Date of Issuance:

7 MAR 1995

NOTICE OF PESTICIDE:

 x Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Unicorn Ear
Miticide IV

Name and Address of Registrant (include ZIP Code):

Unicorn Laboratories
One Corporate Drive, Suite 400
Clearwater, FL 34622

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes:

- a. Revise the EPA Registration Number to read, "EPA Reg. No. 28293-262".
- b. Under the Directions for Use, remove all references to Ear Mite Control.
- c. Submit efficacy data or cite a product with the same use pattern prior to seeking approval to add control for ear mites.

Signature of Approving Official:

Rebecca S. Cool

Date:

3/6/95

page 2
EPA Reg. No.

3. Submit five copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

UNICORN EAR MITICIDE IV

For Use on Cats and Dogs
and on Premises to Control Fleas and Ticks

An Aqueous, Stable, Micro Emulsion

Alcohol Free ■ Prevents Dry Skin

Kills Ear Mite on Dogs, Cats & Rabbits

ACTIVE INGREDIENTS:

Pyrethrins.....	0.05%
Piperonyl Butoxide, Technical*.....	0.50%

INERT INGREDIENTS:..... 99.45%

*Equivalent to min. 0.40% (butylcarbityl) (6-propylpiperonyl) ether and 0.10% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back/Side Panel for Additional Precautions

NET CONTENTS:

Manufactured by:
Unicorn Laboratories
One Corporate Drive, Suite 400
Clearwater, Florida 34622-5587

BEST AVAILABLE COPY

EPA Reg. No. 28293-

ACCEPTED
with COMMENTS
in EPA Letter Dated

7 MAR 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

28293-262

EPA EST. NO. 62478-FL-1A
44616-MO-1B

Subscript used is first letter
of run code on container.

PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid inhalation of vapors. Wash contaminated skin promptly with soap and water after use. Avoid spraying food or foodstuffs. Remove birds and cover fish aquariums before spraying.

Statement of Practical Treatment

IF SWALLOWED: Contact a physician or Poison Control Center. Induce vomiting by touching back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES OR ON SKIN: Flush affected areas with plenty of water. Contact a physician if irritation persists.

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a cool dry place inaccessible to children and pets and away from heat and direct sunlight. Protect from freezing.

DISPOSAL: Do not reuse empty container. Wrap container in several layers of newspaper and discard in trash.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USE

ON PETS

Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and under body. While spraying, fluff the hair so that spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes, face or on genitals. Repeat as necessary. Do not spray on nursing females or pets under six weeks of age.

FOR EAR MITE CONTROL

DIAGNOSIS: Ear mites usually induce the formation of a dry, dark brown, waxy exudate with crusts in the ears of dogs, cats, rabbits. Ear mites may be detected by placing some of the ear exudate on a dark surface and carefully watching for the movement of tiny white specks away from the exudate. Inflamed, watery, or blocked ear canals indicate a more serious condition which requires the services of a veterinarian.

BEST AVAILABLE COPY

647

APPLICATION: While holding pet firmly, fill each ear canal half full of product and massage base of ear to insure that insecticidal action penetrates ear wax. Repeat treatment every other day until condition is relieved. Improvement is usually noted after two applications.

ON LIVESTOCK

To protect cattle, horses and hogs from **Horn Flies, House Flies, Mosquitoes** and **Gnats**, apply a light mist sufficient to wet the surface of the hair. To control **Stable Flies, Horse Flies** and **Deer Flies**, apply at a rate of 2 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control **Blood Sucking Lice**, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

To control **Poultry Lice**, spray roosts walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of **Bedbugs** and **Mites** in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the **Bedbugs** and **Mites** hide.

ON PREMISES

To control **Fleas** and **Brown Dog Ticks** on premises: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

BEST AVAILABLE COPY