

28293-268

3/24/2015

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460-0001

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

March 24, 2015

SENT BY E-MAIL

Ms. Rebecca Horton
Agent for Phaeton Corporation
P.O. Box 1019
Salem, VA 24153

Subject: Amended Label for Unicorn permethrin Multi-Use Insecticide Spray I
EPA Reg. No.28293-268
Decision Number: D413435

Dear Ms. Horton:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the reregistration of the referenced product in connection with the permethrin RED(s), and has concluded that your submission is acceptable.

PLEASE NOTE: This product is not being reregistered under sections 3(c)5 and 4(g) of FIFRA at this time.

A copy of your label stamped "Acceptable" is enclosed along with copies of the acute toxicity and product chemistry reviews completed for the subject product. Products shipped after 12 months from the date of this amendment or the next printing of the label whichever occurs first, must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e).


If you have any questions about this letter, please contact Linda A. DeLuise at (703) 305-5428 or via e-mail at deluise.linda@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Richard Gebken".

Richard Gebken
Product Manager 10
Invertebrate & Vertebrate Branch 2
Office of Pesticide Programs

2/13



[MASTER LABEL – COMMERCIAL/INDUSTRIAL USE]

UNICORN PERMETHRIN MULTI-USE INSECTICIDE SPRAY I

Optional Marketing Statements:

- FOR USE ON DOGS AND ON PREMISES TO CONTROL FLEAS AND TICKS
- CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS
- Effective Treatment for Sarcoptic Mange.
- CONTROLS RESIDENTIAL, INDUSTRIAL AND GARDEN INSECTS
- CONTROLS PESTS FOR UP TO 4 WEEKS
- FOR USE ON LIVESTOCK AND LIVESTOCK PREMISES
- KILLS LICE AND THEIR EGGS
- FOR USE ON BEDDING, FURNITURE AND OTHER INANIMATE OBJECTS INFESTED WITH LICE
- NOT FOR USE ON HUMANS

ACTIVE INGREDIENT:

Permethrin* 0.50%

OTHER INGREDIENTS:**..... 99.50%

Total: 100.00%

*Cis/Trans Ratio: Min. 35% (±) Cis
Max. 65% (±) Trans

**Contains petroleum distillate.

Do Not Use on Cats

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back (Side) Panel for Additional Precautionary Statements and First Aid

NET WEIGHT:

EPA Reg. No. 28293-268
March 16, 2015 – reregis revised

EPA Est. No. _____
Page 1 of 12 Pages

READ ENTIRE LABEL BEFORE EACH USE.
FOR PET USE – USE ONLY ON DOGS.

FIRST AID	
If on Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact [x-xxx-xxx-xxxx] for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Contains petroleum distillate – vomiting may cause aspiration pneumonia.	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear long sleeved shirt and long pants, socks and shoes, and chemical resistant gloves (such as those made out of Barrier Laminate, Nitrile Rubber, Neoprene Rubber or Viton, selection category E).

ANIMALS: Do not spray directly in/on eyes, mouth or genitalia of pets. Do not use on puppies under twelve (12) weeks of age. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Do Not Use on Cats. May be toxic or potentially fatal if applied to or ingested by cats. Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movements. If this occurs, immediate veterinary care should be provided.



User Safety Requirements: Follow manufacturer’s instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other

absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations: User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. User should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

Physical and Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperature above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions

Remove pets and birds and cover fish aquaria before space spraying or surface applications. Except when applying to pets or livestock, do not apply this product in a way that will contact workers or other persons, either directly or through drift. Except when applying to pets or livestock, only protected handlers may be in the area during application. Except when applying to pets or livestock, do not enter or allow others to enter until sprays have dried, and aerosols have dispersed, and the treated area has been thoroughly ventilated. Do not use in aircraft. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Do not apply when food is present. Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Nonfood areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling.)

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches.

HOUSEFLIES, FRUIT FLIES, MOSQUITOES, GNATS, WASPS, HORNETS, BEES, BLACK FLIES, SKIPPER FLIES, BLOW FLIES AND SMALL FLYING MOTHS: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Retreat if infestation occurs, but not more than once every two weeks.

COCKROACHES, WATERBUGS, PALMETTO BUGS, SILVERFISH, FIREBRATS, CRICKETS, SCORPIONS, SPIDERS, MILLIPEDES, CENTIPEDES, SOWBUGS AND PILLBUGS: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray when ever possible. Retreat if infestation occurs, but not more than once every two weeks.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Retreat if infestation occurs, but not more than once every two weeks.

INDIAN MEAL MOTHS, ANGOUMOIS GRAIN MOTHS: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Retreat if infestation occurs, but not more than once every two weeks.

EXPOSED STAGES OF CONFUSED FLOUR BEETLES, SAW TOOTHED GRAIN BEETLES, RED FLOUR BEETLES, RICE WEEVILS, GRANARY WEEVILS, GRAIN MITES: Spray into cracks and crevices, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Retreat if infestation occurs, but not more than once every two weeks. Infested stored food products in the home should be disposed of.

CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets. Retreat if infestation occurs, but not more than once every two weeks.

BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray onto all joints. Treat baseboards, moldings and floors. If lice infestations should occur on humans, use a product labeled for use on humans. Spray to the point of dampness but not runoff. After mattress is dry, cover with mattress cover and sheet. Do not use mattress without cover. Retreat if infestation occurs, but not more than once every two weeks.

To control Clothes Moths and Carpet Beetles on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage directing the spray into cracks, joints and crevices. Spray to the point of dampness but not runoff. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining.

IN ANIMAL QUARTERS AND MILK ROOMS: For rapid control of House Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees, and Small-Flying Moths, close window and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. When used in dairy barns or facilities, close milk bulk tank lids to prevent contamination from spray and from dead or falling insects.

TO TREAT LIVESTOCK: To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitoes, and Gnats: Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.

FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment.

TO CONTROL BLOOD SUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

FOR CONTROL OF BEDBUGS AND MITES IN POULTRY HOUSES, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide. Spray to the point of dampness but not runoff.

FLEAS AND TICKS: (ADULT AND LARVAE): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use. Allow all sprayed articles to dry thoroughly before use.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry thoroughly before use.

TO CONTROL FLEA, TICKS AND LICE ON DOGS: Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Retreat if infestation occurs, but not more than once every two weeks. Do not use on puppies under twelve (12) weeks of age.

TREATMENT FOR SARCOPTIC MANGE OF DOGS: Clip hair around all affected areas. Then wash the animal thoroughly with soap and water. Allow animal to dry before treating. Hold can in an upright

position 3" to 6" from the dog and spray the affected area. Lightly wet the skin. Avoid soaking the skin and treating more than 1/3 of the animal at any one time. Remove and destroy infested bedding.

OUTDOORS:

Do not make applications during rain. Do not wet plants to point of runoff or drip. Do not water the treated area to the point of runoff. All outdoor applications must be limited to spot or crack-and-crevice treatments only except for the following permitted uses:

- 1) Treatment to soil or vegetation around structures
- 2) Applications to lawns, turf, and other vegetation

All outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.

FOR THE CONTROL OF GARDEN INSECTS: For outdoor use only as an aid in controlling these insects. For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, exposed Thrips, Spiders, Red Mites and Leafminers.

May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, pavement Ants, Clover Mites and Army Worms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90° F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

STORAGE AND DISPOSAL

Pesticide Storage: Store in a cool, dry area. Do not transport or store below 32° F.

Product Disposal: If partly filled – Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Disposal: Do Not Puncture or Incinerate! If empty – Place in trash or offer for recycling if available.

Notice: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

**Phaeton Corporation
P.O. Box 290
Madison, GA 30650**

[Lot number added at production.]

8/13

[SUBLABEL A – RESIDENTIAL USE]

UNICORN PERMETHRIN MULTI-USE INSECTICIDE SPRAY I

Optional Marketing Statements:

FOR USE ON DOGS AND ON PREMISES TO CONTROL FLEAS AND TICKS
CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS
Effective Treatment for Sarcoptic Mange.
CONTROLS RESIDENTIAL AND GARDEN INSECTS
CONTROLS PESTS FOR UP TO 4 WEEKS
FOR USE ON NON-COMMERCIAL LIVESTOCK AND LIVESTOCK PREMISES
KILLS LICE AND THEIR EGGS
FOR USE ON BEDDING, FURNITURE AND OTHER INANIMATE OBJECTS INFESTED
WITH LICE
NOT FOR USE ON HUMANS
FOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENT:

Permethrin*	0.50%
OTHER INGREDIENTS**:	99.50%
Total:	100.00%

*Cis/Trans Ratio: Min. 35% (±) Cis
Max. 65% (±) Trans

**Contains petroleum distillate.

Do Not Use on Cats

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back (Side) Panel for Additional Precautionary Statements and First Aid

NET WEIGHT:

READ ENTIRE LABEL BEFORE EACH USE.
FOR PET USE – USE ONLY ON DOGS.

FIRST AID	
If on Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact [x-xxx-xxx-xxxx] for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Contains petroleum distillate – vomiting may cause aspiration pneumonia.	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear long sleeved shirt and long pants, socks and shoes. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ANIMALS: Do not spray directly in/on eyes, mouth or genitalia of pets. Do not use on puppies under twelve (12) weeks of age. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Do Not Use on Cats. May be toxic or potentially fatal if applied to or ingested by cats. Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movements. If this occurs, immediate veterinary care should be provided.



Environmental Hazards

This product is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

10/13

Physical and Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperature above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions

Remove pets and birds and cover fish aquaria before space spraying or surface applications. Except when applying to pets, companion animals or non-commercial livestock, do not apply this product in a way that will contact workers or other persons, either directly or through drift. Except when applying to pets, companion animals or non-commercial livestock, only protected handlers may be in the area during application. Except when applying to pets, companion animals or non-commercial livestock, do not enter or allow others to enter until sprays have dried, and aerosols have dispersed, and the treated area has been thoroughly ventilated. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches.

HOUSEFLIES, FRUIT FLIES, MOSQUITOES, GNATS, WASPS, HORNETS, BEES, BLACK FLIES, SKIPPER FLIES, BLOW FLIES AND SMALL FLYING MOTHS: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Retreat if infestation occurs, but not more than once every two weeks.

COCKROACHES, WATERBUGS, PALMETTO BUGS, SILVERFISH, FIREBRATS, CRICKETS, SCORPIONS, SPIDERS, MILLIPEDES, CENTIPEDES, SOWBUGS AND PILLBUGS: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray when ever possible. Retreat if infestation occurs, but not more than once every two weeks.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Retreat if infestation occurs, but not more than once every two weeks.

INDIAN MEAL MOTHS, ANGOUMOIS GRAIN MOTHS: Close doors and windows, direct spray into all parts of room especially around stored food product containers and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Retreat if infestation occurs, but not more than once every two weeks.

EXPOSED STAGES OF CONFUSED FLOUR BEETLES, SAW TOOTHED GRAIN BEETLES, RED FLOUR BEETLES, RICE WEEVILS, GRANARY WEEVILS, GRAIN MITES: Spray into cracks and crevices where these insects tend to congregate. Spray 30 seconds for each 100 feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Retreat if infestation occurs, but not more than once every two weeks. Infested stored food products should be disposed of.

CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets. Retreat if infestation occurs, but not more than once every two weeks.

BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray onto all joints. Treat baseboards, moldings and floors. If lice infestations should occur on humans, use a product labeled for use on humans. Spray to the point of dampness but not runoff. After mattress is dry, cover with mattress cover and sheet. Do not use mattress without cover. Retreat if infestation occurs, but not more than once every two weeks.

To control Clothes Moths and Carpet Beetles on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage directing the spray into cracks, joints and crevices. Spray to the point of dampness but not runoff. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining.

IN COMPANION AND NON-COMMERCIAL ANIMAL QUARTERS AND MILK ROOMS: For rapid control of House Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees, and Small-Flying Moths, close window and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. When used in dairy barns or facilities, close milk bulk tank lids to prevent contamination from spray and from dead or falling insects.

TO TREAT COMPANION AND NON-COMMERCIAL LIVESTOCK: To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitoes, and Gnats: Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.

FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment.

TO CONTROL BLOOD SUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

FOR CONTROL OF BEDBUGS AND MITES IN NON-COMMERCIAL POULTRY HOUSES, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide. Spray to the point of dampness but not runoff.

FLEAS AND TICKS: (ADULT AND LARVAE): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use. Allow all sprayed articles to dry thoroughly before use.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry thoroughly before use.

TO CONTROL FLEA, TICKS AND LICE ON DOGS: Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Retreat if infestation occurs, but not more than once every two weeks. Do not use on puppies under twelve (12) weeks of age.

TREATMENT FOR SARCOPTIC MANGE OF DOGS: Clip hair around all affected areas. Then wash the animal thoroughly with soap and water. Allow animal to dry before treating. Hold can in an upright position 3" to 6" from the dog and spray the affected area. Lightly wet the skin. Avoid soaking the skin and treating more than 1/3 of the animal at any one time. Remove and destroy infested bedding.

OUTDOORS:

Do not make applications during rain. Do not wet plants to point of runoff or drip. Do not water the treated area to the point of runoff. All outdoor applications must be limited to spot or crack-and-crevice treatments only except for the following permitted uses:

- 1) Treatment to soil or vegetation around structures
- 2) Applications to lawns, turf, and other vegetation

All outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.

FOR THE CONTROL OF GARDEN INSECTS: For outdoor use only as an aid in controlling these insects. For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, exposed Thrips, Spiders, Red Mites and Leafminers.

May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, pavement Ants, Clover Mites and Army Worms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90° F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

STORAGE AND DISPOSAL

Pesticide Storage: Store in a cool, dry area. Do not transport or store below 32° F.

Product Disposal: If partly filled – Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Disposal: Do Not Puncture or Incinerate! If empty – Place in trash or offer for recycling if available.

Notice: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Phaeton Corporation
P.O. Box 290
Madison, GA 30650

[Lot number added at production.]