5-

9 23 2013

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

Ms. Rebecca Horton Agent for Phaeton Corporation P.O. Box 1019 Salem, VA 24153 OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

SEP 2 3 2013

Subject: Label Notification(s) for Pesticide Registration Notice 98-10

Dear Ms. Horton:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated July 8, 2013 for the following product(s):

Unicorn 2.5% Permethrin Concentrate/Repellent EPA Reg. No. 28293-271

The Registration Division (RD) has conducted a review of this request for applicability under PR Notice 98-10 and finds that the label change(s) requested falls within the scope of PRN 98-10. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, please contact Linda A. DeLuise at 703-305-5428.

Sincerely,

Richard J. Gebken Product Manager Insecticide Branch Registration Division (7504P)

	\langle					(2
	Environmenta	United States	- · ·	Form Ar		Reg	jistra endr	2070-00/ Ition nent	80. Approvel expires 2-28-99 OPP Identifier Number
	-	Applicati	on for Pesti	cide - Sec	tion	1			
I. Company/Product Numbe 28293-271	70		2. EPA Product Manager Gebken			roposed Classification			
. Company/Product (Name Unicorn 2.5% Permeth) rin Concentrate/Re	pellent	PM# Team 10						
5. Name and Address of Ap Phaeton Corporation P.O. Box 290 Madison, GA 30650		ode)	(b)(i), to: EPA	pedited Re my product Reg. No luct Name	is sim	ilar d	NOT	ifaloxo	n FIFRA Section 3(c)(3) pmposition and labeling 013
<u>ب</u>	· · · ·		Section						
Agency letter dated Me Too" Application. We Too" Application.					or the Confidential Statement of her understands that if this				
		·	Section -	111					
. Material This Product Wil									
hild-Resistant Peckeging Yes V No Certification must	Unit Packaging Yes Vo If "Yes" Unit Packaging wgt.	No. per container	Water Soluble Yes No If "Yes" Package wgt	No. per containe		2. 1		Containe Metal Plastic Glass Paper Other (r Specify)
e submitted			·			`			
	ontainer		tail Container 8 fl oz to 1 gal	[- 1	of Lab	el Directi	ons
. Manner in Which Label is	ATTIXED TO Product	Paper Stenc	iraph glued iled	Othe	r			· ····	25 · · · · · · · · · · · · · · · · · · ·
			Section -	IV		<u></u>			····
Contact Point (Complete	items directly below f	or identificatio			if nec	essary,	, to pr	ocess thi	s application.)
Name Rebecca M. Horton			Title Consultant/Agent			Telephone No. (Include Aree Code) 540-375-8826			
l certify that the state I acknowledge that an both under applicable	y knowlinglly felse or		all attachments t						6. Date Application Received (Stamped)
Signature Kultura Mi	Hortiz		3. Title Consultant/Agent						
Rebecca M. Horton			5. Date Ju	ly 8, 2013					

Pegistrations esign, Jnc.

P.O. Box 1019 Solem, VA 24153-1019 phone 540.375.8826 fax 540.375.8827 USPS PRIORITY DELIVERY CONFIRMATION #9405 5036 9930 0488 9026 35 July 10, 2013

Document Processing Desk (NOTIF) Office of Pesticide Programs/Registration Division (7504P) U.S. Environmental Protection Agency Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, DC 20460

Re: Notification: EPA Reg. No. 28293-271

Attn: Richard Gebken, PM Team 10

On behalf of Phaeton Corporation of Madison, GA, enclosed is notification of administrative label changes for the above referenced pesticide registration, including optional marketing statements consistent with currently approved label use and optional use of supplemental distributor website.

The following documents are enclosed:

- 1) EPA Form 8570-1, Application for Pesticide Notification (2 copies)
- 2) Copy of the label with areas of revision highlighted.

Written confirmation of acceptance of this notification is requested.

Best regards,

ilucca M Hoston

Rebecca M. Horton Consultant/Agent for PHAETON CORPORATION

e-mail: hortonb@ntelos.net

/bmh Enclosures

Federal & State Pesticide/Fertilizer Registration Services *plus* typesetting/design of product labeling

Established 1996

Unicorn 2.5% Permethrin Concentrate /Repellent

For Use Around Residential [Home] Areas Only (editors note: [] brackets indicate alternate/optional text)

Optional Marketing Statements:

Up to 4 Week Control

Kills [Controls] Mosquitoes, Fleas, Ticks and Ants

Water Based

[Also] Kills [Fleas,] [Ticks,] Japanese Beetles, [Mosquitoes], [Fire Ants] and many other insects [garden pests as listed]

Kills and Repels

Kills Wood Infesting Insects

Kills [Deer] Ticks that [which] may carry Lyme Disease

Kills Carpenter Bees & Beetles

[Also] Stops[s] Bugs in Your [Also use on] [Use on]

- Lawns
- Trees & Shrubs
- Roses & Flowers
- Fruits & Nuts
- Vegetables
- Gardens

Kills Termites, Carpenter Ants, Carpenter Bees, Powderpost Beetles and other listed insects Treat outside surfaces of buildings, porches, wooden decks, patios. fences, foundations, wood piles and tree stumps *(editors note: for final printed labels with wood infesting insects directions)* Spray Around Yard Perimeter

32 oz. Treats Up to 5,000 Square Feet

Hose End Use – Always Shake Well Before Using

Easy To Use Hose-End Spray

Kills 60+ Insects

Contains Permethrin, a long lasting Insecticide which provides contact and residual control for up to 4 weeks of many hard-to-kill sucking and chewing insects.

PRODUCT USAGE [FACTS] [QUICK FACTS]

KILLS: [Ants,] Aphids, Beetles, [Chinch Bugs, Crickets, Japanese Beetles,] [Cutworms,] [Earwigs,] [Fleas,] [Japanese Beetle Grubs,] Mealybugs, [Mites,] [Mole Crickets,] [Mosquitoes,] [Sod Webworms,] Spider Mites, [Tent Caterpillars,] [Ticks,] Whiteflies and many other[s] [garden [nuisance] pests] [more insects] as listed.

Also kills wood-infesting insects including termites and Carpenter Ants. (editors note: for final printed labels with wood infesting insects directions)

USE ON [WHERE TO USE]: Fruits [Fruit Trees], Vegetables, Roses, Flowers, Lawns, Trees, and Shrubs, [Flowers] [and as a residual spray to outside surfaces of buildings, foundations, porches, patios, window frames, and other areas where pests congregate.] [Outdoors around the home]

AMOUNT TO USE: See detailed instruction inside label. [Roses, Flowers, Trees, Shrubs, Fruits, Vegetables: 1-3 fl. oz. per gallon of water] [Lawns: 2-6 fl. oz. per 1,000 sq. ft.] [Nuisance pests: 6 2/3 oz. per 400 linear ft. by 6 inch spray band] [See directions for specific application rates.]

EPA REG NO. 28293-271 July 8, 2013 - notif

. -

EPA EST. NO. [Subscript used is first letter of run code on container.] Page 1 of 14 Pages

NOTIFICATION

SEP 2 3 2013

PRODUCT FACTS [QUICK FACTS]

- KILLS Lawn Pests: Chinch bugs, Cutworms, Mole Crickets, Japanese Beetle Grubs & Sod Webworms
 - Tree, Shrub & Flower pests: Aphids, Beetles, Tent caterpillars, Mealybugs, Mites, Whiteflies & other garden pests
 - Nuisance Pests (outdoors): Ants, Earwigs, Fleas, Mosquitoes and other nuisance pests
- WHERE TO USE On Roses, Flowers Trees, Shrubs & Lawns

• Outdoors around the home

(Optional text below in italics can be used for final printed labels with wood infesting insects directions)

[PRODUCT USAGE [QUICK FACTS]

KILLS: Termites (including dry wood termites), Carpenter Ants, Carpenter Bees, Powderpost Beetles and other wood infesting pests.

USE ON: Outside surface of buildings, porches, wooden decks and patios, wooden fences, window frames, doorways, foundations, eaves, patios, garages, wood piles, tree stumps and other areas where you may find these pests.

AMOUNT TO USE: Mix 10 fl. oz. with 1 gallon of water to treat 100 sq. ft.]

(Optional text below in italics used when packaged in a hose-end applicator container only.) *PRODUCT USAGE [QUICK FACTS]*

KILLS: Ants, Chinch Bugs, Crickets, Japanese Beetles, Mosquitoes, Ticks, Whiteflies, and many other insects as listed [other listed insects].

USE ON: Lawns, Trees, Shrubs, Roses, Flowers, and as a residual spray to outside surfaces of buildings, porches, patios, window frames, and other areas where pests congregate. AMOUNT TO USE: No mixing required. Connect to hose and spray. [Ready to use Hose-End Spray]

(Optional text below in italics used when packaged in certain 32 fl. oz. hose-end applicator container only.)

Kills: Kills troublesome and damaging insects as listed.

Use On: Lawn and other outdoor areas where pests congregate.

Amount to Use: NO MIXING. Connect to hose and spray. Treats up to 5,000 sq. ft.

ACTIVE INGREDIENT:

Permethrin*	
OTHER INGREDIENTS:	
	$1\overline{00.0\%}$

*Cis/trans isomer ratio: Min. 35% (±) cis, Max. 65% (±) trans

KEEP OUT OF REACH OF CHILDREN CAUTION

See Side/Back/Other Label Panel(s) for Additional Precautionary Statements & First Aid

NET CONTENTS:

EPA EST. NO. [Subscript used is first letter of run code on container.] Page 2 of 14 Pages

ÉPA REG NO. 28293-271 July 8, 2013 - notif

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID				
If on skin or	• Take off contaminated clothing.			
clothing	• Rinse skin immediately with plenty of water for 15-20 minutes.			
	• Call a poison control center or doctor for treatment advice.			
Have the product container or label with you when calling a poison control center or				
doctor, or going for treatment. [You may also contact 1-800-222-1222 (or specific				
company number	r) for emergency medical treatment information.]			

Environmental Hazards

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions

Do not allow adults, children or pets to enter the treated area until vapors, mists and aerosols have dispersed and the treated area has been thoroughly ventilated. Do not spray animals. Do not water the treated area to the point of run-off. Do not make applications during rain. All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- 1) Treatment to soil or vegetation around structures;
- 2) Applications to lawns, turf, and other vegetation;

3) Applications to building foundations, up to a maximum height of 3 feet. Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches, and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application. Not for use in outdoor residential misting systems.

This product kills troublesome outdoor insects which destroy lawns, roses, flowers, trees and shrubs: Ants, Bagworms, Aphids, Borers, Armyworms, Chinch Bugs, Inchworms, Spider mites, Green Fruitworms, Beetles, Fleas, Cicadas, Thrips, Leafhoppers, Cutworms, Sod webworms, Leafrollers, Whiteflies, Moths, Weevils.

EPA REG NO. 28293-271 July 8, 2013 - notif EPA EST. NO. ______ [Subscript used is first letter of run code on container.] Page 3 of 14 Pages Also, kills Alfalfa loopers, Brown dog ticks Budworms, Cabbageworms, Citrus black flies, Cluster flies, Codling moths, Corn earworms, Crickets, Earwigs, Elm spanworms, Fall cankerworms, Gnats, Grasshoppers, House flies, Japanese beetles, Leafminers, Lone star ticks, Lygus bugs, Mealybugs, Millipedes, Mole crickets, Mosquitoes, Navy orangeworms, Scales, Sowbugs (pillbugs), Spring cankerworms, Tent caterpillars, Ticks including Deer tick (which may carry Lyme disease) Tobacco budworms and many others.

THREE EASY STEPS TO MIX AND APPLY:

Step #1: For lawns and vegetables, measure the area to be sprayed. Calculate square feet by multiplying length by width.

Step #2: Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.

Step #3: Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use. Make applications when insects first appear. Repeat applications as directed. See directions for complete insect list.

Use dilution rates as indicated. 1 fl. oz. = 6 tsp. or 2 Tbsp; 8 fl. oz. = $\frac{1}{2}$ pint or 1 cup. Food utensils such as teaspoons or measuring cups should not be used for food purposes after use with insecticides.

NUISANCE PESTS IN OUTDOOR AREAS

TREATMENT	PESTS	REMARKS [How to Use]	USE RATE
AREAS	[Controlled]	[General Directions]	
Outside surfaces Of buildings	Ants (including foraging fire ants), Chiggers, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs), Mosquitoes	To help prevent infestation of buildings, treat the building foundation to a height of 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to	6 2/3 fl. oz. in 1 quart of water, treating 400 linear feet with a 6- inch spray band
Other outdoor areas	Ants (including foraging fire ants), Chiggers, Crickets, Earwigs, Fleas, Lone star ticks, Ticks, including Deer tick (which may carry Lyme disease) Cluster flies, Flying moths, Gnats, House Flies, Mosquitoes Tent caterpillars	 maintain effectiveness. For treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For Ants, thoroughly wet hills and runways. Repeat application as reinfestation occurs. For use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Repeat as necessary. Application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent, however, they should die within several hours. 	

EPA REG NO. 28293-271 July 8, 2013 - notif

WOOD-INFESTING INSECTS INCLUDING TERMITES** (OUTDOOR USE):

WOOD INFESTING INSECTS INCLUDING TERMITES: For Beetles (Furniture, Deathwatch, False powderpost, Longhorned, Powderpost), Carpenter ants (and other wood-infesting ants), Carpenter bees and Termites (for localized control only) (including dry-wood termites):

Treat outside surfaces of building, porches, wooden decks and patios, wooden fences, window frames, doorways, foundations, eaves, patios, garages, garage areas and other areas where you may find these pests.

Mix 10 fl. oz. with 1 gallon of water to treat 100 sq. ft. Spray infested surfaces until thoroughly wet, avoiding dripping and runoff. Apply evenly on wood surfaces. To treat small areas, apply by brushing or spraying the diluted spray. For large areas, use coarse, low-pressure spray. Repeat treatment as needed to maintain effectiveness. Do not spray into electrical outlets or utilities.

Spray infested areas and areas likely to be infested, such as wood surfaces, voids and channels in damaged wood, spaces between wooden parts of a structure, and between wood and foundations where infestation is likely to occur, also around fence posts, wood piles (do not use on firewood) and tree stumps.

Suitable protective clothing should be worn in confined or overhead areas to prevent exposure to eyes and skin. The use of gloves and footwear, a long-sleeved shirt and long pants or coveralls are recommended. Do not permit contact with treated surfaces until spray has dried. Treated confined areas, such as detached garages, under porches and decks, should be ventilated during and immediately following application before the areas are re-occupied.

For carpenter ants, spray around doors and windows and other places where ants may enter or crawl and hide. Also treat cracks and crevices or openings into wall voids or other areas where these ants may be present.

For carpenter bees, spray or inject into holes and tunnels already made by the bees with a tank sprayer, and spray areas where they are beginning to build tunnels into the wood.

For termites** (including dry-wood termites): treat as for other wood-infesting insects. This product will kill workers and winged reproductive forms of termites in the treated areas. Re-treat as needed to maintain effectiveness. For localized control only. For active infestation get a professional inspection.

** USE OF THIS TYPE OF PRODUCT TO CONTROL (KILL) TERMITES SHOULD NOT BE CONSIDERED A SUBSTITUTE FOR SOIL TREATMENT OF SUBTERRANEAN TERMITES DONE BY A PROFESSIONAL PEST CONTROL OPERATOR.

		AE LAWNS	
TREATMENT	PESTS [Controlled]	REMARKS [How to Use]	USE RATE
AREAS [Plants]		[General Directions]	
Bent, Bermuda, Bluegrass, Dichondra, Fescue, Irish Moss, Merion, St. Augustine	Armyworms, Brown dog ticks, Chiggers, Chinch bugs, Cutworms, Fleas, Japanese beetle grubs, Mole crickets, Mosquitoes, Sod webworms, Ticks, including Deer Tick (which may carry Lyme disease)	Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For heavy infestations, repeat application after 2 weeks.	6 fl. oz. in 10 gals. Of water to cover 1,000 sq. ft.
	Ants (including foraging fire ants), Crickets, Grasshoppers		2 fl. oz. in 3 gallons of water to cover 1,000 sq. ft.
Bent, Bermuda, Bluegrass, Dichondra, Fescue, Irish Moss, Merion, St. Augustine	Fire ants (mound treatment)	Apply 1 gallon of solution as a gentle rain to each fire ant mound using a sprinkler can. Thoroughly wet the mound and surrounding area to a 4 foot diameter. For best results, apply in cool weather, 65°-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity, and treat all colonies which may not as yet have constructed a mound. Larger quantities of solution can be prepared for multiple mound treatments. Equipment to deliver the diluted product as a gentle rain at the rate of 1 gallon per mound is still required. Do not use equipment that produces a pressurized spray as pressurized sprays may disturb fire ants and cause migration, reducing product effectiveness.	6 fl. oz. in 1 gallon of water to treat one mound

HOME LAWNS

(

EPA REG NO. 28293-271 July 8, 2013 - notif EPA EST. NO. ______ [Subscript used is first letter of run code on container.] Page 7 of 14 Pages

TREATMENT	PESTS [Controlled]	REMARKS [How to Use]	USE RATE		
AREAS [Plants]		[General Directions]			
Arizona cypress, Azalea,	Aphids, Bagworms,	Apply as a thorough spray, wetting	1 fl. oz. in 1		
Birch, Cherry, Non-	Cicadas, Exposed	leaves and branches to the dripping	gallon of water		
bearing citrus, Conifers,	thrips, Fall	point. Try to penetrate dense foliage.			
Elm, English ivy,	cankerworms,	Spray in the late afternoon or evening,	· · ·		
Euonymus, Weeping fig,	Inchworms,	when the temperature ranges from 50°			
Fir, Honeysuckle, Lilac,	Leafminers,	to 75° F and when there is little or no			
Mock-orange, Oak,	Leafrollers, Mealybugs,	wind. Spray at the first sign of			
Palm, Philodendron,	Spider mites,	insects. Repeat as necessary; use			
Pine, Poinsettia, Tulip,	Whiteflies	intervals of 4-8 days. Application can			
Poplar, Rhododendron,		be made up to the day of collection.			
Taxus					
Cherry, Honeysuckle	Fall webworms				
Conifers	Budworms				
Elm	Elm leaf beetles, Elm				
	spanworms, Fall				
	webworms				
Douglas and Grand firs	Tussock moths				
Non-bearing citrus	Citrus black flies				
Oak	Oakworms				
Pine	Pine beetles, Pine				
	moths, Pine				
· · · · ·	needleminers, Needle				
	scales				

TREES & SHRUBS

(

11/17

ROSES & FLOWERS

TREATMENT	PESTS [Controlled]	REMARKS [How to Use]	USE RATE
AREAS [Plants]		[General Directions]	
Ageratum, Aster,	Aphids, Bagworms,	Apply as a thorough spray, wetting	1 fl. oz. in 1
Azalea, Begonia, Coleus,	Cicadas, Exposed	the plants to the dripping point. Try	gallon of water
Common ninebark and	thrips, Fall	to hit underside of leaves and	
snowberry, Exacum,	cankerworms, Fall	penetrate dense foliage. Spray in the	
Gladiolus, Gold bells,	webworms, Inchworms,	late afternoon or evening, when the	
Hypoestes, Ivy, Lilac,	Leafminers,	temperature ranges from 50° to 75° F	
Marigold, Mock-orange,	Leafrollers, Mealybugs,	and when there is little or no wind.	
Nannyberry, Orchid,	Spider mites,	Spray at the first sign of insects.	
Pansy, Pea shrub,	Whiteflies	Repeat as necessary; use intervals of	
Petunia, Poinsettia,	1	4-8 days. Application can be made	
Rhododendron, Rose,		up to the day of collection.	
Snapdragon, Zinnia			

EPA EST. NO. ______ [Subscript used is first letter of run code on container.] Page 8 of 14 Pages

	FRUIT	& NUT TREES	
TREATMENT	PESTS [Controlled]	REMARKS [How to Use]	USE RATE
AREAS [Plants]		[General Directions]	
Almonds	Navel orangeworms, Peach twig borers	Apply when insects appear. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Do not make more than 2 applications per season. Can be applied up to 7 days prior to harvest.	2 fl. oz. in 1 gallon water
Apples	Green fruitworms, Leafrollers (Oblique- banded; Redbanded), Plum curculio, Rosy apple aphids, Spotted tentiform leafminers, Tarnished plant bugs, White apple leafhoppers	Apply when insects appear. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Repeat as required to maintain control. Do not make more than 3 applications. Do not apply after petal fall.	
Peaches	Borers (Lesser peachtree, Peach twig), Green fruitworms, Plum curculio, Oriental fruit moths, Tarnished plant bugs	Apply when insects appear. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.	
Pears	Pear psylia	Apply when insects appear. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Apply during the dormant through delayed dormant growth period only. Do not make more than 2 applications per season.	
Pears (summer)	Codling moths, Green fruitwoms, Pear psylla	Apply when insects appear. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Do not make more than 3 applications per season. Can be applied up to 14 days prior to harvest.	

FDIIIT & NUT TOFFS

(

EPA REG NO. 28293-271 July 8, 2013 - notif

1

EPA EST. NO. [Subscript used is first letter of run code on container.] Page 9 of 14 Pages

		GETABLES	
TREATMENT	PESTS [Controlled]	REMARKS [How to Use]	USE RATE
AREAS [Plants]		[General Directions]	
Asparagus (pre- harvest)	Asparagus beetles, Cutworms	Apply for uniform coverage as needed. Do not make more than 4 applications per crop. Can be applied up to 3 days prior to harvest.	$1\frac{1}{2}$ fl. oz. in 1 gallon of water to cover 1000 sq. ft.
Asparagus (post- harvest)	Beetles (Adult Japanese, Larval stages of asparagus), Lygus bugs, Tarnished plant bugs	Treat ferns or bush growth after spear harvest when insects are present.	
Celery	Armyworms, (beet, Fall, Southern), Cabbage loopers, Cutworms (black, granulate), Vegetable leafminers	Apply every 3-5 days or as needed. Can be applied up to 3 days prior to harvest. Do not make more than 10 applications per season.	
Broccoli, Brussels sprouts, Cabbage, Cauliflower	Beet armyworms, Cabbage aphides and loopers, Diamondback moths, Imported cabbageworms	Apply every 5 days or as needed. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts, and Broccoli and 10 applications on Cabbage.	1 fl. oz. in 1 gallon of water to cover 1,000 sq. ft.
Eggplant	Colorado Potato beetles, Vegetable leafminers	Apply for uniform coverage as needed. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest.	3 fl. oz. in 1 gallon of water to cover 1,000 sq. ft.
Horseradish (foliar application)	Imported crucifer weevils	Apply as needed. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.	
Head lettuce	Corn earworms, Beet armyworms, Loopers (Alfalfa, Cabbage), Tobacco budworms	Apply every 5-10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest.	2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft.
Potatoes	Beetles (Colorade potato, Potato flea), Beet armyworms, Cabbage loopers, Cutworms, European corn borers, Leafhoppers (Aster, Potato), Potato aphids, Potato psyllids, Potato tuberworms, Tarnished plant bugs	Apply as needed. Do not make more than 12 applications. Can be applied up to 7 days prior to harvest.	1 ¹ / ₂ fl. oz. in 1 gallon of water to cover 1,000 sq. ft.
Peppers	Cabbage loopers, Corn earworms, European corn borers, Flea beetles, Pepper weevils, Vegetable leafminers	Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest.	2 fl. oz. in 1 gallon of water to cover 1,000 sq. ft.

VEGETABLES

(

.

.

(

EPA REG NO. 28293-271 July 8, 2013 - notif EPA EST. NO. _____ [Subscript used is first letter of run code on container.] Page 10 of 14 Pages

Spinach	Beet armyworms, Cabbage loopers, Corn earworms, Cutworms, European corn borers, Green cloverworms, Leafhoppers	Apply for uniform coverage when needed, but not more than 7 applications per season. Can be applied up to 7 days prior to harvest.	1 ¹ / ₂ fl. oz in 1 gallon of water to cover 1,000 sq. ft.
Sweet Corn	Armyworms (Fall, Southern), Corn earworms, European corn borers	Apply every 5 days and when needed, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.	
Tomatoes (Do not apply on Cherry Tomatoes and varieties yielding tomatoes less than one inch in diameter)	Armyworms, (Beet, Southern), Cabbage loopers, Colorado potato beetles, Granulate cutworms, Hornworms, Tomato Fruitworms, Tomato pinworms	Apply as needed. Do not make more than 6 applications per season. Can be applied up to day of harvest.	1 ¹ / ₂ fl. oz in 1 gallon of water to cover 1,000 sq. ft.

14/17

ORNAMENTAL HOUSE PLANTS

INDOORS: To protect plants in home greenhouses, lathhouses, and indoor landscapes

TREATMENT	PESTS [Controlled]	REMARKS [How to Use]	USE RATE
AREAS [Plants]		[General Directions]	
Ageratum, Ardisia,	Aphids, Armyworms,	Contact and residual spray. Use	1 fl. oz. in 1
Aster, Azalea, Baby's	Corn earworms,	sufficient spray to obtain full	gallon of water
breath, Begonia,	Exposed thrips, Fall	coverage of all foliage. Adjust spray	
Birdsnest fern, Bleeding	cankerworms,	volume used to plant size. Spray	
heart, Carnation,	Inchworms, Japanese	directly on insects. If application is to	
Chrysanthemum, Coleus,	beetles, Leafminers,	be made indoors, remove infested	
Crown of thorns,	Leafrollers, Mealybugs,	plants from living areas prior to	
Cockscomb, Cyclamen,	Scales, Spider mites,	application. Spray plants in an area	
Dracaena, Dumbcane,	Spring cankerworms,	not likely to be occupied by people or	
English ivy, Fuchsia,	Whiteflies	pets (particularly fish). When leaves	
Gladiolus, Grape ivy,		dry, plants can be returned to original	
Hypoestes, Ivy, Juniper,		location. Do not use more product	
Lily, Marigold, Orchid,		than recommended, as this may cause	
Palm, Pansy, Peperomia,	,	some plant injury. Use with care on	
Petunia, Philodendron,		ferns, palms and fuchsia. As with all	
Piggy-back plant,		plants listed above, especially	
Poinsettia, Portulaca,		carnations, chrysanthemums and	
Pothos, Prayer plant,		roses, there exists a wide variety of	
Purple passion, Rabbits		species and cultivars, and therefore, a	
foot fern, Rose,		wide variety of susceptibility to	
Schefflera, Snapdragon,		injury. To ascertain possible	
Snake plant, Statice,		phytotoxicity, spray a few plants as	
Velvet plant, Verbena,		described above, wait 24 hours, and	
Zinnia	·	check for possible injury. Repeat	
Chrysanthemums	Armyworms,	applications may be necessary to	
	Cabbageworms,	control citrus blackflies on non-	
	Cutworms, Leafrollers	bearing citrus and adult Japanese	
Velvet plants	Fungus gnats	beetles on roses.	

EPA REG NO. 28293-271 July 8, 2013 - notif

÷ 11

i e subs

EPA EST. NO. [Subscript used is first letter of run code on container.] Page 11 of 14 Pages HOSE END SPRAYER: Easy to Use. For Application to Lawns, Outdoor Ornamentals, Outside Surfaces of Buildings, and Other Outdoor Residential Areas. 32 oz. Treats Up to 5,000 Square Feet

- 1. Make sure Knob is twisted fully clockwise to OFF position. Shake well before attaching to hose.
- 2. Turn Product Control Button clockwise until the flat portion is flush with the lock tab, then push button all the way in. Avoid squeezing bottle.
- 3. Turn water on at faucet, aim nozzle toward lawn and turn Knob counterclockwise to begin spraying.
- 4. When finished push Product Control Button to OFF position from opposite side. Return Knob and faucet to OFF position. Discharge water pressure by turning Knob ON and OFF again. Turn Product Control Button counterclockwise away from lock tab to prevent accidental discharge. Store out of reach of children.

(Alternate Hose End Sprayer Directions – Used in accordance with type of Hose End packaged) HOSE END SPRAYER: Easy to Use. For Application to Lawns, Outdoor Ornamentals, Outside Surfaces of Buildings, and Other Outdoor Residential Areas.

32 oz. Treats Up to 5,000 Square Feet

[Snap off tab on top of sprayer before first use.] [Editor Note: when using K1 hose end]

- 1. Make sure knob Control lever is in the OFF position. Shake well and attach to hose.
- 2. Turn water on at faucet, aim nozzle toward lawn and turn knob Control lever to ON to begin spraying.
- 3. When finished turn Control lever to OFF position. Turn faucet to OFF Position. Discharge residual water pressure by turning ON and OFF again. Store out of reach of children.

(Alternate Hose End Sprayer Directions – Use in accordance with type of Hose End packaged) HOSE END SPRAYER: Easy to Use. For Application to Lawns, Outdoor Ornamentals, Outside Surfaces of Buildings, and Other Outdoor Residential Areas.

32 oz. Treats Up to 5,000 Square Feet

INSTRUCTIONS

- 1. Make sure [white] Water [Control] Valve [on top of bottle] is in the OFF or back position and the red Product Mixing Valve is [pointed] in the CLOSED or upright position.
- 2. Shake the product well and firmly attach to garden hose.
- 3. Turn water on at faucet and aim nozzle toward lawn. To begin water flow, turn [white] Water Valve ON by pushing forward. To begin spraying mixture, rotate [red] Product Mixing Valve OPEN by turning it ¹/₄ of a turn counter clockwise.
- 4. Walk at a steady pace spraying in a side to side arc slightly overlapping treated areas. [Refer to graduated side of container to guide coverage.]
- 5. To stop spraying, turn the [red] Product Mixing Valve to the CLOSED or upright position and return the [white] Water Valve to the OFF or back position.
- 6. For product storage, both valves should be in the CLOSED/OFF position.

(Editors Note: The following two sets of alternate hose end sprayer directions are for specific customer packaging and used only be that customer on their final printed labels.)

EPA REG NO. 28293-271 July 8, 2013 - notif

la cara a ana a ana an i

.

EPA EST. NO. ______ [Subscript used is first letter of run code on container.] Page 12 of 14 Pages (Alternate Hose End Sprayer Directions – use in accordance with type of Hose End packaged) [FOR USE WITH THE Lock 'N Spray® SYSTEM

Lock 'N Spray makes lawn and garden care easy. Simply attach this cartridge to the Lock 'N Spray sprayer (sold separately). Eliminates mixing, measuring and messy clean up, Automatically mixes the right amount of product and water.]

HOW TO USE THE Lock	'N Spray® SYSTEM		
CONNECT	Connect sprayer to hose.		
	Attach cartridge to sprayer. Turn cartridge until it clicks		
	into place.		
SPRAY .	To begin spraying, rotate sprayer nozzle tip to direct spray up or down, or remove for a 20 to 25 ft. jet spray. Turn on water and squeeze trigger.		
FINISH	• To stop spraying, release trigger, unlock cartridge and remove from sprayer. (No need to turn off water.)		
	• To clean sprayer, flush with water, squeezing and releasing trigger in 2 short bursts. Turn off water.		
	• To disconnect sprayer, relieve water pressure by squeezing trigger.		
Cartridge reseals itself auto	omatically – No spills, no leaks!		

(Alternate Hose End Sprayer Directions – Use in accordance with type of hose End packaged) [HOW TO USE THE Ready-Spray® SYSTEM

CONNECT:

- 1. Shake well before using.
- 2. Connect sprayer to hose. Be sure larger green ON/OFF switch located at hose end is set to "OFF"
- 3. Turn on water.
- 4. Activate smaller green switch at sprayer opening to "OPEN" position. (Coin or key may be needed to activate switch.)

SPRAY:

- 1. To begin spraying, point nozzle in the direction you want to spray.
- 2. Turn larger green ON/OFF switch at hose end counterclockwise to on position.
- 3. Spray evenly over measured area.

FINISH:

- 1. To stop spraying, turn larger green ON/OFF switch clockwise to "OFF".
- 2. Deactivate smaller green switch at sprayer opening to "CLOSED" position.
- 3. Turn off water.
- 4. Relieve water pressure by turning larger green ON/OFF switch counterclockwise to on position until water slows to a drip. Then, turn same ON/OFF switch back to "OFF".
- 5. Disconnect sprayer from hose.

EPA REG NO. 28293-271 July 8, 2013 - notif EPA EST. NO. ______ [Subscript used is first letter of run code on container.] Page 13 of 14 Pages (Alternate Hose End Sprayer Directions – Use in accordance with type of Hose End packaged) HOW TO USE: Always keep nozzle pointed away from people and pets.

- 1) Shake well before using. Connect sprayer to hose. Be sure the larger ON/OFF dial located at the hose end is set to "OFF" (tab is vertical when "OFF").
- 2) Turn on water at faucet.
- 3) Turn the smaller, star shaped button so that the flat side is aligned with the lock tab and push to the "OPEN" position.
- 4) To begin spraying, turn the larger ON/OFF dial at the hose end counterclockwise to the "ON" position.
- 5) To STOP spraying, turn the larger ON/OFF dial to the "OFF" position. Push the star shaped button to the "CLOSED" position. Twist the star shaped button so that the flat side is no longer aligned with the lock tab. Turn water off at faucet. To relieve pressure, turn the larger ON/OFF dial to the "ON" position then back to the "OFF" position. Disconnect sprayer from hose.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry area away from heat or open flame.

PESTICIDE DISPOSAL: *If partly filled* – Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: *If empty* – Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

PHAETON CORPORATION

P.O. Box 290, Madison, GA 30650

(Lot no. added at production)

[Optional supplemental distributor website: www.enforcer.com]

EPA REG NO. 28293-271 July 8, 2013 - notif EPA EST. NO. [Subscript used is first letter of run code on container.] Page 14 of 14 Pages