

UNICORN

SYNERGIZED POUR-ON INSECTICIDE III

Kills lice and flies on cattle

For Use on Beef Cattle, Lactating and Non-lactating Dairy Cattle, and Calves as a Pour-On, Mist Spray or Back Rubber

Controls keds on sheep and lambs when applied as a pour-on

Controls flies on horses when applied as a wipe-on, pour-on or ready to use spray

Concentrated Dose

Synergized Formula For Effective Control

ACTIVE INGREDIENTS:

Permethrin* -----	7.4%
Piperonyl Butoxide Technical** -----	7.4%
INERT INGREDIENTS:*** -----	85.2%
Total:	100.0%

* Cis/trans ratio: Min. 35% (±) cis
 Max. 65% (±) trans

** Equivalent to Min. 5.92% (butylcarbityl)(6-propylpiperonyl) ether and 1.48% related compounds.

*** Contains Petroleum Distillate

Keep Out of Reach of Children


CAUTION

See Side/Back Panel for Additional Precautionary Statements

NET CONTENTS:

Manufactured by:

Unicorn Laboratories
12385 Automobile Blvd.
Clearwater, Florida 33762



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes and clothing. Avoid breathing vapor or mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling.

FIRST AID	
If on Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

Environmental Hazards

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep container sealed when not in use. Do not store near food or feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT NOTE: Concentrated formulation - Use only with an accurate measuring container or applicator. **EXCEPT FOR BACK RUBBER/READY TO USE:** No dilution necessary. Apply as a pour-on, mist spray or back rubber on beef cattle, lactating and non-lactating dairy cattle and calves; as a pour-on to sheep and lambs; and as a wipe-on, pour-on or ready to use spray to horses.

APPLY TO	TARGET INSECTS	APPLICATION INSTRUCTIONS
Lactating and Non-Lactating Dairy Cattle, Beef Cattle and Calves	Lice, Horn Flies, Face Flies. Aids in control of Horse Flies, Stable Flies, House Flies, Mosquitoes and Black Flies. Control Horn Flies up to 6 weeks after application.	<p>DOSAGE: Apply 1.5 ml per 100 lbs. body weight of animal up to a maximum of 15 ml for any one animal when applied as pour-on or spray. To control heavy horn fly infestations, this product can be applied at a rate of 2.0 ml per 100 lbs. body weight, up to a maximum of 20 ml for any one animal.</p> <p>POUR-ON: Pour correct dose along back and down face.</p> <p>READY-TO-USE SPRAY: Use undiluted in a mist sprayer to apply correct dose. Apply directly to neck, face, back, legs and ears.</p> <p>BACK RUBBER USE: Mix 64 ml (2.1 fl.oz.) per gallon of mineral oil. Keep rubbing device charged. Results improved by daily forced use.</p>
Sheep and Lambs	Sheep Keds and Lice.	<p>POUR-ON: Pour along back. Apply 1 ml per 50 lb. body weight of animal, up to a maximum of 12 ml for any one animal. For optimum control, all animals in the flock should be treated after shearing.</p>
Horses and Foals	Stable Flies, Horn Flies, Face Flies, Deer Flies, House Flies, Eye Gnats, Lice and Ticks. Aids in the control of Horse Flies, Mosquitoes and Black Flies.	<p>WIPE-ON: Apply 8 ml to 16 ml as a wipe on. Dampen an applicator mitt, cloth or toweling (turkish) with product. Rub over hair with special attention to the legs, shoulders, neck and facial areas where flies tend to congregate. Be especially careful not to get in horse's eyes.</p> <p>READY-TO-USE SPRAY: Use this product after riding or exercise. Use undiluted in a mist sprayer, applying 8 ml to 16 ml per animal. Apply directly onto attached ticks.</p>
Horses	Stable Flies, Horn Flies, Face Flies, Deer Flies, House Flies, Eye Gnats, Lice and Ticks. Aids in the control of Horse Flies, Mosquitoes and Black Flies.	<p>POUR-ON: After riding or exercise, pour 8 ml to 16 ml of product along the back and down the face of MATURE horses, being careful to avoid the eyes. Do not use this method of application on foals.</p>
PREMISES – In and around Horse, Beef, Dairy, Swine, Sheep and Poultry Premises, Animal Hospital Pens and Kennels and "outside" Meat Processing Premises.	House Flies, Stable Flies, Face Flies, Gnats, Mosquitoes, Black Flies, Fleas, Little House Flies, (<i>Fannia spp.</i>). Aids in control of Cockroaches, Ants, Spiders, and Crickets.	<p>For use as a Ready-To-Use Spot Spray or Premise Spray: Use undiluted in a mist sprayer. Apply directly to surface to leave a residual insecticidal coating, paying particular attention to areas where insects crawl or alight. 500 ml will treat approximately 7,200 square feet.</p>

4 8 4

For cattle and sheep, repeat treatment as needed, but no more often than once every two weeks. For optimum lice control, two treatments at a 14-day interval are recommended.

Special Notes: This product is not effective in controlling cattle grubs. This is an oil-base, ready-to-use product, that may leave an oily appearance on the hair coat of some animals. This product should be used in an integrated pest management system which may involve repeat treatments and the use of other pest control practices.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.