


PM 3

28,293 - 294

3/9/98

1/6

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <u> X </u> Registration <u> </u> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number:	Date of Issuance:
	28293-294	MAR 9 1998
	Term of Issuance: Conditional	
Name of Pesticide Product:		
Unicorn .20% Permethrin WB II		
Name and Address of Registrant (include ZIP Code):		
Unicorn Laboratories 12385 Automobile Blvd. Clearwater, FL 33762		
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.		
Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.		
<p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Make the labeling changes listed below before you release the product for shipment: <ol style="list-style-type: none"> a. Add the phrase "EPA Registration No. 28293-294". b. On the front panel delete "Provides 60 Day (2 Months) Control of Fleas" or cite efficacy data to support this claim. c. Under Statement of Practical Treatment/First Aid add "IF INHALED: Remove effected person to fresh air. Apply artificial respiration if indicated." 		
Signature of Approving Official:		Date:

2/
16

d. Under Enviromental Hazards Change "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark." to "Do not apply directly to water."

e. Under INDOOR APPLICATION add "Do not apply to classrooms while in use. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed. Not for use in commercial food/feed handling establishments, restaurants or other places where food/feed is commercially prepared or processed."

f. Under ON DOGS after "Reapply every two (2) weeks as needed." add "Use protective gloves or mitts when applying product."

g. Under STORAGE AND DISPOSAL add "STORAGE: Store in a cool dry area away from heat or open flame. Store above 32 degrees fahrenheit."

h. Under directions for control of garden insects add "Not for Use on Plants being grown for sale or other commercial use."

i. Under Directions for Use modify the second sentence to read "Affix trigger pump or other type spray head to container for application by homeowners indoors."

3. Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

3/6

UNICORN

.20% PERMETHRIN WB II

- ◆ For Domestic Use both Indoors and Outdoors ✓
 - ◆ A Ready-To-Use Stable Emulsion Insecticide ✓
 - ◆ Provides 60 Day (2 Months) Control of Cockroaches ✓
 - ◆ Provides 60 Day (2 Months) Control of Fleas No
 - ◆ For Control of Fleas, Brown Dog Ticks and Lice on Premises and on Dogs ✓
 - ◆ Controls Fleas and Ticks on Dogs for 35 Days
 - ◆ Aids in Control of Ticks on Dogs for an additional 2 Weeks ✓
 - ◆ For Control of Pests on Livestock ✓
- ◆ Quick Kill Formula ◆ Long Lasting Control

ACTIVE INGREDIENT:

Permethrin * 0.20%

INERT INGREDIENTS: 99.80%

Total: 100.00%

* (3-phenoxyphenyl) methyl (±) cis/trans 3-(2,2-dichloroethenyl)
2,2-dimethylcyclopropanecarboxylate.

Cis/trans ratio: Min. 35% (±) cis
 Max. 65% (±) trans

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side/back panel for Additional Precautionary Statements.

NET CONTENTS:

Manufactured by:

Unicorn Laboratories
12385 Automobile Blvd.
Clearwater, FL 33762
(813) 578-4545

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR 9 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for use as insecticide
registered under EPA Reg. No.

28293-294

EPA REG. NO. 28293-~~XXXXX~~ *FOU*

- 9/16
- READ ENTIRE LABEL BEFORE EACH USE.
 - FOR PET USE - USE ONLY ON DOGS.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry. Sensitivities may occur after using **ANY** pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

of inhalation

Statement of Practical Treatment/ First Aid

- IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.
- IF ON SKIN:** Remove contaminated clothing and wash affected areas with soap and warm water.
- IF SWALLOWED:** Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

Environmental Hazards

This product is extremely toxic to fish. Do not apply directly to water, ~~to areas where surface water is present or to intertidal areas below the mean high water mark.~~ Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

SHAKE WELL BEFORE USING

INDOOR APPLICATION

For spot application only. Do not use as a space spray indoors. *Do not etc.*

Surface Spraying:

To control **Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadollos:** Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

5/6

To control **Carpet Beetles**: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control **Bedbugs**: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control **Fleas** (adults and larvae) and **Ticks** (adults and larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

To control **Clothes Moths**: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible.

Apply spray liberally to empty chests, closets, bureaus and other storages, directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To kill **Lice** and **Louse Eggs**: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

ON LIVESTOCK

To protect cattle (beef and dairy), goats, sheep, hogs and horses from **Horn Flies**, **Face Flies**, **House Flies**, **Mosquitoes** and **Gnats**: Apply a light mist sufficient to wet the surface of the hair. For face flies, Spray face and head, but do not spray into eyes.

To control **Stable Flies**, **Horse Flies** and **Deer Flies**: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control **Blood-sucking Lice**: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control **Poultry Lice**: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To control **Bedbugs** and **Mites** in poultry houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS

To control **Fleas**, **Ticks** and **Lice**: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin.

9/16

Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Reapply every two (2) weeks as needed.

NOTE: Do not use on puppies under twelve (12) weeks of age. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication.

DO NOT USE ON CATS OR KITTENS.

OUTDOOR APPLICATION

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

FOR CONTROL OF GARDEN INSECTS:

For use on roses, dahlia, asters and other ornamentals to kill **Japanese Beetles** hit by spray as well as **Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites and Leafminers**. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against **Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms**.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

DISPOSAL: Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.