


Martin's

SEVIN®

Brand Carbaryl Insecticide

5% Dust

Ready to Use
Vegetables -- Gardens -- Lawns
Poultry -- Horses -- Dogs -- Cats

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate).....	5%
INERT INGREDIENTS.....	95%
	100%

CAUTION**KEEP OUT OF REACH OF CHILDREN**

EPA REG. NO. 28293-301 (11/1/99 AMEND)

EPA EST. NO. 299-TX-1

READ ENTIRE LABEL BEFORE EACH USE
USE ONLY ON POULTRY, HORSES, DOGS, AND CATS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS – CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS: This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This pesticide is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

FIRST AID: If swallowed, induce vomiting and seek medical attention immediately. If inhaled, remove victim to fresh air. Apply artificial respiration if indicated. If in eyes, flush eyes with plenty of water. Call physician if irritation persists. If on skin, wash skin thoroughly with soap and water.

NOTE TO PHYSICIANS AND VETERINARIANS: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is an antidote.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USER SAFETY REQUIREMENTS FOR GARDENS

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product. Change clothing as soon as possible after use. Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet. Do not allow children or pets to contact treated area until dust has settled.

Application: Begin application when insect damage is first observed. Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week.

Vegetables: Corn, Eggplant, Pepper, Tomato – To control Corn Earworms, European corn Borers, Sap Beetles, Armyworms, Tomato Fruitworms, Flea Beetles, Leafhoppers, Tomato Hornworms, and Stink Bugs, dust upper and lower leaf surfaces thoroughly. When Used on Sweet Corn for control of Corn Earworms, application of this product to the tassel region of corn during pollen shed period will seriously reduce bee population. Do not harvest sooner than 2 days after application to corn or sooner than 3 days after application to eggplant, pepper or tomato.

Beans – To control Mexican Bean Beetles, Bean Leaf Beetles, Cucumber Beetles, Leafhoppers, flea Beetles, Japanese Beetles, Corn Earworms, Lygus Bugs and Lima Bean Pod Borers, dust upper and lower leaf surfaces thoroughly. Do not harvest sooner than 3 days after application to fresh beans or sooner than 21 days after application to dried beans.

Cucumber, Summer Squash – To control Pickleworms, Melonworms, Cucumber Beetles, Squash Bugs and Flea Beetles, dust upper and lower leaf surfaces thoroughly. Do not harvest sooner than 3 days after application.

Potato – To control Colorado Potato Beetles, Leafhoppers, Flea Beetles and Lygus Bugs, dust upper and lower leaf surfaces thoroughly. Do not harvest sooner than 7 days after application.

SUBSEQUENT PLANTINGS OF FOOD AND FEED CROPS IN TREATED SOIL MUST BE MADE WITH CROPS ON THIS OR OTHER CARBARYL LABELS.

Lawns – To control Chinch Bugs, Armyworms, Ants, Fleas, Sod Webworms, Lawn Moths, Millipedes and Earwigs, apply 1 lb. to 200 sq. ft. For best results, apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 or 3 weeks if necessary.

Poultry: Litter Treatment – Apply 1 lb. to 40 sq. ft. to control Northern Fowl Mites, Chicken Mites, Lice, Fleas and Bed Bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs. Do not house birds in treated houses within 7 days of slaughter.

USER SAFETY REQUIREMENTS FOR PETS

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product. Change clothing as soon as possible after use. Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.

Dogs & Cats: To control Brown Dog Ticks and Fleas, use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work down to the skin. This product should not be applied by children. Do not use on pregnant dogs or cats. Do not use on kittens or puppies under 12 weeks of age. Dust bedding and doghouse using a shaker can or other dust applicator. Do not repeat treatment for 7 days. Do not use in homes. DO NOT APPLY TO RUGS AND CARPETS AS STAINING MAY OCCUR. Consult a veterinarian before using this product on debilitated, aged, or animals on medication.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. However, flea and tick collars may be immediately replaced.

Horses: For control of Hornflies and suppression of Stable Flies – Wearing household latex or rubber gloves, rub or brush 2 oz. of powder per animal into the hair on back and neck. Stable Fly treatment of legs of animals is supplemental to premise treatment for Stable Fly control. Do not repeat application more than once a week. This product should not be applied by children. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing, or animals on medication.

STORAGE & DISPOSAL: **Storage** – Do not contaminate water, food or feed by storage or disposal. Store in original container in a dry place until used. Store out of reach of children, pets and animals. **Disposal** – Do not reuse empty container. Completely empty can, then dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

SEVIN is a registered trademark of Rhone-Poulenc for carbaryl insecticide.

MARTIN'S is a registered trademark of C.J. Martin Company

**Unicorn Laboratories
12385 Automobile Blvd.
Clearwater, FL 33762
(727) 578-4545**

