

### Unicorn .25% Permethrin Granules

- [KILLS INSECTS IN LAWNS]
- [LAWN & GARDEN INSECT CONTROL]
- [TREATS UP TO 5,000 10,000 15,000 SQ.FT.]
- [PROTECTS YOUR HOME FROM INVADING PESTS]
- [10 POUNDS TREATS 5,000 SQ.FT.]
- [CONTROLS SURFACE AND FOLIAGE FEEDING PESTS]
- [FOR GRASS AND DICHONDRA LAWNS]
- [CONTAINS SYNTHETIC PYRETHROID]
- [FOR OUTDOOR USE ON HOME LAWNS]
- [CONTAINS PERMETHRIN]
- [READY TO USE]
- [KILLS: Ants, Armyworms (Inc. Fall, Yellow-striped, etc.), Bear Tick, Bermudagrass mites, Billbugs (adult), Black Legged Tick, Boxelder Bugs, Brown Dog Ticks, Centipedes, Chiggers, Chinch Bugs, Clover mites, Cockroaches, Crickets, Cutworms, Darkling Ground Beetles, Deer Tick (Ixodes dammini), Digger wasps, Dung Beetles (Ataenius Spretulus), Essex Skipper, European Crane Fly Larvae, Field Crickets, Fiery Skipper, Fire Ants, Flea Beetle (In dichondra), Fleas, Grasshopper, Grubs, Hyperodes (Turfgrass Weevil), Japanese Beetle (adult), Lawn Moths, Leafhoppers, Lone Star Tick, Scorpions, Silverfish & Firebrats, Sod Webworms, Sowbugs, Spiders (Inc. Black Widow), Spittlebugs, Springtails, Ticks, Vegetable Weevils (In dichondra), Western Black Legged Tick (Ixodes pacificus)]
- [PROVIDES CONTACT AND RESIDUAL CONTROL FOR UP TO 4 WEEKS]
- [CONTAINS PERMETHRIN, A LONG LASTING INSECTICIDE WHICH PROVIDES CONTACT AND RESIDUAL CONTROL OF OUTDOOR INSECT PESTS.]
- [KILLS ANTS, FLEAS, BROWN DOG TICKS, CRICKETS, CHINCH BUGS, CUTWORMS, BILLBUGS, SOD WEBWORMS, GRUBS AND OTHER LISTED INSECTS.]
- [LONG LASTING INSECT CONTROL]

**ACTIVE INGREDIENT:**

\*Permethrin.....0.25%

INERT INGREDIENTS: .....99.75%

TOTAL .....100.00%

\*(3-phenoxyphenyl) methyl (±) cis/trans 3-(2,2-dichloroethenyl) -2,2-dimethyl cyclopropanecarboxylate  
Cis/trans ratio: Maximum 55% (±) cis and minimum 45% (±) trans


KEEP OUT OF REACH OF CHILDREN

### CAUTION

See Side/Back Panels (Label) for Additional Precautionary Statements.

Net Contents:

Manufactured by:  
Unicorn Laboratories  
12385 Automobile Blvd  
Clearwater, FL 33762



<b>FIRST AID</b>	
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Avoid contamination of food or feed. Utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticides.

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish. Do not apply directly to water. Drifts and runoffs from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Unicorn .25% Permethrin Granules is recommended for the control of lawn insects, fleas, and ticks outdoors in residential areas. Repeated applications are made by broadcast treatment with a spreader to lawns, under shrubbery, and other ornamental and turf grass areas, band treatments are recommended around house foundation and spot applications are made to ant mounds and wasp nests.

**STORAGE AND DISPOSAL**

**STORAGE:** Store product in its original container in a cool, dry locked place out of reach of children.

**DISPOSAL:** Completely empty bag into application equipment. Do not reuse empty container. Wrap bag and put in trash collection.

**GENERAL USE DIRECTIONS [THREE EASY STEPS TO USE THIS PRODUCT]**

1. [Step #1] Determine size of area to be treated – measure and multiply length times width.
2. [Step #2] Determine appropriate spreader setting using chart.
3. [Step #3] Treat as directed – thorough coverage is important. Water in where indicated.

Keep children and pets off treated areas until this material is washed into turf and grass is dry.

Treatment Areas	Pests	Use Rate	How to Use
<p><b>OVERALL LAWN TREATMENT FOR GRASS AND DICHONDRA LAWNS</b></p>	<p>Ants, Armyworms (Inc. Fall, Yellow-striped, etc.), Bermudagrass mites, Boxelder Bugs, Centipedes, Chiggers, Chinch Bugs, Clover mites, Cockroaches, Crickets, Cutworms, Darkling Ground Beetles, Digger wasps, Essex Skipper, European Crane Fly Larvae, Field Crickets, Fiery Skipper, Flea Beetle (In dichondra), Grasshopper, Hyperodes (Turfgrass Weevil), Lawn Moths, Leafhoppers, Scorpions, Silverfish &amp; Firebrats, Sod Webworms, Sowbugs, Spiders (Inc. Black Widow), Spittlebugs, Springtails, Vegetable Weevils (In dichondra)</p>	<p>Treat when insects first appear using granules at the rate of 2 pounds per 1,000 sq.ft. or, for heavy lawns such as St. Augustine grass, 3 pounds per 1,000 sq.ft. using a fertilizer spreader. See spreader setting chart below.</p>	<p>Apply this product uniformly to home lawns and under shrubbery. To help obtain uniform distribution, apply one-half required amount of product in one direction, then apply the balance at a right angle. Immediately after application, water the lawn lightly.  <b>European Crane fly larvae (Leatherjackets)</b> – the best time to apply is in May or June when the insects are young.  <b>Chinch Bugs</b> – Start treatments when temperatures reach 80°F or above. A second application may be necessary in 6 weeks if insect activity continues.</p>
	<p>Billbugs, Grubs, Japanese Beetle (adult), Dung Beetles (<i>Ataenius Spretulus</i>)</p>	<p>Use granules at the rate of 2 ½ pounds per 1,000 sq.ft.</p>	<p>Water grass thoroughly after application. For <b>Billbug</b> control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. For <b>Grub</b> control, except <b>Dung Beetle</b>, treat any time between late July and early October. For <b>Dung Beetle Grub</b> control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with ¼ to ½ inch of water after application. Treat again in late summer or early fall if a second generation occurs.</p>

Treatment Areas	Pests	Use Rate	How to Use
SPOT TREATMENT	Ants nesting outdoors including: Acrobat, Argentine, Big Headed, Cornfield, Field, Harvester, Leafcutter, Little Black, Pavement and Pyramid Ants	Product should be applied to thoroughly cover the nest (1 to 2 teaspoons) and sprinkled gently with water.	If possible, it is best to try to locate nest and treat this first. Nests will usually be found in the ground but may be concealed under stones, boards, slabs or in stumps or logs. In addition to treating ant nests or mounds, entry points of buildings (used by ants) and the soil surface along the perimeter of the building should be treated.
	Digger Wasps	Apply 1 to 2 teaspoons of granules in and around each individual ground nest opening.	Make the application at dusk when the wasps are not active.
SPOT TREATMENT (continued)	Imported Fire Ants	Apply one cup over, and 2 feet around each fire ant mound.	Immediately after application, water the granules into the soil gently with a light sprinkling to avoid disturbing the colony. Use at least 1 gallon of water per mound, but no more than 2 gallons. Use of high pressure watering equipment may disturb the colony and cause migration of the ants, thus reducing the effectiveness of the treatment. Apply in cool weather (65-80°F), or in early morning or late evening for best results. Treat new mounds as they appear.
BAND (PERIMETER) TREATMENT AROUND HOUSE FOUNDATION	Ants, Boxelder Bugs, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Field Crickets, Fleas, Scorpions, Sowbugs, Spiders (Inc. Black Widow), Springtails (Collembola) and Ticks	Treat at a rate of 1 lb. per 500 sq.ft. or 2 lbs. per 1,000 sq.ft.	To help prevent infestation of buildings, apply a 5 foot band of granules next to the house foundation. Distribute granules uniformly. Water lightly after application.

**FLEA AND TICK CONTROL (OUTDOORS)**

In summer months, fleas and ticks frequently become established in lawns and pet resting areas surrounding the home. For best control of Fleas and Ticks (including but not limited to Bear Tick, Black-legged Tick, Brown Dog Tick, Deer Tick (Ixodes dammini), Lone Star Tick, Western Black-legged Tick, Wood Tick): a regular treatment program should be followed. Treat once a month when pests first appear in late spring (May-June) and continue until late summer (September-October). Fall temperatures below 50°F decrease pest activity. Eliminating weeds around the home and mowing lawns frequently reduces the hiding places for these pests.

Note: Dogs, cats, their kennel or bedding area should also be treated with a registered product. In addition, if fleas and ticks are indoors on carpets or pet resting areas, use a registered product such as (*product brand name*), ideally in combination with an Insect Growth Regulator such as (*product brand name*). It is very important that all three sites – outdoors, indoors and also the pet be treated with appropriate labeled products to achieve satisfactory control. If any of these areas is not treated, it could serve as a source of reinfestation to the entire premises.

**SOD WEBWORMS (CRAMBUS SPP)**

Sod Webworms are the light brown caterpillars ¼ to ¾ inch long, that hide in the grass during the day and feed on the grass blades at night. Adult Webworms, usually referred to as Lawn Moths (about 1 inch in length with long snout) emerge from the grass at dusk. They fly over the lawn laying eggs that later hatch into destructive caterpillars. Damage appears first as small brown spots which, if not treated, may rapidly develop into large brown areas of severely damaged or dead grass. If Lawn Moths (Sod Webworm Adults) are noticed in large numbers at dusk, or at first sign of damage to lawn, treat entire lawn evenly.

**SPREADER SETTINGS**

Spreader Make and Model	Grass or Dichondra Lawns at 2 lbs. per 1000 sq.ft.	White Grubs, Billbugs at 2 ½ lbs. per 1000 sq.ft.	St. Augustine Lawns at 3 lbs. per 1000 sq.ft.
Cyclone Broadcast	3	4	5
Greenview Drop Spred III	3	4	5
Greenview Jet Spred III	4	5	6
Republic Drop EZ530	6 1/2	7	8
Republic Broadcast EZ540	2	3	4
Scotts AG3	6	6 ½	7
Scotts SG3	4	4 ½	5
Scotts EG3	24	25	26
Scotts PF4	6	6 ½	7
Sears Drop 1927	5 1/2	6	7
Sears Broadcast 1922	5 1/2	6	7

Note: Spreader settings given are approximate and variation can occur between identical models. User should calibrate spreader by accurately weighing a few pounds into your spreader and applying to a measured area. Then adjust spreader setting if necessary to assure correct rate of application.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty or any other warranty of Merchantability or Fitness for a Particular Purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Brackets "[ ]" denote optional wording.