



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

May 9, 2025

James McFadden
Director, Regulatory Affairs
Phaeton Corp, dba Unicorn Laboratories
1501 East Woodfield Road, Suite 200 West
Schaumburg, Illinois 60173

Subject: Label Amendment - Registration Review Mitigation for Permethrin
Product Name: UNICORN MULTI-USE .25% PERMETHRIN DUST
EPA Registration Number: 28293-352
Application Date: 12/29/2020
Decision Number: 569312
Case Number: 475921

Dear James McFadden:


The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Permethrin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Carolyn Smith via email at smith.carolyn@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Jaclyn Pyne". The signature is written in a cursive, flowing style.

Jaclyn Pyne, Team Leader
Risk Management and Implementation Branch 3
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

PERMETHRIN	GROUP	3A	INSECTICIDE
------------	-------	----	-------------

UNICORN MULTI-USE .25% PERMETHRIN DUST

MASTER LABEL

SUB-LABEL A: COMMERCIAL USE

SUB-LABEL B: RESIDENTIAL USE

ACTIVE INGREDIENT:

Permethrin*0.25%

OTHER INGREDIENTS:.....99.75%

Total:100.00%

*Cis/trans ratio: Min.35% (±) cis, Max. 65% (±) trans


KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side/Back Label Panel for Additional Precautionary Statements and First Aid

NET WEIGHT: [1, 2 or 5 LB]

Phaeton Corporation
P.O. Box 290
Madison, GA 30650



EPA Reg. No. 28293-352

EPA Est. No. 58996-MO-1A; 44616-MO-1B; 44616-GA-1C

Subscript used is first letter of run code on container

May 1, 2025 - revised

Page 1 of 15 Pages

[SUB-LABEL A: COMMERCIAL USE]

PERMETHRIN	GROUP	3A	INSECTICIDE
------------	-------	----	-------------

UNICORN MULTI-USE .25% PERMETHRIN DUST

Optional Marketing Statements:

[] Indicates alternate or optional wording
For Use On [Can be used on]: Livestock
For Control of Horn Flies and Lice on Beef and Dairy Cattle
Aids in reducing Face Fly population on Beef and Dairy Cattle
For Control of Lice on Swine
For Control of Northern Fowl Mites on Poultry
For use in dust bags, shaker can or mechanical dust applicator
Easy-to-use powder duster
No unpleasant odor.

ACTIVE INGREDIENT:

Permethrin*0.25%

OTHER INGREDIENTS:.....99.75%

Total:100.00%

*Cis/trans ratio: Min.35% (±) cis, Max. 65% (±) trans

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side/Back Label Panel for Additional Precautionary Statements and First Aid

NET WEIGHT: [1, 2 or 5 LB]

READ ENTIRE LABEL BEFORE EACH USE.

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice

EPA Reg. No. 28293-352

EPA Est. No. 58996-MO-1A; 44616-MO-1B; 44616-GA-1C

Subscript used is first letter of run code on container

May 1, 2025 - revised

Page 2 of 15 Pages

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 [xxx-xxx-xxxx] for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT:

Some materials that are chemical resistant to this product are barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride, viton. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

Loaders, applicators, and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves

Wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Application with aerial or motorized ground equipment is prohibited.

USER SAFETY RECOMMENDATIONS

User should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

User should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Environmental Hazards

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

Under some conditions, it may also have a potential for transport into surface water in runoff (primarily absorbed to suspended soil particles), for several months or more after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, and areas overlying extremely shallow groundwater, with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For Indoor Use Only

RESISTANCE-MANAGEMENT

For resistance management, the Unicorn Multi-Use .25% Permethrin Dust contains Group 3A insecticides. Any insect population may contain individuals naturally resistant to the Unicorn Multi-Use .25% Permethrin Dust and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly. Appropriate resistance-management strategies should be followed. To delay insecticide resistance, take the following steps:

- Rotate the use of the Unicorn Multi-Use .25% Permethrin Dust or other Group 3A insecticides with different groups that control the same pests.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, record keeping and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist for any additional pesticide resistance-management

EPA Reg. No. 28293-352

EPA Est. No. 58996-MO-1A; 44616-MO-1B; 44616-GA-1C

Subscript used is first letter of run code on container

May 1, 2025 - revised

Page 4 of 15 Pages

and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance, contact Phaeton at 1-800-XXX-XXXX.

Restrictions

Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).. Rainfall within 24 hours after application may cause unintended runoff of pesticide application. Do not apply when the wind speed is greater than 15 mph. Do not water the treated area to the point of run-off.

Use of handheld power duster equipment is prohibited. Except when treating livestock, do not apply this product in a way that will contact workers or other persons, either directly or through drift. Except when treating livestock, only protected handlers may be in the area during application. Do not breathe dusts. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.



When used in dairy barns or facilities: close milk bulk tank lids to prevent contamination from dust and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.

Follow proper disposal procedures on this label.

Siga las indicaciones del etiquetado para el desecho apropiado del producto.

BEEF AND DAIRY CATTLE AND HORSES:

Can be used in dust bags, shaker can and mechanical dust applicator. Horn Flies, Lice, Face Flies: Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through which the animals must pass daily for water, feed, or minerals. Bags may also be placed in loafing sheds or in front of mineral feeders. For dairy cows, bags may be suspended in the exit through which the cows leave the milking barn. The bags should hang 4 to 6 inches below the back line of the cattle. For reduction of Face Flies, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted. Horn Flies, Lice - Direct Application: Apply 2 oz. of dust per animal by shaker can over the head, neck, shoulders and back and tail head.

HORSES: Dust horse for Hornflies with shaker can or dust glove along back, neck and legs.

SWINE: Lice on Swine - Direct Application: Apply only 1 ounce per head as a uniform coat to the head, shoulder and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary, but not more often than once every 10 days. In severe infestation, both animals and the bedding may be treated. Do not ship animals for slaughter within 5 days of treatment.

POULTRY: To control Northern Fowl Mites, apply at a rate of ¼ lb per 25 birds. Ensure thorough treatment of vent area.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry). Do not pour or dispose down-the-drain or sewer. Call your local solid waste Agency for local disposal options.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Completely empty can by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Phaeton Corporation
1501 E. Woodfield Rd, Ste 200 West
Schaumburg, IL 60173

[Lot number added at production]

[SUB-LABEL B: RESIDENTIAL USE]

UNICORN MULTI-USE .25% PERMETHRIN DUST

Optional Marketing Statements:

[] Indicates alternate or optional wording

For Use On: [Can be used on]

- Non-Commercial Livestock and Companion Animals
- Dogs
- Home Ornamentals
- Home Gardens

Kills [over] 60 [60+] [types of] Insects

[on Vegetables, Ornamentals, Fruits]

[Kills Bugs (Apple Aphid, Armyworm, Artichoke Plume Moth, Asparagus Beetle, Aster leafhopper, Beet Armyworm, Cabbage Aphid, Cabbage Looper, Codling Moth, Colorado Potato Beetle, Corn Earworm, Corn Rootworm (Adult), Cucumber Beetle (Adult), Cutworm, Cutworm Complex, Diamondback Moth, European Corn Borer, Fall Armyworm, Filbertworm, Flea Beetle, Fruit Fly, Granulate Cutworm, Green Fruitworm, Hornworm, Imported Cabbageworm, Japanese Beetle, Leafhopper, Leafminer, Lepidopterous Larvae, Lesser Peachtree Borer, Lygus Bug, Melon Worm, Mirid Bug, Navel Orangeworm, Obliquebanded Leafroller, Omnivorous Leafroller, Onion Maggot (Adult), Onion Thrips, Oriental Fruit Moth, Peach Twig Borer, Pear Psylla, Pepper Weevil, Pickle Worm, Plant Bug (including Lygus and Stink Bug), Plum Curculio, Potato Aphid, Potato Flea Beetle, Potato Leafhopper, Potato Psyllid, Potato Tuberworm, Redbanded Leafroller, Rindworm, Rose Chafer, Southern Armyworm, Spotted Tentiform Leafminer, Squash Bug, Squash Vine Borer, Stink Bug, Tarnished Plant Bug, Thrips, Tomato Fruitworm, Tomato Pinworm, Vegetable Leafminer, Walnut Husk Fly, White Apple Leafhopper)]

[Kills Beetles, Caterpillars & Bugs]

Controls Fleas & Ticks on Dogs [Use on Dogs to Kill Fleas, Ticks and Lice]

Effective Long Lasting Control

[For Home Gardens – Vegetables, Fruit Trees and Ornamental Plants]

[Use on Home Garden Fruits, Roses, Vegetables & Ornamental Plants]

[Product Facts: Kills Insects Beetles (including Japanese & Cucumber Beetles), Caterpillars (including Armyworm, Beet Armyworm, Cabbage Looper, Corn Earworm, Cutworm, Diamondback Moth, Fall Armyworm, Hornworm, Imported Cabbageworm, Lepidopterous Larvae, Lesser Peachtree Borer, Oriental Fruit Moth, Peach Twig Borer, Southern Armyworm, Tomato Fruitworm, Tomato Pinworm), Leafhoppers & other Garden insects [and other insects as listed] Plus Fleas & Ticks

Where to Use • In your Garden on Vegetables, Fruits & Ornamentals (including Roses) • On Dogs]

ACTIVE INGREDIENT:

Permethrin*0.25%

OTHER INGREDIENTS:99.75%

Total:100.00%

*Cis/trans ratio: Min.35% (±) cis, Max. 65% (±) trans

KEEP OUT OF REACH OF CHILDREN

EPA Reg. No. 28293-352

EPA Est. No. 58996-MO-1_A; 44616-MO-1_B; 44616-GA-1_C

Subscript used is first letter of run code on container

May 1, 2025 - revised

Page 7 of 15 Pages

CAUTION

See Side/Back Label Panel for Additional Precautionary Statements and First Aid

NET WEIGHT: [1, 2 or 5 LB]

READ ENTIRE LABEL BEFORE EACH USE.

FOR PET USE – USE ONLY ON DOGS. DO NOT USE ON CATS.

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 [xxx-xxx-xxxx] for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

All cat warning labels will be placed within a shaded box within the Precautionary Statements on the label. The coloring of the box and text will be such to provide a contrasting color to other label text. The shaded box must contain the following.

“DO NOT USE ON CATS. May be toxic or potentially fatal if applied to or ingested by cats. Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movements. If this occurs, immediate veterinary care should be provided.”

Each label must include at least one cat slash icon. This icon may be in the shaded box or in other areas on the label.

Each individual tube must be labeled with the “DO NOT USE ON CATS” statement or cat slash icon.

Environmental Hazards

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide of the treatment area. This pesticide is highly toxic to bees exposed to direct treatment

EPA Reg. No. 28293-352

EPA Est. No. 58996-MO-1_A; 44616-MO-1_B; 44616-GA-1_C

Subscript used is first letter of run code on container

May 1, 2025 - revised

Page 8 of 15 Pages

on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For Both Indoor and Outdoor Use

Restrictions

Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application. Do not apply when the wind speed is greater than 15 mph.

Do not water the treated area to the point of run-off. Use of handheld power duster equipment is prohibited. Except when applying to pets, companion animals or non-commercial livestock, do not apply this product in a way that will contact adults, children or pets, either directly or through drift. Except when applying to pets, companion animals or non-commercial livestock, do not allow adults, children, or pets to enter the treated area until dusts have settled. Do not breathe dusts.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.



Follow proper disposal procedures on this label.

Siga las indicaciones del etiquetado para el desecho apropiado del producto.

Applications to dogs and livestock must be done indoors.

ON PETS, COMPANION ANIMALS AND NON-COMMERCIAL LIVESTOCK:

Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

PETS – USE ON DOGS ONLY. DO NOT USE ON CATS.

To control Fleas, Ticks and Lice on Dogs, dust animals thoroughly, working the dust into the hair down to the skin. Pay close attention to legs and feet. Use ½ oz. on animals 20 pounds and under and 1 oz. on pets over 20 pounds. Avoid getting dust in pet's eyes. Reapply every two (2) weeks as needed. Do not use on dogs under twelve (12) weeks of age. Dust sleeping area including bedding and cracks and crevices.

CATTLE AND HORSES:

Can be used in dust bags, shaker can and mechanical dust applicator. Horn Flies, Lice, Face Flies: Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through which the animals must pass daily for water, feed, or minerals. Bags may also be placed in loafing sheds or in front on mineral feeders. For dairy [milk] cows, bags may be suspended in the exit through which the cows leave the milking barn. The bags should hang 4 to 6 inches below the back line of the cattle. For reduction of Face Flies, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted. Horn Flies, Lice - Direct Application: Apply 2 oz. of dust per animal by shaker can over the head, neck, shoulders and back and tail head.

HORSES: Dust horse for Hornflies with shaker can or dust glove along back, neck and legs.

SWINE: Lice on Swine - Direct Application: Apply only 1 ounce per head as a uniform coat to the head, should and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary, but not more often than once every 10 days. In severe infestation, both animals and the bedding may be treated. Do not slaughter animals within 5 days of treatment.

POULTRY: To control Northern Fowl Mites, apply at a rate of ¼ lb per 25 birds. Ensure thorough treatment of vent area.

HOME GARDENS: For Use on Home Gardens. Not for Use on Plants being grown for Sale or other Commercial Use.

[HOW TO APPLY]

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. [Wearing household latex or rubber gloves, open shaker side of cap and shake gently to apply. (directions to be used when packaged in an applicator container – the statement used depends upon type of container packaged.))] Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. (Many insects hide on the underside of the leaf). Apply when air is calm to avoid drift and contact with eyes and skin. Controls a wide range of insect pests.

[WHEN TO APPLY]

Use at first signs of insect infestation, preferably before damage from feeding occurs. Note the pre-harvest (time between use and harvest) interval with each crop listed.

Directions for Application

Crop	Target Pest [Insects Controlled]	REMARKS [Applications Per Season (Max)] [Days To Wait To Harvest] [as separate columns]
Cole Crops		

Broccoli	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Brussel Sprouts	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Cabbage	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.
Cauliflower	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Chinese Broccoli* *Not for use in California	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Chinese Cabbage (tight-heading varieties only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.
Collards (for use in AR, AZ, GA, IL, NC, OK, SC & TX only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Cucurbit Vegetables		
Cucurbit Vegetables*	Leafminer Squash Bug Cabbage Looper Cucumber Beetle (adult) Cutworm Melon Worm Pickle Worm Plant Bug (including Lygus and Stink Bug) Rindworm Squash Vine Borer	Dust lightly. Apply up to 8 times per season. May be applied until harvest.
*Cucurbit Vegetables are: Chayote (fruit), Chinese Waxgourd, citron melon, cucumber (gherkin), gourds (edible includes hyotan, cucuzza, hechima), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelons (includes true cantaloupe, cantaloupe, casaba, crenshaw, golden pershaw, honeydew melons, honey balls,		

mange melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, pumpkin, squash (summer such as crookneck, scallop, straightneck, vegetable marrow, zucchini), squash (winter such as butternut, calabaza, hubbard, acorn, spaghetti), watermelon (includes hybrids).		
Fruiting Vegetables		
Eggplant	Colorado Potato Beetle Vegetable Leafminer Cabbage Looper Flea Beetle	Dust lightly. Apply up to 10 times per season. May be applied until 3 days before harvest.
Peppers (Bell)	Pepper Weevil Vegetable Leafminer Cabbage Looper Flea Beetle European Corn Borer	Dust lightly. Apply up to 8 times per season. May be applied until 3 days before harvest.
Tomatoes	Tomato Fruitworm Cabbage Looper Tomato Pinworm Vegetable Leafminer Beet Armyworm Southern Armyworm Colorado Potato Beetle Granulate Cutworm Hornworm	Dust lightly. Apply up to 6 times per season. May be applied until day of harvest.
Leafy Vegetables		
Leafy Vegetables* (Except brassica)	Vegetable Leafminer Cabbage Looper Cutworm Complex Fall Armyworm Beet Armyworm Corn Earworm Leafhopper European Corn Borer	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.
*Leafy vegetables are: Amaranth, arugula, cardoon, celery, Chinese celery, celtuce, chervil, corn salad, chrysanthemum, edible-leaved chrysanthemum, garland, cress (garden, yellow rocket and winter), dandelion, dock (sorrel), endive, fennel, lettuce (head and leaf), orach, parsley, purslane (winter and garden), radicchio, rhubarb, spinach, spinach vine (Malabar spinach and Indian spinach), spinach (New Zealand), Swiss Chard.		
Root, Tuber, and Bulb Vegetables		
Onions (Dry bulb and Garlic)	Onions Thrips Armyworm Cutworm Leafminer Onion Maggot (Adult) Stink Bug	Dust Lightly. Apply up to 6 times per season. May be applied until 1 day before harvest.

Potatoes	Colorado Potato Beetle Potato Leafhopper Aster Leafhopper Potato Flea Beetle Potato Aphid Potato Tuberworm Potato Psyllid Cabbage Looper	Dust lightly. Apply up to 8 times per season. May be applied until 14 days before harvest.
Turnips (FL, GA, IL, IN, OK, SC, TX, WA only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid	Dust lightly. May be applied until 1 day before harvest.
(TX, OK, IL, IN only)		Apply up to 8 times per season
(SC, GA, FL, WA only)		Apply up to 4 times per season
Cereal Grains		
Sweet Corn	Corn Earworm European Corn Borer Fall Armyworm Southern Armyworm Corn Rootworm (Adult)	Dust lightly. Apply up to 6 times per season. May be applied until 1 day before harvest.
(FL only)	Cutworm Aster Leafhopper Flea Beetle	Apply up to 8 times per season
Other Crops		
Artichokes	Artichoke Plume Moth	Dust lightly. Apply up to 6 times per season. May be applied until 1 day before harvest.
Asparagus	Cutworm Complex Asparagus Beetle	Dust lightly. Apply up to 4 times per season. May be applied until 1 day before harvest.
Ornamentals		
Asparagus Fern	Japanese Beetle Asparagus Beetle Lygus Bug	Dust lightly. Apply to the fern stage of asparagus plant after speer harvest when adult stage of the asparagus beetle, lygus bugs and the adult stage of the Japanese beetle are present.
Roses	Beet Armyworm Cabbage Looper	Dust lightly.

	Omnivorous Leafroller	
Tree Crops		
Dust lightly, using a dust gun or rotary duster [an appropriate duster as available].		
Almonds	Navel Orangeworm Peach Twig Borer	Dust lightly. Do not apply after petal fall.
Apples	Apple Aphid Redbanded Leafroller Obliquebanded Leafroller Plum Curculio White Apple Leafhopper Spotted Tentiform Leafminer Tarnished Plant Bug	Dust lightly. Do not apply after petal fall.
Avocados	Thrips Leafhopper Mirid Bug Lepidopterous Larvae	Dust lightly. Apply up to 6 times per season. May be applied until 7 days before harvest.
Cherries	Lesser Peachtree Borer Rose Chafer Green Fruitworm Redbanded Leafroller Plum Curculio Tarnished Plant Bug	Dust lightly. Apply when insects appear. In the East: do not exceed 6 applications per season, with no more than 4 applications after petal fall. In the West: do not apply more than 4 applications per season, with no more than 3 applications after petal fall.
Filberts	Filbertworm Obliquebanded Leafroller	Dust lightly. Apply up to 4 times per season. May be applied until 14 days before harvest.
Papaya (For use in Florida only)	Fruit Fly	Dust lightly. Apply up to 6 times per season. May be applied until 7 days before harvest.
Peaches	Peach Twig Borer Oriental Fruit Moth Green Fruitworm Lesser Peachtree Borer Plum Curculio Tarnished Plant Bug Rose Chafer	Dust lightly. Apply up to 5 times per season. May be applied until 14 days before harvest.
Pears (Dormant)	Pear Psylla	Dust lightly. Apply up to 2 times per season. May be applied during the dormant through the delayed dormant growth periods only.
Pears (Pre-Bloom)	Pear Psylla Codling Moth Green Fruitworm	Dust lightly. Apply up to 2 times per season. May be applied from dormant through bud burst.

Pistachios	Peach Twig Borer Navel Orangeworm	Dust lightly. Apply up to 2 items per season. May be applied until 10% hull split.
Walnuts	Navel Orangeworm Codling Moth Walnut Hush Fly	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.

Alternate or optional wording below in brackets “[]” to be used before Directions for Use or on front panel.

[Product Usage: Home Gardens – Kills 60+ Insects including Apple Aphids, Armyworms, Cabbage Loopers, Colorado Potato Beetles, Cutworms, Leafminers, Japanese Beetles and other insects as listed.

Use On – Vegetables, Fruit and Ornamental Plants

Usage – Dust Lightly

Pet Use – Kills - Fleas, Ticks, Lice

Use On – Dogs [Companion Animals and Non-commercial livestock]

Usage – Dust Animals Thoroughly]

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry area away from heat or open flame.

PESTICIDE DISPOSAL: If partly filled – Do not pour or dispose down-the-drain or sewer. Call your local solid waste Agency for local disposal options.

CONTAINER DISPOSAL: If empty – Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Phaeton Corporation
1501 E. Woodfield Rd, Ste 200 West
Schaumburg, IL 60173

[Lot number added at production]