

# UNICORN 45% PERMETHRIN FLY & TICK INSECTICIDE

PROTECTS HORSES FROM FLIES, GNATS, MOSQUITOES AND TICKS

- KILLS AND REPELS TICKS FOR UP TO [4 WEEKS] [3 TO 4 WEEKS] INCLUDING DEER TICKS (vector of LYME DISEASE), BROWN DOG TICKS AND AMERICAN DOG TICKS
- PROTECTS HORSES FROM HOUSE FLIES, STABLE FLIES, FACE FLIES, HORN FLIES, EYE GNATS, AND TICKS
- KILLS AND REPELS HOUSE FLIES, STABLE FLIES, FACE FLIES, HORN FLIES, EYE GNATS, AND TICKS ON HORSES
- AIDS IN THE CONTROL OF HORSE FLIES, DEER FLIES, MOSQUITOES AND BLACK FLIES ON HORSES

**ACTIVE INGREDIENT:**

Permethrin\* ..... 45.0%

**OTHER INGREDIENTS:** ..... 55.0%

**Total:** ..... 100.0%


\*(CAS #52645-53-1)

**KEEP OUT OF REACH OF CHILDREN**


## CAUTION

**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

NET CONTENTS: 3 [ 0.34 fl. oz. (10.0 mL) ] Applicators or 6 [ 0.34 fl. oz. (10.0 mL) ] Applicators or 1.01 fl.oz. (30 mL) Dropper Bottle or 2.03 fl.oz. (60 mL) Dropper Bottle, or 4.06 fl.oz (120 mL) Wick Applicator Bottle with measured dose chamber



**Manufactured By:**  
Unicorn Laboratories  
12385 Automobile Blvd.  
Clearwater, FL 33762  
(727) 578-4545



**READ ENTIRE LABEL BEFORE EACH USE  
USE ONLY ON HORSES OVER 12 WEEKS OF AGE**

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

**Hazards to Humans:** Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling.

**Hazards to Domestic Animals: FOR EXTERNAL USE ON HORSES ONLY.** Do not use on foals under 12 weeks of age. Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing horses. Consult a veterinarian before using on horses with known organ dysfunction. **DO NOT USE ON ANIMALS OTHER THAN HORSES.** Certain medications can interact with pesticides. It is advisable to consult a veterinarian before using this product with any other pesticide or drug.

| <b>FIRST AID</b>   |  |
|--|--|
| <b>If swallowed</b>  | <ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> |
| <b>If in eyes</b>  | <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>If on skin or Clothing</b>  | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 – 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>  |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. |  |

**Adverse Reactions:** Some animals may be sensitive to ingredients in this product. Reactions in horses may include skin sensitivity. Horses may show lethargy, increased pruritis (itchiness), erythema (redness), rash and hair discoloration or hair loss at the application site. Observe the horse following treatment. Sensitivity may occur after using ANY pesticide product on horses. If signs of sensitivity occur bathe your horse with a mild, non-insecticidal shampoo and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

### **ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish. Do not add directly to water. Do not contaminate water when disposing of product or packaging.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **HORSES**

**DO NOT USE ON HORSES INTENDED FOR FOOD USE**  
**DO NOT USE ON FOALS UNDER 12 WEEKS (3 MONTHS) OF AGE**

Remove a product tube from package and hold in an upright position. Twist or cut off top of applicator tube. Remove cap carefully. Apply one tube (10.0 mL) of product solution as follows:

- 5 mL along the back line starting at the poll (top of head behind the ears), along the base of the mane, down the dorsal midline to the dock (base of the tail).
- 1 mL to the top of the forehead under the forelock.
- 1 mL as a stripe to the front of each foreleg, from the ankle up to the knee (2 mL total).
- 1 mL as a stripe to the front of each hindleg, from the ankle up to the hock (2 mL total).
- Do not allow the product to contact horses eyes or mucous membranes.
- Wrap tube and put in trash.
- Do not reapply for 14 days (2 weeks).

**\*\*DROPPER BOTTLE MULTI-DOSE PACKAGING:**

Apply 10.0 mL of product solution as follows:

- 5 mL along the back line starting at the poll (top of head behind the ears), along the base of the mane, down the dorsal midline to the dock (base of the tail).
- 1 mL to the top of the forehead under the forelock.
- 1 mL as a stripe to the front of each foreleg, from the ankle up to the knee (2 mL total).
- 1 mL as a stripe to the front of each hindleg, from the ankle up to the hock (2 mL total).
- Do not allow the product to contact horses eyes or mucous membranes.
- Do not reapply for 14 days (2 weeks).

Insert dropper tube into insecticide solution, squeeze bulb and then release to fill tube barrel to 1 mL or 5 mL line for application to locations on horse as described. Position dropper tube end in horse's hair next to skin and squeeze bulb to apply all of the solution to the horse's skin.

**\*\*Note: For Sale to Veterinarians Only.**

**WICK APPLICATOR BOTTLE MULTI-DOSE PACKAGING:**

Apply 10.0 mL of product solution as follows:

Remove protective cap, squeeze container to fill measuring chamber to 10 mL line. Invert bottle and then rub wick across horse's hair next to skin applying to locations on horse as described.

- 5 mL along the back line starting at the poll (top of head behind the ears), along the base of the mane, down the dorsal midline to the dock (base of the tail).
- 1 mL to the top of the forehead under the forelock.
- 1 mL as a stripe to the front of each foreleg, from the ankle up to the knee (2 mL total).
- 1 mL as a stripe to the front of each hindleg, from the ankle up to the hock (2 mL total).
- Do not allow the product to contact horses eyes or mucous membranes.
- Do not reapply for 14 days (2 weeks).

**SUGGESTED MINIMAL LABELING  
FOR EACH INDIVIDUAL TUBE**

**ACTIVE INGREDIENT: Permethrin ..... 45%**

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

**DO NOT USE ON ANIMALS OTHER THAN HORSES**

**Read label before use.  
10 cc [0.34 fl oz] [10 mL]**

**EPA Reg. No. 28293-XXX**

**EPA Est. No. 62478-FL-1**

414

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in cool, dry place. Protect from freezing.

**Disposal:** If empty – Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

Never place unused product down any indoor or outdoor drain.

**Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.**