

GENERAL FUMIGATION DIRECTIONS

For specific information on dosage, uses, insect problems, procedures, or safety precautions, consult "Hy-Yield 98-2".

This fumigant may be used in box cars, warehouses, fumigation vaults, food and cereal mills, ships and bulk grain storages. Do not fumigate if grain moisture is high or if grain temperature is low (below 60°F) or if there is excessive dockage.

INSECTS CONTROLLED: Granary weevil, grain beetle, rice weevil, cadellid, mites, bran bugs, grain borers, mealworms, Indian meal moth. Kills insects in all life stages. Kills rats and mice in fumigated area.

SPACE FUMIGATION: The usual dosage rate for each 1000 cu. ft. will range from 1 to 3 lbs. 12 to 24 hours exposure depending on tightness of structure and kind and amount of commodity to be stored.

BULK GRAIN DOSAGE RATES

Generally above 65°F


Shelled corn 2 lbs. each 1000 cu. ft.
Wheat similar 3 lbs. each 1000 cu. ft.
Small grains 3 lbs. each 1000 cu. ft.
Millet grain 4 lbs. each 1000 cu. ft.
Sorghum 4 lbs. each 1000 cu. ft.

CAUTION: Extinguish all fires, such as pilot lights and glowing heating units, before fumigation.

Do not use in spaces higher than recommended. Do not use in spaces where there is a possibility of fire. Do not use in spaces where there is a possibility of explosion.

Read directions carefully. Do not use in spaces where there is a possibility of fire. Do not use in spaces where there is a possibility of explosion. Do not use in spaces where there is a possibility of fire. Do not use in spaces where there is a possibility of explosion.

CAUTION: The pack is for use in spaces where there is a possibility of fire. Do not use in spaces where there is a possibility of explosion. Do not use in spaces where there is a possibility of fire. Do not use in spaces where there is a possibility of explosion.



ANTIDOTE

- Remove victim from fumigation immediately. • Keep victim warm and dry. • Give artificial respiration if breathing has stopped. • Call a physician immediately.

WARNING: Poisonous to humans and animals

- Avoid contact with skin and eyes. • Contact with mouth of fumigant should be avoided. • Do not eat, drink, or smoke while using. • If contact with skin occurs, wash with soap and water for at least 15 minutes. Remove contaminated clothing and shoes. • Do not use in areas where food is stored. • Do not use in areas where children or animals are present.

For more information on the use of this fumigant, consult the "Hy-Yield 98-2" manual.

98-2

CONTAINS 2% CHLOROPICRIN

ACTIVE INGREDIENTS:

	By Wt.
Methyl bromide	98%
Chloropicrin	2%
Total	100

KEEP OUT OF REACH OF CHILDREN
POISON

NET WEIGHT 25 LBS.

MADE IN U.S.A.

HY-YIELD INCORPORATED

HY-YIELD INCORPORATED

SOIL FUMIGATION

The following are general directions for the use of this fumigant in soil fumigation.

SOIL FUMIGATION USES: This soil fumigant kills weed seeds and certain soil fungi, nematodes and other soil pests. It is also used to control tobacco, vegetable and tree seedlings where soil is infested with insects and nematodes. It is also used to control weeds and other pests in vegetable and tree seedlings.

SOIL PREPARATION: Soil should be prepared before fumigation. Do not fumigate soil that is too wet.

EXPOSURE PERIODS: Exposure periods should be between 6 and 24 hours depending on temperature.

EXPOSURE PERIODS: Exposure periods should be between 6 and 24 hours depending on temperature.

DOSAGE: Dosage should be between 1 and 3 lbs. per 1000 cu. ft.

WHEN TO PLANT: Plant should be planted 3 to 5 days after seed setting out vegetable grow.

APPLICATION OF 98-2: Method of application should be as follows: 1. Apply 1 lb. per 1000 cu. ft. of soil. 2. Apply 1 lb. per 1000 cu. ft. of soil. 3. Apply 1 lb. per 1000 cu. ft. of soil.

APPLICATION OF 98-2: Method of application should be as follows: 1. Apply 1 lb. per 1000 cu. ft. of soil. 2. Apply 1 lb. per 1000 cu. ft. of soil. 3. Apply 1 lb. per 1000 cu. ft. of soil.


APPLICATION OF 98-2: Method of application should be as follows: 1. Apply 1 lb. per 1000 cu. ft. of soil. 2. Apply 1 lb. per 1000 cu. ft. of soil. 3. Apply 1 lb. per 1000 cu. ft. of soil.

APPLICATION OF 98-2: Method of application should be as follows: 1. Apply 1 lb. per 1000 cu. ft. of soil. 2. Apply 1 lb. per 1000 cu. ft. of soil. 3. Apply 1 lb. per 1000 cu. ft. of soil.

APPLICATION OF 98-2: Method of application should be as follows: 1. Apply 1 lb. per 1000 cu. ft. of soil. 2. Apply 1 lb. per 1000 cu. ft. of soil. 3. Apply 1 lb. per 1000 cu. ft. of soil.

APPLICATION OF 98-2: Method of application should be as follows: 1. Apply 1 lb. per 1000 cu. ft. of soil. 2. Apply 1 lb. per 1000 cu. ft. of soil. 3. Apply 1 lb. per 1000 cu. ft. of soil.

APPLICATION OF 98-2: Method of application should be as follows: 1. Apply 1 lb. per 1000 cu. ft. of soil. 2. Apply 1 lb. per 1000 cu. ft. of soil. 3. Apply 1 lb. per 1000 cu. ft. of soil.



CONTAINS 2% CHLOROPICRIN

Methyl bromide
Chloropicrin

98%
2%

100%

KEEP OUT OF REACH OF CHILDREN
POISON

1994-1995 2011-12

HY-VECT, INCORPORATED

The following information is being furnished to you for your information only. It is not intended to be used for any other purpose. It is not to be used for any other purpose. It is not to be used for any other purpose.

SOME PLANT GRAFT USES — Rhizopus is an effective fungicidal fungicide for the control of nematodes, weed seeds and other soil borne diseases causing fungi in seed beds and potting soils. It contributes to a maximum production of healthy vigorous tobacco, vegetable transplants, nursery and forest tree seedlings. Where slips, cuttings, sprouts, c. seeds are used as in greenhouses, nurseries, pot greens, lawns and ornamental gardens, healthy and vigorous rooting is established quickly.

SOIL PREPARATION: Plow or otherwise work soil to a fine tilth. Add fertilizer manure, or compost work in well. Soil should be moist and friable for best results. Do not fumigate when soil is dry or excessively wet.

4.401 v. Do not fumigate when soil temperature is below 50 F. 982 Use of vegetable products is limited to the production of transplants only.

EXPOSURE PERIODS: Expose for 48 hours when soil temperature is between 50° F and 60° F and for 24 hours when above 60° F.

DOSAGE - From 10 mg/kg up to 60 mg/kg
for each 100 lb of body weight

WHEN TO PLANT After the soil has warmed sufficiently for 3 days before seeding or 5 to 7 days before setting out vegetative growth.

method consists of surveying the population by making the randomised telephone calls to the households and by interviewing a randomly selected individual under the telephone. The sample size was at least 200-400 persons for each age group. The mean of the telephone number was randomly selected by computer. The telephone number was called by computer. The telephone number was called by computer. The telephone number was called by computer.

There is a growing concern that the current system of public provision of health care in the United Kingdom is not sustainable. The current system is based on a combination of general taxation and contributions from employers and employees. This system has led to a steady increase in the size of the public sector, which is now estimated to be around 25% of GDP. This is a significant increase from the 1970s, when the public sector was around 15% of GDP. The current system is also facing a number of challenges, including a rising cost of health care, a shortage of health care workers, and a growing demand for health care services. These challenges are likely to lead to a further increase in the size of the public sector, which could have significant implications for the economy and the welfare of the population. This paper examines the current system of public provision of health care in the United Kingdom, and discusses the challenges it faces. It also considers some of the options for reforming the system, and the implications of these options for the economy and the welfare of the population.

Trademark and Copyright © 1995 by
 International Technology Corporation
 All rights reserved. This document is the property of International Technology Corporation and is loaned to your organization. It and its contents are not to be distributed outside your organization. For more information on this issue, contact us at:
 International Technology Corporation
 10000 Highway 100, Suite 100
 New York, NY 10020-1000

[illegible]

CONTROL OF ARMILLARIA MELLEA OAK ROOT FUNGUS IN DECIDUOUS FRUITS AND NUTS CITRUS AND VINEYARDS

For a full description of the treatment Cross which are being tested, see the following table. We will have harvest data for the first harvest of at least 24 points. Main effect of the treatment data will be discussed in the following.

2: 100% H_2SO_4 (100 ml) and 100 ml 50% H_2SO_4 by choice

wood at an average 18% increase up to a density of 0.1 trees m⁻². Covered a 10% side with an empty space, used a 10% side with Armadillo (Mellie) and a 10% side with grass, and a 10% side with groves and a 10% side with the orchards and a preparation of the soil (0.1 m² m⁻²) to use where trees are used.

On 10/10/2010, 24 months.

[illegible]

1. *Chlorophyll *a** was determined by the method of Arar and Collins (1971). The concentration of chlorophyll *a* was determined by measuring the optical density of the extract at 663 nm. The concentration of chlorophyll *a* was determined by measuring the optical density of the extract at 663 nm.

[illegible]

the 1990s, the number of people in the world who are under 15 years of age is expected to increase from 1.1 billion to 1.5 billion. The number of people aged 65 and over is expected to increase from 200 million to 400 million. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion.