



67-33

PREPLANT SOIL FUMIGANT

ACTIVE INGREDIENTS:	By Wt.
Methyl bromide	67%
Chloropicrin	33
	Total 100

For control of nematodes, wireworms, weed seed, fungus diseases, grubs, ants and soil-borne insects. (Example: white grubs)

67-33 FUMIGATION DIRECTIONS RECOMMENDED APPLICATION RATES

For control of nematodes, wireworms, weed seed, fungus diseases, grubs, ants and soil-borne insects. (Example: white grubs)

For control of nematodes, wireworms, weed seed, fungus diseases, grubs, ants and soil-borne insects. (Example: white grubs)

For control of nematodes, wireworms, weed seed, fungus diseases, grubs, ants and soil-borne insects. (Example: white grubs)

For control of nematodes, wireworms, weed seed, fungus diseases, grubs, ants and soil-borne insects. (Example: white grubs)

NOTICE

For control of nematodes, wireworms, weed seed, fungus diseases, grubs, ants and soil-borne insects. (Example: white grubs)

NOTICE

For control of nematodes, wireworms, weed seed, fungus diseases, grubs, ants and soil-borne insects. (Example: white grubs)

AGG...
...
...

67-33

PREPLANT SOIL FUMIGANT

ACTIVE INGREDIENTS:

	By Wt.
Methoxy bromide	67%
Chloropicrin	33%
Total	100%

For control of nematodes, wireworms, weed seed, fungus diseases, grubs, ants and soil-borne insects. (Example: white grubs)

DANGER

KEEP OUT OF REACH OF CHILDREN

POISON

NET WEIGHT _____

EPA REG. NO. 20870

GROSS WEIGHT _____

BY YIELD, INCORPORATED

100 B

100 B



67-33 FUMIGATION DIRECTIONS RECOMMENDED APPLICATION RATES

Soil treatment for
Strawberries 225 to 360 lb Acre

Exposure to fumigant for 48 hours. Aerate 2
weeks after setting transplants in treated
area.

Soil treatment for
Tomatoes 225 to 325 lb Acre

Exposure to fumigant for 48 hours. Aerate 2
weeks after setting transplants in treated
area.

CITRUS SOIL FOR CONTROL OF *PHY-* *TOPHTHORA* IN FLORIDA SANDY SOILS

This is a preplant or replant treatment. Trees
which are planted in this treated soil will not
have harvestable fruit for a period of at least
24 months. Use a minimum of 1.1 to 1.2 to 100
sq. ft. Exposure to fumigation for 96 hours.
Treated area with a 4 m tarpaulin.
Do not re-plant to a depth of 4 feet.
Aerate 2 weeks after setting transplants in
treated area.

Apply at 100 lb per acre in lines apart to
a depth of 4 inches. Seal fumigant with a
tarpaulin. If a leak is immediately behind
the tarpaulin, the tarpaulin must be removed
and replaced with the recommended fumigant.
Seal the tarpaulin for 2 weeks.
Do not re-plant to a depth of 4 feet.

Apply at 100 lb per acre in lines apart to
a depth of 4 inches. Seal fumigant with a
tarpaulin. If a leak is immediately behind
the tarpaulin, the tarpaulin must be removed
and replaced with the recommended fumigant.
Seal the tarpaulin for 2 weeks.
Do not re-plant to a depth of 4 feet.

Apply at 100 lb per acre in lines apart to
a depth of 4 inches. Seal fumigant with a
tarpaulin. If a leak is immediately behind
the tarpaulin, the tarpaulin must be removed
and replaced with the recommended fumigant.
Seal the tarpaulin for 2 weeks.
Do not re-plant to a depth of 4 feet.

Apply at 100 lb per acre in lines apart to
a depth of 4 inches. Seal fumigant with a
tarpaulin. If a leak is immediately behind
the tarpaulin, the tarpaulin must be removed
and replaced with the recommended fumigant.
Seal the tarpaulin for 2 weeks.
Do not re-plant to a depth of 4 feet.

POISON
DANGER

MAY BURN SKIN AND EYES
POISONOUS IF SWALLOWED

DO NOT BREATHE VAPORS
DO NOT GET IN EYES OR ON CLOTHING

Wash thoroughly with soap and water
if you get this material on your skin or
clothing. If you get this material in your
eyes, wash them with clean water for
15 minutes. If you get this material
in your mouth, do not swallow. Rinse
your mouth with water.

POISON


SEND FOR DATA SHEET AND FUMIGATION
KIT

For more information on the use of
67-33, contact your local Extension
Agent or the Florida Department of
Agriculture, Bureau of Plant Industry,
Tallahassee, Florida.

NOTICE

Following soil fumigation, undesirable
concentrations of chloropicrin may drift to
nearby areas. If this occurs, immediately
cover treated area with a plastic tarpaulin.
Leave tarpaulin on overnight and remove
during the daytime. Recover the area if
irritation indicates that the undesirable
concentration is continuing. These concentra-
tions are most likely to occur in the evening
at night when the air is most likely to be
still and overcast. Do not make application
of 67-33 when there is a strong wind or
when there is a heavy overcast.

NOTE CAREFULLY: Fertilizer containing
the heavy metal cadmium should not be
applied to soil treated with 67-33. If
fertilizer is applied to soil treated with
67-33, it should be applied after the
Fertilizer containing the heavy metal
cadmium should not be used. Apply only
fertilizer after the chloropicrin has
dissipated and the soil temperature is at
least 50°F. Do not apply fertilizer to
soil treated with 67-33 until the soil
temperature is at least 50°F. Do not
apply fertilizer to soil treated with 67-33
until the soil temperature is at least 50°F.
Do not apply fertilizer to soil treated
with 67-33 until the soil temperature is
at least 50°F.



side of the plants, which
should be examined and
destroyed immediately
at harvest. Some of these
labels are not available,
however, if you are in
Mexico or Canada.
Do not use this product
if you are in Mexico or
Canada. Do not use this
product if you are in
Mexico or Canada.