

USE DIRECTIONS:

Disinfection

Use at 2 ounces per gallon. This product will be effective against *Pseudomonas aeruginosa* PRD-10 at this level. This has been substantiated by the AOAC Use Dilution Confirmation Test. At this level, the product is fungicidal against pathogenic fungi.

For best results, use RC-2 with a cloth or mop and apply to walls, floors and other hard surfaces.

For heavy soil or organic matter, a pre-cleaning step is recommended.

Sanitization

If used for sanitization of previously cleaned food equipment or food contact items, limit the level of active quaternaries to 200 ppm (2.8 ounces per 5 gallons of water). **NO RINSE REQUIRED.**

This product when used on environmental, inanimate, hard surfaces at 2 ounces per gallon of water, is effective against Influenza A₂, Herpes Simplex, Adenovirus type 2, and Vaccinia viruses.

PHENOL COEFFICIENTS:

Guaranteed	Minimum
S. aureus	44
S. typhosa	28

Rinse empty container thoroughly with water before discarding or returning drum to drum reconditioner.

(RC - 2)

**CLEANER
DEODORIZER**

**DISINFECTANT
FUNGICIDE**

Designed for Hospital and Institutional use.

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
dimethyl benzyl ammonium chlorides 2.25%
n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl
ethylbenzyl ammonium chlorides 2.25%
Sodium Carbonate 3.00%
Tetrasodium ethylenediamine tetraacetate 1.00%

INERT INGREDIENTS:..... 91.50%

100.0%

WARNING

KEEP OUT OF REACH OF CHILDREN

See right side panel for additional precautionary and first aid statements.

EPA Reg. No. 29780-5

EPA EST. 29780-GA-1

Net contents gal.

CENTENNIAL CHEMICALS, INC.
1094 Huff Road, N. W.
Atlanta, Georgia 30318

RC-2 is a cleaner, disinfectant, and sanitizer, designed for institutional use such as Hospitals, Nursing Homes and Institutions where housekeeping will be of prime importance in controlling the hazard of cross-infection from environmental surfaces. RC-2 can provide the assurance you need in this area, when used as directed. For use on floors, walls, sink tops, garbage pails, telephones, rest rooms, bed pans and operating rooms. Effective against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Pseudomonas aeruginosa*. Also effective against pathogenic fungi, *Trichophyton interdigitale*.

When used for sanitization of previously cleaned food processing equipment and food contact items, limit active quaternaries to 200 ppm. At this level, **NO POTABLE WATER RINSE** is required.

WARNING

Keep Out of Reach of Children. Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

