CAPTAN 5001 SEED PROTECTANT

SLURRY APPLICATION FOR SEED DISEASE CONTROL

DYE ADDED



7.5 GAL. NET 70 LBS. NET

ACTIVE INGREDIENT:

CAPTAN *	32%
INERT INGREDIENTS68	
[*] য় (নাল এটেনতাশ্র (লিপ্র Thio) -4-CYCLOHEXENE-1, 2-DICARBOXIMIDE	100%
ACONTAINS DED CARTAN DED CALLONS	

C YION

KEEP OUT OF REACH OF CHILDREN

CAUSES SKIN AND EYE IRRITATION.

MAY PE HARMFUL IF SWALLOWED OR INHALED.

TALL TO MITACT WITH SKIN, EYES OR CLOTHING.

AVOID BREATHING DUST WEAT RESPIRATOR WASH THOROUGHLY AFTER HANDLING

DI N T STURE OR USE NEAR FOOD OR FEED

N ASE OF CONTACT IMMEDIATELY FLUSH EYES WITH PRENTY OF WATER FOR AT LEAST IS MEDUTES WASH SKIN WITH SOAP AND WATER.

CAPTAN 5001 SEED PROTECTANT IS A MICROFINE EMULSIFIABLE LIQUID ESPEC-

TALLY FORMULATED FOR THE TREATMENT OF SEED, PRIOR TO STOPAGE AND

PLANTING, TO PROTECT SEED FROM MOLDS AND OTHER FUNGICAUSING STORAGE

LOSS AND TO PROTECT SEED FROM MEET P. RNE AND SOIL BORNE FUNGI WHICH

CAUSE SEED DECAY AND SEEDLING BLIGHT &

COMPATIBILITY: CAPTAN 5001 CAN BE USED IN COMBINATION WITH MOST COM-

MONLY USED SEED PROTECTANTS, BOTH INSECTICIDES AND FUNGICIDES AS WELL

AS FERTILIZERS DO NOT USE WITH STRONGLY ALKALINE MATERIALS

PLANT SAFETY: AT THE RECOMMENDED DOSAGE CAPTAN 5001 IS NON INJUDIOUS

FROM THE STANDPOINT OF SEED INJURY OR YIELD REDUCTION. THE GERMINATION OF

TREATED SEED IS FREQUENTLY STIMULATED GIVING A SUPERIOR RESULT AS COM

FOITO MESULTS FROM OTHER SEED FUNGICIDES.

Cameraman's Note

Poor Copy

GENERAL USE PRECAUTIONS

DIRECTIONS FOR USE

SLURRY TREATMENT, MIX THE RECOMMENDED AMOUNT OF THIS MATERIAL WITH THE REQUIRED AMOUNT OF WATER FOR THE SLURRY EQUIPMENT TO BE USED.

CAUSES SKIN AND EYE IRRITATION. MAY BE HARMFUL IF SWALLOWED OR INHALED.

A FREE ONTACT WITH SKIN, EYES OR CLOTHING.

AVOID BREATHING DUST WEAT RESPIRATOR WASH THOROUGHLY AFTER HANDLING DU N T STURE OR USE NEAR FOOD OR FEED

N ANE OF CONTACT PUMERINATELY FLUSH EYES WITH PRENTY OF WATER FOR AT CHAST IS MURULES. WASH SKIN WITH SOAP AND WATER.

GENERAL USE PRECAUTIONS

READ THRECTIONS BEFORE USING AND USE ONLY AS SPECIFIED ON THIS LABEL.

TO A HOUSE OUT OF LAKES, STREAMS OR PONDS. DO NOT CONTAMINATE

TO A HOUSE AND OF EQUIPMENT OR DISPOSAL OF WASTES.

IMPUREARTY TOFATED SEED MUST NOT BE USED FOR HUMAN CONSUMPTION NOT AS AN MAN FEND. BADS JONE MINIST TREATED SEED SHOULD BE LABELED!

TREATED SEED DO NOT USE FOR FEED, FOOD OR OIL PURPOSES

THE FEDERAL GO PERMENT AND SOME STATES HAVE ISSUED REGULATIONS FOR THE ACT OF THE ACT AND THE ACT AND

PROTECT FROM EXCESSIVE HEAT . PRODUCT FREEZING TEMPERATURE 320 CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES. WEIGHT PER GALLON EQUALS 9.3 LBS.

E.P. A. REG. NO. 29964-1

MADE IN U.S.A. BY
PIONEER HI-BRED INTERNATIONAL, INC.
DES MOINES, IOWA 50308

MONLY USED SEED PROTECTANTS, BOTH INSECTICIDES AND FUTURITIES AS WELLS.
AS FERTILIZERS DO NOT USE WITH STRONGLY ALKALINE MATERIALS.

FROM THE STANDPOINT OF SEED INJURY OR YIELD REDUCTION. THE GERMINATION OF TREATED SEED IS FREQUENTLY STIMULATED GIVING A SUPERIOR RESULT AS COM

(IND) 29944-1

DIRECTIONS FOR USE

SLURRY TREATMENT: MIX THE RECOMMENDED AMOUNT OF THIS MATERIAL WITH THE REQUIRED AMOUNT OF WATER FOR THE SLURRY EQUIPMENT TO BE USED.

RECOMMENDATIONS

FOR CONTROL OF MANY SEED ROTS, SEEDLING POIS, ROOT ROTS, SOIL BORNE FUNGI AND KERNEL SMUT OF SORGHUM

AMOUNT OF CAPTAP BOOL BY WEIGHT TO USE ON REEDS

SEED	SLURRY
ALFALFA, CLOVER, LESPEDEZA, TREFOIL	12.5 07 /100 Lt.
BARLEY, OATS, WHEAT	1.67 OZ./BU.
CORN - FIELD	1.67 02./BU.
SWEET	5.83 QZ./100 E.
QRASSES	12.5 07./100 60.
SORGHUM (HULL-LESS)	3.12 02./BU.
SOYBEANS	2.6 OZ./BU.

FURTHER SPECIFIC INFORMATION ON THIS SUBJECT MAY BE OBTALLED FROM STATE AGRICULTURAL EXPERIMENT STATIONS OR THE STATE AGRICULTURAL EXPENSION SERVICE.

NOTICE: PIONEER HI-BRED INTERNATIONAL, INC. MAKES NO WAR-RANTY, EXPRESS OR IMPLIED, INCLUDING THE WARRANTIES OF MERCHAN-TABILITY AND/OR FITNESS FOR ANY PARTICULAR PURPOSE, CONCERNING THIS MATERIAL, EXCEPT THOSE WHICH ARE CONTAINED ON PIONEER'S LABEL.

Cameraman's Not

Poor

\$11072