


PYRELLIN E.C.

Pyrellin E.C. is a broad spectrum insecticide utilizing natural pyrethrins for quick knockdown, rotenone for longer kill with aphid control, and essential oils for insect activation, gustatory inhibition, and insect repellence.

- SAFETY with proven EFFECTIVENESS
- Over 20 years use in rigorous and demanding South Florida agriculture
- Cost effective
- Approved for preharvest use

WARRANTY

WEBB WRIGHT CORPORATION warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of WEBB WRIGHT CORP. In no case shall WEBB WRIGHT CORP. be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. WEBB WRIGHT CORP. MAKES NO WARRANTY, EXPRESS OR IMPLIED, INCLUDING THE MERCHANTABILITY AND/OR FITNESS FOR ANY PARTICULAR PURPOSE, CONCERNING THIS MATERIAL, EXCEPT THOSE WHICH ARE CONTAINED ON THE LABEL.



Pyrellin E.C.

EMULSIFIABLE CONCENTRATE INSECTICIDE

ACTIVE INGREDIENTS:

Pyrethrin	0.60%
Permethrin	0.50%
Other Carb Resins	0.50%
INERT INGREDIENTS	98.40%

Contents Per Gallon:
0.042 lbs. of Pyrethrin
0.035 lbs. of Permethrin
0.035 lbs. of Other Carb Resins

KEEP OUT OF REACH OF CHILDREN
CAUTION
 SEE REVERSE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS 2.5 GALLONS

Manufactured by:
WEBB WRIGHT CORPORATION



P.O. Box 1572
Ft. Myers, FL 33908

EPA Establishment Number 30573FL01

EPA Registration Number 30573-2

**PRECAUTIONARY STATEMENT
STATEMENT OF PRACTICAL TREATMENT**

IF SWALLOWED: Contact a physician at once.
IF INHALED: Remove from contact & seek medical attention.
IF ON SKIN: Flush immediately with copious amounts of water.
IF IN EYES: Flush immediately with copious amounts of water. If irritation develops seek medical attention.

HAZARDOUS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin or eyes. In case of contact, immediately flush eyes or skin with plenty of water. For eyes get medical attention if irritation develops or persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, ponds & streams. Do not contaminate water by cleaning equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Keep away from heat and open flame.

DIRECTIONS FOR USE

General Classification


It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. SEE DIRECTIONS IN THE ENCLOSED CIRCULAR.

STORAGE AND DISPOSAL

STORAGE: Keep out of heat and cold. Store only in the original container. Keep away from fertilizer, food, other pesticides & feed. Keep out reach of children & animals.


DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



Pyrellin E.C.





Webb E.C.

DISINFESTANT CONCENTRATE INSECTICIDE

INGREDIENTS:

one ounce	0.60%	Contents Per Gallon
Cube Resins	0.50%	0.042 lbs. of Pyrethrins
Other Cube Resins	0.50%	0.035 lbs. of Rotenone
INGREDIENTS	98.40%	0.035 lbs. of Other Cube Resins

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

WARNING: Contact a physician at once.
SYMPTOMS: Remove from contact & seek medical attention.
TREATMENT: Flush immediately with copious amounts of water.
ADDITIONAL: Flush immediately with copious amounts of water. If symptoms develop, seek medical attention.

Additional Precautionary Statements

WEBB WRIGHT CORPORATION

P. O. BOX 1572
 FT. MYERS, FL 33902
 EPA REG. NO. 30573-2

GROWING CROPS (Outdoors and in Greenhouses)

VEGETABLE CROPS:

Crop	Insect	Rate Pints/Acre	Additional Instructions
Asparagus	Aphids	1-2	Apply in 25 to 100 gallons of water by ground sprayer or in 1 to 2 gallons of water by Low Volume apparatus, or in 1 to 2 pints of vegetable oil if applied with ULV spray equipment
Beans (Green, Lima, Manj. Snap)	Asparagus Beetles	-	
Beets (red)	Alfalfa Web worms	-	Apply in intervals of 7 days or less
Cabbage (Chinese, sul. NAPA red)	Bean Beetles	-	
Carrots	Beet Webworms	-	Repeat as necessary to maintain control
Celery	Cabbage Worms	-	
Corn	Celery Leaf miners	-	Apply as necessary to maintain control
Cucumber	Garden Flea hoppers	-	
Eggplant	Harlequin Bugs	-	Apply as necessary to maintain control
Lettuce (Head)	Japanese Beetles	-	
Lettuce	Leafhoppers	-	Apply as necessary to maintain control
Escarole	Leaf miners	-	
Leaf But	Loopers	-	Apply as necessary to maintain control
tercrunch	Lygus Bugs	-	
Romaine)	Mites	-	Apply as necessary to maintain control
Onions (Green Dry Bulb)	Oblique banded Leafrollers	-	
Pepper	Plant Bugs	-	Apply as necessary to maintain control
Potato	Stink Bugs	-	
Squash (Summer, Winter)	Thrips	-	Apply as necessary to maintain control
Acorn	White Flies	-	
Butternut	Cucumber Beetles	1-2	3 to 4 applications or more as eggs start to hatch and repeat at 5 to 7 day intervals
Crookneck, Hubbard)	European Corn Borers	-	
Spaghetti Squash)	Flea Beetles	-	Apply as necessary to maintain control
Zucchini)	Squash Bugs	-	
Spinach	Squash Vine Borers	-	Apply as necessary to maintain control
Tomato	Vegetable Weevils	-	
Artichokes (Globe Jerusalem)	Aphids	1-2	Apply as necessary to maintain control
Broccoli (Chinese, rab)	Leafhoppers	-	
Brussel Sprouts	Leaf miners	-	Apply as necessary to maintain control
Cauliflower	Loopers	-	
Bok Choy (Kai)	Lygus Bugs	-	Apply as necessary to maintain control
	Mites	-	
	Plant Bugs	-	Apply as necessary to maintain control
	Thrips	-	

Dates (also Chinese)
 Dewberries
 Elderberries
 Ginkgo
 Guava
 Huckleberries
 Kiwi
 Loganberries
 Longan Fruit
 Loquat
 Mango
 Mulberries
 Papayas
 Passion Fruit
 Persimmons
 Pineapple
 Pitanga Cherry
 Prickly Pear
 Quince
 Sapodilla
 Sapote (green, white, black)

Crop	Insect	Rate Pints/Acre	Additional Instructions
Almonds	Aphids	1-2	Apply in 100 gallons of water as needed for control
Beechnuts	Mites	-	
Butternuts	Thrips	-	Apply as necessary to maintain control
Cashews	White Flies	-	
Chestnuts			Apply as necessary to maintain control
Filberts			
Hazelnuts			Apply as necessary to maintain control
Hickory Nuts			
Lychee			Apply as necessary to maintain control
Macadamia Nuts			
Pecans			Apply as necessary to maintain control
Pistachio			
Pomegranate			Apply as necessary to maintain control
Tamarnad			
Walnuts			

HERBS AND SPICES:

Crop	Insect	Rate Pints/Acre	Additional Instructions
Anise	Aphids	1-2	Apply in 25 to 100 gallons water by ground sprayer or in 1 to 2 gallons of water by Low Volume apparatus, or in 1 to 2 pints of vegetable oil if applied with ULV spray equipment
Basil	Loopers	-	
Caraway	Mites	-	Apply in intervals of 7 days or less
Chicory	Plant Bugs	-	
Chives	Thrips	-	Apply as necessary to maintain control
Coriander (Cilantro)	White Flies	-	
Cumin			Apply in intervals of 7 days or less
Dill			
Fennel			Repeat as necessary to maintain control
Ginger			
Ginseng			Repeat as necessary to maintain control
Marjoram			

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin or eyes. In case of contact, immediately flush eyes or skin with plenty of water. For eyes get medical attention if irritation develops or persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, ponds & streams. Do not contaminate water by cleaning equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Keep away from heat & open flame.

GENERAL CLASSIFICATION DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in such a manner as to directly or through drift expose workers or other persons.

STORAGE AND DISPOSAL

STORAGE: Keep out of heat and cold. Store only in the original container. Keep away from fertilizer, food, other pesticides & feed. Keep out reach of children & animals.

DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS

USED ALONE

Pyrethrin E.C. can be used at rates of 1 to 2 pints per acre in 25 to 100 gallons of water by ground sprayers at any time during the growing season including use as a pre-harvest insecticide up to and including the day of harvest in those areas where aerial application is allowed. Pyrethrin E.C. should be applied in rates of 1 to 2 pints per acre with a minimum of 50 gallons of water. Repeat as needed for control. Constant spray tank agitation is required during spraying when diluting with water.

Superior control may be obtained by applying Pyrethrin E.C. in Low Volume, small droplet (70-80 micron) depending on existing weather conditions; sprays of 1 to 2 pints Pyrethrin E.C. with 1 to 2 gallons of water or in ULV (ultra low volume) small droplet sprays of 1 to 2 pints Pyrethrin E.C. with 1 to 2 pints of vegetable oil where such sprays are approved. Small droplet sprays may also minimize risk of phytotoxicity.

ULV sprays (recommended particle size 50-75 micron) applied neat (vegetable oil dilute no water) aid leaf penetration for control of mites.

COMBINED WITH OTHER INSECTICIDES

Choi, Pak Choy,
Pak Toy)
Curdoo
Cassavas (Yucca)
Cauliflower
Celery
Cherimoya
Chick Peas
Collards
Dakon
Endive
Escarole
Garlic
Greens (Dandelion, Turnip,
Mustard, Beet,
China, Poke)

Hops
Horseradish
Kale
Kohlrabi
Leeks
Lentils
Malanga
Melons (Cantaloupe, Casabas,
Crenshaws,
Honeydews,
Muskmelons,
Persian,
Watermelons)

Okra
Parsnip
Peas (Field,
Blackeye,
Green Snow,
Peppers (Bell,
Chili non bell)

Pumpkins
Radichio
Radish
Rhubarb
Rutabagas
Salsify
Shallots
Swiss Chard
Taro (Dasheen)
Turnip
Yams

Dry Beans
Peanuts
Soybeans
Subarbeans
Sunflower

Aphids
Bean Beetles
Beet Webworms
Caterpillars
Loopers
Mexican Bean Beetles
Mites
Thrips

1 - 2
Apply as necessary to maintain control

Mustard
Oregano
Paprika
Parsley
Rosemary
Sage
Savory
Tarragon
Thyme

FIELD CROPS:

Crop	Insect	Rate Pints/Acre	
Alfalfa	Aphids	1 - 2	Apply water
Barley	Armyworms	-	1 to 2
Clover	Cinch Bugs	-	Volume
Field Corn	Flies	-	pints of
Grass (Bermuda)	Mites	-	with ULV
Blue Fescue)	Thrips	-	
Joyoba			
Millet			Apply less
Rice			
Sorghum			Repeat
Wheat			tain control

MISCELLANEOUS:

Crop	Insect	Rate Pints/Acre	
Cotton	Aphids	1 - 2	Apply water
	Fleahoppers	-	1 to 2
	Loopers	-	Volume
	Mites	-	pints of
	Pink Boll Worms	-	with ULV
	Thrips	-	
			Apply less
			Repeat
			tain control

WARRANTY

WEBB WRIGHT CORPORATION warrants that this product conforms to the description on the label thereof and is reasonably fit for purposes specified when used in accordance with the directions on the label under normal use conditions. WEBB WRIGHT CORPORATION does not warrant that it can or will eliminate all risks inherently associated with use of this product. The presence of other unintended consequences may result because of conditions, presence of other materials, or the manner of use or application which are beyond the control of WEBB WRIGHT CORP. In no case shall WEBB WRIGHT CORP. be liable for consequential, special or indirect damages.

IN BARNY, MILKING PARLORS, MILK ROOMS, DRINKS AND FEEDING TROUGS

To control house flies, horn flies, stable flies, horse flies, ear flies, biting and sucking lice, poultry lice, bedbugs, mites, fleas, ticks, mosquitoes, and gnats dilute at the rate of 4 fluid ounces per gallon of water and apply as a fog or fine mist, directing the spray above livestock and poultry toward the ceiling and upper corners until the area is filled with mist using about two ounces of diluted material per 1000 cubic feet of space. For best results, close doors and windows before spraying and keep them closed for ten to fifteen minutes. Applicator should vacate treated area and ventilate before reoccupying. Repeat application as necessary.

ON HARVESTED TOMATOES AND FRUIT (Including Grapes)

To control vinegar flies and fruit flies dilute at the rate of 1 part Pyrethrin E.C. with 600 parts of water (1.5 pints per 100 gallons or 2 Tablespoons per 5 gallons of water). Keep the emulsion thoroughly mixed in the spray tank. Make treatment as follows:

1. Apply liberally to tomatoes and fruits in baskets, on trucks or in plants. Use sprayers at a high pressure for applying at the rate of five or six pints of the diluted spray to a two-ton load of tomatoes. Direct the spray for maximum coverage of the baskets or hampers. It is important to spray between and beneath the containers.
2. Spray the raw stock stacked in the yard.
3. Dip the baskets in the diluted spray, after dumping the fruit, to kill adhering larvae and pupae.

AS A GRAIN PROTECTANT*

Pyrethrin E.C. may be used to protect against grain storage insects at a dilution rate of 1 part Pyrethrin E.C. with 15 parts of water (1 pint with 1 gallon 7 pints of water). Apply directly to the grain at a rate of 4 to 5 gallons of diluted material per 1000 bushels as the grain is carried along a belt or as it enters the auger or elevator. Pyrethrin E.C. may be used in combination with a registered fumigant for use on heavily infested stored products.

Monthly inspections should be made. If the top two or three inches are found to be reinfested, re-treat, applying at the rate of 1 to 2 gallons of diluted material per 1000 bushels of stored product.

* Some yellowing or discoloration may occur due to natural plant pigments of untreated pyrethrum extract used in Pyrethrin E.C. If in doubt, test on a sample quantity before general application.

MOSQUITO CONTROL

May be used on croplands as an adulticide to protect agricultural workers during harvesting. Use at 2 to 12 ounces per acre when applied by fixed wing aircraft, helicopters or truck mounted equipment.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated

Blueberries	Maggots		1 to 2 gallons of water by
Cherries	Cherry Fruit		Low-Volume apparatus, or
Currents	Flies	-	drift to 2 pints of vegetable
Gooseberries	Gooseberry	-	oil (applied with ULV spray
Grapes	Fruit Worms	-	equipment)
Peaches	Imported Currant	-	
Pears	Worms	-	
Plums	Japanese Beetles	-	Apply in intervals of 7
Raspberries	Leafhoppers	-	days or less
	Mites	-	
	Red-necked	-	Repeat as necessary to
	Borers	-	maintain control
	Raspberry Fruit	-	
	Worms	-	
	Rose Chafers	-	
	Thrips	-	
Strawberries	Mites	1 - 2	Apply in 100 gallons of water
	Strawberry Root	-	as needed for control
	Worms	-	
	Strawberry Leaf	-	
	Rollers	-	
	Thrips	-	
Citrus	Aphids	1 - 2	Apply as necessary to main
	Caterpillars	-	tain control
	Fuller Rose	-	
	Beetle	-	
	Thrips	-	
	White Flies	-	
Acerola	Aphids	1 - 2	Apply as necessary to main
Anon	Caterpillars	-	tain control
Apricots	Fireworms	-	
Atemoya	Fruit Flies	-	
Avocadoes	Leafhoppers	-	
Boysenberries	Mites	-	
Cactus Fruit	Thrips	-	
Calamondin	Weevils	-	
Apricots	Fireworms	-	
Atemoya	Fruit Flies	-	
Avocadoes	Leafhoppers	-	
Boysenberries	Mites	-	
Cactus Fruit	Thrips	-	
Calamondin	Weevils	-	
Carambola	White Flies	-	
Crabapples			
Cranberries			

