

-Fruit, Nut & Vine Crops continued-

Crop	Insect	Rate Pints/Acre	Additional Instructions
Cranberries			
Dates (also Chinese)			
Dezberries			
Elderberries			
Guava			
Huckleberries			
Kiwi			
Loganberries			
Longan Fruit			
Loquat			
Mango			
Mulberries			
Papayas			
Passion Fruit			
Persimmons			
Pineapple			
Pitanga Cherry			
Prickly Pear			
Quince			
Sapodilla			
Sapota			
Almonds	Aphids	1-2	Apply in 25 to 100 gallons of water as needed for control
Beechnuts	Mites	.	
Butternuts	Thrips	.	
Cashews	White Flies	.	
Chestnuts			
Filberts			
Hazelnuts			
Hickory Nuts			
Lychee			
Macadamia Nuts			
Pecans			
Pistachio			
Pomegranate			
Tamarind			
Walnuts			

**HERBS & SPICES**

Crop	Insects	Rate Pints/Acre	Additional Instructions
Anise	Aphids	1-2	Apply in 25 to 100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment.
Basil	Loopers	.	
Caraway	Mites	.	
Chicory	Plant Bugs	.	
Chives	Thrips	.	
Coriander (Cilantro)	White Flies	.	Apply in intervals of 7 days or less

-Herbs & Spices continued next page-

-Herbs & Spices continued-

Crop	Insect	Rate Pints/Acre	Additional Instructions
Cumin			Repeat as necessary to maintain control
Dill			
Fennel			
Ginger			
Ginseng			
Marjoram			
Mint			
Mustard			
Oragano			
Paprika			
Parsley			
Rosemary			
Sage			
Savory			
Tarragon			
Thyme			

**FIELD CROPS**

Crop	Insect	Rate Pints/Acre	Additional Instructions
Alfalfa	Aphids	1-2	Apply in 25-100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment
Barley	Armyworms	.	
Clover	Chinch Bugs	.	
Field Corn	Flies	.	Apply in intervals of 7 days or less
Grass (Bermuda, Blue, Fescue)	Mites	.	
Jobo	Thrips	.	Repeat as necessary to maintain control
Millet			
Rice			
Sorghum			
Wheat			
Cotton	Aphids	1-2	Apply in 25-100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment.
	Fleahoppers	.	
	Loopers	.	
	Mites	.	
	Pink Boll Worms	.	Apply in intervals of 7 days or less
	Thrips	.	Repeat as necessary to maintain control.
Dry Beans	Aphids	1-2	Apply as necessary to maintain control
Peanuts	Bean Beetles	.	
Soybeans	Beet Webworms	.	
Sugarbeets	Caterpillars	.	
Sunflowers	Loopers	.	
	Mexican Bean Beetles	.	
	Mites	.	
	Thrips	.	

**MISCELL**

Crops  
Ornamentals

The re  
for ag  
on far  
In clo

•IN BU  
To contr  
mosqui  
above li  
material  
to fifteen

•ON I  
To con  
to 2 ta

**BEST COPY AVAILABLE**

668

**MISCELLANEOUS**

Crops	Insects	Rate Pints/Acre	Additional Instructions
Ornamentals	Aphids	1-2	Apply in 25-100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment
	Armyworms	.	
	Caterpillars	.	Apply in intervals of 7 days or less.
	Crunch Bugs	.	
	Flashoppers	.	
	Fruit Flies	.	Repeat as necessary to maintain control
	Japanese Beetles	.	
	Leafhoppers	.	
	Leaf-miners	.	
	Leafrollers	.	
	Loopers	.	
	Lygus Bugs	.	
	Mealy Bugs	.	
	Mites	.	
	Plant Bugs	.	
Thrips	.		
White Flies	.		

*-Non-Agricultural Uses continued-*

Make treatment as follows:

- 1.) Apply liberally to tomatoes and fruits on baskets, on trucks or in plants. Use sprayers at a high pressure for applying at the rate of five or six pints of the diluted spray to a two-ton load of tomatoes. Direct the spray for maximum coverage of the baskets or hampers. It is important to spray between and beneath the containers.
- 2.) Spray the raw stock stacked in the yard.
- 3.) Dip the baskets in the diluted spray, after dumping the fruit, to kill adhering larvae and pupae.

**-AS A GRAIN PROTECTANT-**

Pyrethrin E.C. may be used to protect against grain storage insects at a dilution rate of 1 part Pyrethrin E.C. with 15 parts of water (1 pint with 1 gallon 7 pints of water). Apply directly to the grain at a rate of 4 to 5 gallons of diluted material per 1000 bushels as the grain is carried along a belt or as it enters the auger or elevator. Pyrethrin E.C. may be used in combination with a registered fumigant for use on heavily infested stored products.

Monthly inspections should be made. If the top two or three inches are found to be reinfested, re-treat, applying at the rate of 1 to 2 gallons of diluted material per 1000 bushels of stored product.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

In closed or confined areas the applicator should vacate treated areas and ventilate before reoccupying.

**STORAGE AND DISPOSAL**

**STORAGE:** Keep out of heat and cold. Store only in the original container. Keep away from fertilizer, food, other pesticides & feed. Keep out of reach of children & animals.

**DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**NON-AGRICULTURAL USES**

**• IN BARN, MILKING PARLORS, MILK ROOMS, DAIRIES AND POULTRY HOUSES**

To control house flies, horn flies, stable flies, horse flies, deer flies, biting and sucking lice, poultry lice, bedbugs, mites, fleas, ticks, mosquitoes, and gnats dilute at the rate of 4 fluid ounces per gallon of water and apply as a fog or fine mist directing the spray above livestock and poultry toward the ceiling and upper corners until the area is filled with mist using about two ounces of diluted material per 1000 cubic feet of space. For best results, close doors and windows before spraying and keep them closed for ten to fifteen minutes. Applicator should vacate treated area and ventilate before reoccupying. Repeat application as necessary.

**• ON HARVESTED TOMATOES AND FRUIT (Including Grapes)**

To control vinegar flies and fruit flies dilute at the rate of 1 part Pyrethrin E.C. with 600 parts of water (1.5 pints per 100 gallons or 2 tablespoons per 5 gallons of water). Keep the emulsion thoroughly mixed in the spray tank.

**WARRANTY**

WEBB WRIGHT CORPORATION warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of WEBB WRIGHT CORP. In no case shall WEBB WRIGHT CORP. be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. WEBB WRIGHT CORP. MAKES NO WARRANTY, EXPRESS OR IMPLIED, INCLUDING THE MERCHANTABILITY AND/OR FITNESS FOR ANY PARTICULAR PURPOSE, CONCERNING THIS MATERIAL, EXCEPT THOSE WHICH ARE CONTAINED ON THE LABEL.



P.O. BOX 1572  
FT. MYERS, FL 33902  
EPA REG. NO. 30573-2

PY DIR 1-95

*-con. next page-*

**BEST COPY AVAILABLE**

6

b  
b  
b

Webb Wright Corp.  
P.O. Box 1572  
Fort Meyers, FL 33902

38

Attention: Mr. William Wright

Subject: Pyrellin E.C.  
EPA Registration No. 30573-2  
Your amended application of March 3, 1995  
Your letters of March 29 and 30, 1995

The amended application referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you make the following changes:

A. Confidential Statement of Formula (CSF)

The Confidential Statement of Formula (CSF), dated April 22, 1994, is acceptable. A copy of this CSF has been placed in our file for the subject product. All previous CSF's are obsolete.

B. Revised Labeling

Submit three (3) copies of revised labeling with the following changes:

1. Revise your "Hazards to Humans and Domestic Animals" statement as follows:

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

149

2. Change the "Statement of Practical Treatment" heading to "First Aid" and modify the "IF SWALLOWED" and "IF ON SKIN" statements as follows:

#### FIRST AID

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

3. Add the following information at the end of your section on "Personal Protective Equipment":

#### User Safety Recommendations:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

4. Revise the "Environmental Hazards" section as follows:

#### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

5. Under "DIRECTIONS FOR USE", delete "General Classification".

6. Under "Agricultural Uses", add the following section:

#### Use Restrictions:

Pyrellin E.C. may be used to control the pests listed below at the indicated sites. Do not apply within one day of harvest. Wash any residue from vegetables before cooking.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store only in original container in a dry place inaccessible to children and pets

**PESTICIDE DISPOSAL:**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** [Use the appropriate statement on the attached, Appendix A on Page 7 of PR Notice 83-3.]

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,



Robert A. Forrest  
Product Manager (14)  
Insecticide-Rodenticide Branch  
Registration Division (H7504C)

- Enclosures:**
- 1. Stamped label
  - 2. A-79 Enclosure
  - 3. Appendix A, Page 7 of PR Notice 83-3

Peacock WP#5:F:\USER\DPEACOCK\30573-2.ROT:305-5407,-6600:4/4/95

4 9 9

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING ▷

# Pyrellin® E.C.

**EMULSIFIABLE CONCENTRATE INSECTICIDE**

**ACTIVE INGREDIENTS:**

Pyrethrins .....	0.60%
Rotenone .....	0.50%
Other Cube Resins .....	0.50%
<b>INERT INGREDIENTS .....</b>	<b>98.40%</b>
<b>TOTAL .....</b>	<b>100.00%</b>

**Contents Per Gallon:**

0.042 lbs. of Pyrethrins
0.035 lbs. of Rotenone
0.335 lbs. of Other Cube Resins

**Keep out of reach of children**

**Caution**

SEE INSIDE OF BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

**NET CONTENTS: 2.5 GALLONS**

EPA Reg. No. 30573-2  
 EPA Est. No. 37429-GA-01<sup>®</sup>, 655-GA-01<sup>®</sup>, 789-GA-01<sup>®</sup>  
 Superscript used is first letter of lot number.

Manufactured by:

**WEBB WRIGHT CORPORATION**

P.O. Box 1572  
P. Myers, FL 33902

1-95

ACCEPTED  
 with COMMENTS  
 in EPA Letter Dated  
**APR 4 1995**

Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act  
 as amended, the pesticide  
 registered under EPA Reg. No.  
30573-2

4

**STOP!**  
**READ ALL PRECAUTIONS AND DIRECTIONS FOR USE BEFORE USING**

**PRECAUTIONARY STATEMENTS**  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Contact a physician at once.  
**IF INHALED:** Remove from contact & seek medical attention.  
**IF ON SKIN:** Flush immediately with copious amounts of water.  
**IF IN EYES:** Flush immediately with copious amounts of water. If irritation develops seek medical attention.

**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin or eyes. In case of contact, immediately flush eyes or skin with plenty of water. For eyes get medical attention if irritation develops or persists.

**Personal Protective Equipment:**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category D on an EPA chemical resistance category selection chart.

- 1) Barrier Laminate
- 2) Butyl Rubber

**Applicators and other handlers must wear:**

- 1) Long-sleeved shirt and long pants
- 2) Chemical-resistant gloves, such as Barrier Laminate or Butyl Rubber
- 3) Shoes plus-socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering control statements:**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (D) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

Flammable. Keep away from heat & open flame. Do not use with thermal fogging units.

**DIRECTIONS FOR USE**  
**GENERAL CLASSIFICATION**

It is a violation of Federal law to use this product in a manner inconsistent with its label, state or tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard contains requirements for the protection of agricultural workers on farms and handlers of agricultural pesticides. It contains requirements for training, decontamination, assistance. It also contains specific instructions and exceptions pertaining to the protective equipment (PPE) and restricted-entry interval. The requirements in this section that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift, when they are in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI).

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard with anything that has been treated, such as plants, soil, or water is:

- 1) Coveralls
- 2) Chemical-resistant gloves, such as Barrier Laminate or Butyl Rubber
- 3) Shoes plus socks

**AGRICULTURAL USES**

**•USED ALONE**

Pyrethrin E.C. can be used at rates of 1 to 2 pints per acre in 25 to 100 gallons of water during growing season including use as a pre-harvest insecticide where a 12 hour REI application is allowed. Pyrethrin E.C. should be applied in rates of 1 to 2 pints per acre as needed for control. Constant spray tank agitation is required during spraying.

Superior control may be obtained by applying Pyrethrin E.C. in Low-Volume, small droplet (weather conditions) sprays of 1 to 2 pints Pyrethrin E.C. with 1 to 2 gallons of water or 60 micron) sprays of 1 to 2 pints Pyrethrin E.C. with 1 to 2 pints of vegetable oil which sprays may also minimize risk of phytotoxicity.

ULV sprays (recommended particle size 50-75 micron) applied neat (vegetable oil dilution) of leaf-miner.

**•COMBINED WITH OTHER INSECTICIDES**

Pyrethrin E.C. may be combined with other insecticides for a quicker and more complete control.

**BEST COPY AVAILABLE**

-continued-

problem, and as an insect "exciter" to flush insects out of hiding and into contact with other spray residues. The application should conform to accepted use precautions and directions for both products. Pyrethrin E.C. may be tank mixed at rates from 4 to 16 ounces per acre. Products with which it may be mixed include, but are not limited to, abamectin, azinphos-methyl, Bacillus thuringiensis, carbaryl, chlorpyrifos, insecticidal soaps\*, methomyl, and synthetic pyrethroids.

\* Local water conditions may affect compatibility with insecticidal soap products.

**"SPREADER/STICKER" ADJUVANTS**

If use of a "spreader/sticker" spray adjuvant is desired to improve spray efficiency (often resulting in increased overall efficacy), use a natural vegetable oil product (typically an "85/15" composition with 85% soy bean or cottonseed oil and 15% inerts including emulsifiers). Use of petroleum oil adjuvants increases risk of phytotoxicity.

**ADULTICIDE CONTROL**

May be used on croplands as an adulticide to protect agricultural workers during harvesting when used according to the requirements contained in the Agricultural Use Requirements box on the preceding page. Use at 1/4 to 1 pint per acre when applied by fixed wing aircraft, helicopters or truck mounted equipment.

**USE THROUGH IRRIGATION SYSTEMS**

Apply Pyrethrin E.C. only through sprinkler systems containing anti-siphon and check valves which will prevent water source contamination and overflow of the mix tank and containing interlocking controls between the metering device and the water pump to insure simultaneous shutoff. Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application. Inject Pyrethrin E.C. with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled or drained.

Accurate calibration and distribution are essential for insect control. Crop damage may result from inaccurate calibration or distribution.

**Growing Crops  
(Outdoors and in Greenhouses)**

**VEGETABLE CROPS**

Crop	Insect	Rate Pints/Acres	Additional Instructions
Asparagus	Aphids	1-2	Apply in 25 to 100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment.
Beans (Green Lima, Mung, Snap)	Asparagus Beetles	.	
Beets (red)	Alfalfa Webworms	.	Apply in intervals of 7 days or less.
Cabbage (including Chinese, NAPA, red, sav)	Bean Beetles	.	
Carrots	Beet Webworms	.	Repeat as necessary to maintain control
Celery	Cabbageworms	.	
Corn	Celery Leafhoppers	.	
Cucumbers	Garden Fleahoppers	.	
Eggplant	Hornworms	.	

-continued next page-

-Vegetable Crops continued-

Crop	Insect	Rate Pints/Acres	Additional Instructions
Lettuce (Head Lettuce, Leaf, Buttercrunch, Romaine)	Japanese Beetles	1-2	
Onions (Green, Dry Bulb)	Leafhoppers	.	
Peppers	Leaf-Miners	.	
Potato	Looper	.	
Spinach	Lygus Bugs	.	
Squash (Summer, Winter, Acorn, Butternut, Crookneck, Hubbard, Spaghetti, Zucchini)	Mites	.	
Tomato	Oblique-banded Leafrollers	.	
	Plant Bugs	.	
	Stink Bugs	.	
	Thrips	.	
	White Flies	.	
	Colorado Potato Beetles	1 1/2 - 2	3 to 4 applications or more as eggs start to hatch and repeat at 5 to 7 day intervals.
	Cucumber Beetles	.	
	European Corn Borers	.	
	Flea Beetles	.	
	Squash Bugs	.	
	Squash Vine Borers	.	
	Vegetable Weevils	.	
Artichokes (Globe, Jerusalem)	Aphids	1-2	Apply as necessary to maintain control
Broccoli (Chinese, rib)	Leafhoppers	.	
Brussels Sprouts	Leaf-Miners	.	
Bok Choy (Kai Choy, Pak Choi, Pak Choy, Pak Toy)	Looper	.	
Cardoon	Lygus Bugs	.	
Cassavas (Yucca)	Mites	.	
Cauliflower	Plant Bugs	.	
Celeriac	Thrips	.	
Chic Peas	White Flies	.	
Collards			
Dakota			
Endive			
Escarole			
Garlic			
Greens (Dandelion, Turnip, Mustard, Beet, Chino, Poke)			
Hops			
Horseradish			
Kale			
Kohlrabi			
Leeks			
Lentils			
Malanga			
Melons (Cantaloupe, Casabas, Crenshaws, Honeydews, Muskmelons, Persian, Watermelons)			
Okra			
Parsnip			
Pears (Field, Blackeye, Green, Snow)			
Pumpkins			
Radicchio			

-Vegetable Crops continued next page-

BEST COPY AVAILABLE

629



*-Vegetable Crops continued-*

Crop	Insect	Rate Pints/Acre	Additional Instructions
Radish			
Rhubarb			
Rutabagas			
Salsify			
Shallots			
Swiss Chard			
Taro (Dash-n)			
Turnip			
Yams			

**FRUIT, NUT & VINE CROPS:**

Crop	Insect	Rate Pints/Acre	Additional Instructions
Apples	Aphids	1-2	Apply in 25 to 100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment.
Blackberries	Blueberry Maggots	.	
Blueberries	Cherry Fruit Flies	.	
Cherries	Gooseberry Fruit Worms	.	
Currents	Imported Current Worms	.	
Gooseberries	Japanese Beetles	.	
Grapes	Leafhoppers	.	
Peaches	Mites	.	
Pears	Red Necked Borers	.	
Plums	Raspberry Fruit Worms	.	
Raspberries	Rose Chalcids	.	
	Thrips	.	
Strawberries	Mites	1-2	Apply as necessary to maintain control
	Strawberry Root Worms	.	
	Strawberry Leafrollers	.	
	Thrips	.	
Citrus	Aphids	1-2	Apply as necessary to maintain control
	Caterpillars	.	
	Fuller Rose Beetle	.	
	Thrips	.	
	White Flies	.	
Acerola	Aphids	1-2	Apply as necessary to maintain control
Anon	Caterpillars	.	
Apricots	Fireworms	.	
Atemoya	Fruit Flies	.	
Avocados	Leafhoppers	.	
Boysenberries	Mites	.	
Cactus Fruit	Thrips	.	
Calamondin	Weevils	.	
Carambola	White Flies	.	
Chirimoya			
Crab Apples			

*-Fruit, Nut & Vine Crops continued next page-*

**MISCELLANEOUS**

Crops	Insects	Rate Pints/Acre	Additional Instructions
Ornamentals	Aphids	1-2	Apply in 25-100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with U.V spray equipment.
	Armyworms	.	
	Caterpillars	.	
	Chinch Bugs	.	
	Fleahoppers	.	
	Fruit Flies	.	
	Japanese Beetles	.	
	Leafhoppers	.	
	Leaf-miners	.	
	Leafrollers	.	
	Loopers	.	
	Lygus Bugs	.	
	Mealy Bugs	.	
	Mites	.	
Plant Bugs	.		
Thrips	.		
White Flies	.		

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

In closed or confined areas the applicator should vacate treated areas and ventilate before reoccupying.

**NON-AGRICULTURAL USES**

**•IN BARN, MILKING PARLORS, MILK ROOMS, DAIRIES AND POULTRY HOUSES**

To control house flies, horn flies, stable flies, horse flies, deer flies, biting and sucking lice, poultry lice, bedbugs, mites, fleas, ticks, mosquitoes, and gnats dilute at the rate of 4 fluid ounces per gallon of water and apply as a fog or fine mist, directing the spray above livestock and poultry toward the ceiling and upper corners until the area is filled with mist using about two ounces of diluted material per 1000 cubic feet of space. For best results, close doors and windows before spraying and keep them closed for ten to fifteen minutes. Applicator should vacate treated area and ventilate before reoccupying. Repeat application as necessary.

**•ON HARVESTED TOMATOES AND FRUIT (Including Grapes)**

To control vinegar flies and fruit flies dilute at the rate of 1 part Pyrethrin E.C. with 600 parts of water (1.5 pints per 100 gallons or 2 tablespoons per 5 gallons of water). Keep the emulsion thoroughly mixed in the spray tank.

*-continued next page-*

**BEST COPY AVAILABLE**

ACCEPTED  
with COMMENTS  
in EPA Letter Dated  
**APR 4 1995**