



EMULSIFIABLE CONCENTRATE INSECTICIDE

ACTIVE INGREDIENTS:

Pyrethrins	0.60%
Rotenone	0.50%
Other Associated Resins	0.50%
OTHER INGREDIENTS	98.40%
TOTAL	100.00%

Contents Per Gallon

0.042 lbs. of Pyrethrins
0.035 lbs. of Rotenone
0.035 lbs. of Other Associated Resins

Manufactured by:
WEBB WRIGHT CORPORATION

P.O. Box 1572
Fl. Myers, FL 33902

Keep out of reach of children

CAUTION

SEE ACCOMPANYING BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

FIRST AID

- IF SWALLOWED:** Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF INHALED:** Remove from contact & seek medical attention.
- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.
- IF IN EYES:** Flush immediately with copious amounts of water. If irritation develops seek medical attention.


ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water by cleaning equipment or disposal of wastes.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NET CONTENTS: 2.5 GALLONS

EPA Reg. No. 30573-2
EPA Est. No. 50591-GA-01[®], 655-GA-01[®], 48498-CA-01[®]
Superscript used is first letter of lot number.





CONTENTS MAY HAVE SETTLED.
SHAKE BEFORE USING.

PEEL BACK BOOK HERE AND
RESEAL AFTER OPENING



EMULSIFIABLE CONCENTRATE INSECTICIDE

ACTIVE INGREDIENTS:

Pyrethrins	0.60%	Contents Per Gallon	
Rotenone	0.50%	0.042 lbs. of Pyrethrins	
Other Associated Resins	0.50%	0.035 lbs. of Rotenone	
OTHER INGREDIENTS	98.40%	0.035 lbs. of Other Associated Resins	
TOTAL	100.00%		

Keep out of reach of children

CAUTION

SEE INSIDE OF BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

NET CONTENTS: 2.5 GALLONS

EPA Reg. No. 30573-2

EPA Est. No. 50591-GA-01 655-GA-01 48498-CA-01

Superscript used is first letter of job number.

Manufactured by

**WEBB WRIGHT
CORPORATION**


P.O. Box 1572
Et. Myers, FL 33902

11-00

ACCEPTED

APR 23 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 30573-2





STOP! Read All Precautions & Directions Before Using

PRECAUTIONARY STATEMENTS

Keep Out of Reach of Children

CAUTION

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious person.
IF INHALED: Remove from contact & seek medical attention.
IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.
IF IN EYES: Flush immediately with copious amounts of water. If irritation develops seek medical attention.

HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin or eyes. In case of contact, immediately flush eyes or skin with plenty of water. For eyes get medical attention if irritation develops or persists. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category D on an EPA chemical resistance category selection chart.

- 1) Barrier Laminate
- 2) Butyl Rubber

Applicators and other handlers must wear:

- 1) Long sleeved shirt and long pants
- 2) Chemical-resistant gloves, such as Barrier Laminate or Butyl Rubber
- 3) Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Engineering control statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (D) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment/washwaters.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involve contact with anything that has been treated, such as plants, soil, or water is:

- 1) Coveralls
- 2) Chemical-resistant gloves, such as Barrier Laminate or Butyl Rubber
- 3) Shoes plus socks

AGRICULTURAL USES

~~USE RESTRICTIONS~~

Pyrethrin E.C. may be used to control the pests listed below at the indicated sites. Do not apply within 12 hours of harvest.

~~USED ALONE~~

Pyrethrin E.C. can be used at rates of 1 to 2 pints per acre in 25 to 100 gallons of water by ground sprayers at any time during the growing season including use as a pre-harvest insecticide where a 12 hour REI is maintained. In those areas where aerial application is allowed, Pyrethrin E.C. should be applied in rates of 1 to 2 pints per acre with a minimum of 5 gallons of water. Repeat as needed for control. Constant spray tank agitation is required during spraying when diluting with water.


Superior control may be obtained by applying Pyrethrin E.C. in Low-Volume, small droplet (70-80 micron, depending on existing weather conditions) sprays at the 1 to 2 pint per acre rate with 1 to 2 gallons of water or in ULV (ultra-low volume) small droplet (40-60 micron) sprays at the 1 to 2 pint per acre rate with 1 to 2 pints of vegetable oil where such sprays are approved.

Smaller droplet sprays can often be obtained with conventional sprayers by adjusting to a higher pressure setting (250 - 400 lbs). Small droplet sprays also minimize risk of phytotoxicity.

~~COMBINED WITH OTHER INSECTICIDES~~

Pyrethrin E.C. may be combined with other insecticides for a quicker and more complete kill where insect resistance may be a problem, and as an insect "exciter" to flush insects out of hiding and into contact with other spray residues. The application should conform to accepted use precautions and directions for both products. Pyrethrin E.C. may be tank mixed at rates from 4 to 32 ounces

-continued next page-





-Agricultural Uses continued-

per acre. Pyrellin E.C. is compatible with most widely used insecticides*, though test applications for phytotoxicity are advised when mixing with products containing petroleum distillates.

* Local water conditions may affect compatibility with insecticidal soap products.

"SPREADER/STICKER" ADJUVANTS

Two thirds of Pyrellin E.C.'s volume by weight is comprised of a soybean oil base. This in combination with a 7 1/2% by weight volume of emulsifiers/surfactants will in many cases provide adequate spreader/sticker performance. If additional spreader/sticker capability is desired, use a natural vegetable oil product (typically an "85/15" composition with 85% soybean or cottonseed oil and 15% inerts including emulsifiers). With higher concentrations of vegetable oils, avoid large droplet sprays to lessen risk of phytotoxicity by using higher spray pressures (250-400 lbs). Use of petroleum oil adjuvants is not recommended due to risk of phytotoxicity.

MOSQUITO CONTROL

May be used on croplands as an adulticide to protect agricultural workers during harvesting when used according to the requirements contained in the Agricultural Use Requirements box on the preceding page. Use at 1/4 to 1 pint per acre when applied by fixed wing aircraft, helicopters or truck mounted equipment.

USE THROUGH IRRIGATION SYSTEMS

Apply Pyrellin E.C. only through sprinkler systems containing anti-siphon and check valves which will prevent water source contamination and overflow of the mix tank and containing interlocking controls between the metering device and the water pump to insure simultaneous shutoff. Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application. Inject Pyrellin E.C. with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled or drained.


Accurate calibration and distribution are essential for insect control. Crop damage may result from inaccurate calibration or distribution.

**Growing Crops
(Outdoors and in Greenhouses)**

VEGETABLE CROPS:

Crop	Insect	Rate Pints/Acres	Additional Instructions
Asparagus	Aphids	1-2	Apply in 25 to 100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment.
Beans (Green, Lima, Mung, Snap)	Asparagus Beetles		
Beets (red)	Alfalfa Webworms		Apply in intervals of 7 days or less.
Cabbage (including Chinese, NAPA, red, su)	Bean Beetles		
Carrots	Beet Webworms		Repeat as necessary to maintain control.
Celery	Cabbageworms		
Corn	Celery Leafhoppers		
Cucumbers	Garden Fleahoppers		
Eggplant	Harlequin Bugs		

-continued next page-





-Vegetable Crops continued-

Crop	Insect	Rate Pints/Acres	Additional Instructions
Lettuce (Head Lettuce, Leaf, Buttercrunch, Romaine)	Japanese Beetles	1-2	
Onions (Green, Dry Bulb)	Leafhoppers	.	
Peppers	Leaf-Miners	.	
Potato	Looper	.	
Spinach	Lygus Bugs	.	
Squash (Summer, Winter, Acorn, Butternut)	Mites	.	
Crookneck, Hubbard, Spaghetti, Zucchini	Oblique-banded Leafrollers	.	
Tomato	Plant Bugs	.	
	Stink Bugs	.	
	Thrips	.	
	White Flies	.	
	Colorado Potato Beetles	1 1/2 -2	3 to 4 applications or more as eggs start to hatch and repeat at 5 to 7 day intervals
	Cucumber Beetles	.	
	European Corn Borers	.	
	Flea Beetles	.	
	Squash Bugs	.	
	Squash Vine Borers	.	
	Vegetable Weevils	.	
Artichokes (Globe, Jerusalem)	Aphids	1-2	Apply as necessary to maintain control
Broccoli (Chinese, rat)	Leafhoppers	.	
Brussels Sprouts	Leaf-Miners	.	
Box Choy (Kai Choy, Pak Choy, Pak Choy, Pak Toy)	Looper	.	
Cardoon	Lygus Bugs	.	
Cassava (Yucca)	Mites	.	
Cauliflower	Plant Bugs	.	
Celery	Thrips	.	
Chic Peas	White Flies	.	
Collards			
Daikon			
Endive			
Escarole			
Garlic			
Greens (Dandelion, Turnip, Mustard, Beet, Chins, Poke)			
Hops			
Horseradish			
Kale			
Kohlrab-			
Lentils			
Lentils			
Melanga			
Melons (Cantaloupe, Casabas, Grenshaws, Honeydews)			
Muskmelons, Persian, Watermelons			
Okra			
Parsnip			
Peas (Field, Blackeye, Greer, Snow)			
Pumpkins			
Radicchio			

-Vegetable Crops continued next page-






-Vegetable Crops continued-

Crop	Insect	Rate Pints/Acre	Additional Instructions
Radish			
Rhubarb			
Rutabagas			
Salsify			
Shallots			
Swiss Chard			
Taro (Dasheen)			
Turnip			
Yams			

FRUIT, NUT & VINE CROPS:

Crop	Insect	Rate Pints/Acre	Additional Instructions	
Apples	Aphids	1-2	Apply in 25 to 100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment.	
Blackberries	Blueberry Maggots			
Blueberries	Cherry Fruit Flies			
Cherries	Gooseberry Fruit Worms			
Currants	Imported Currant Worms			
Gooseberries	Japanese Beetles			
Grapes	Leafhoppers			Apply in intervals of 7 days or less.
Peaches	Mites			Repeat as necessary to maintain control.
Pears	Red Necked Borers			
Plums	Raspberry Fruit Worms			
Raspberries	Rose Chafers			
	Thrips			
Strawberries	Mites	1-2	Apply as necessary to maintain control.	
	Strawberry Root Worms			
	Strawberry Leafrollers			
	Thrips			
Carus	Aphids	1-2	Apply as necessary to maintain control.	
	Caterpillars			
	Fuller Rose Beetle			
	Thrips			
	White Flies			
Acerola	Aphids	1-2	Apply as necessary to maintain control.	
Anon	Caterpillars			
Apricots	Fireworms			
Atemoya	Fruit Flies			
Avocados	Leafhoppers			
Boysenberries	Mites			
Cactus Fruit	Thrips			
Calamondin	Weevils			
Carambola	White Flies			
Chirimoya				
Crab Apples				

-Fruit, Nut & Vine Crops continued next page-






-Fruit, Nut & Vine Crops continued-

Crop	Insect	Rate Pints/Acre	Additional Instructions
Dates (also Chinese):			
Dewberries			
Elderberries			
Guava			
Huckleberries			
Kiwi			
Loganberries			
Longan Fruit			
Loquat			
Mango			
Mulberries			
Papayas			
Passion Fruit			
Persimmons			
Pineapple			
Pitanga Cherry			
Prickly Pear			
Quince			
Sapodilla			
Sapote			
Almonds	Aphids	1-2	Apply in 25 to 100 gallons of water as needed for control
Beechnuts	Mites	.	
Butternuts	Thrips	.	
Cashews	White Flies	.	
Chestnuts		.	
Filberts		.	
Hazelnuts		.	
Hickory Nuts		.	
Lycnee		.	
Macadamia Nuts		.	
Pecans		.	
Pistachio		.	
Pomegranate		.	
Tamarind		.	
Walnuts		.	

HERBS & SPICES:

Crop	Insect	Rate Pints/Acre	Additional Instructions
Anise	Aphids	1-2	Apply in 25 to 100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment.
Basil	Loobers	.	
Caraway	Mites	.	
Chicory	Plant Bugs	.	
Chives	Thrips	.	
Corander (Cilantro)	White Flies	.	

-herbs & Spices continued next page-






-Herbs & Spices continued-

Crop	Insect	Rate Pints/Acre	Additional Instructions
Cumin Dill Fennel Ginger Ginseng Marjoram Mint Mustard Oregano Paprika Parsley Rosemary Sage Savon Tarragon Thyme			Repeat as necessary to maintain control.

FIELD CROPS:

Crop	Insect	Rate Pints/Acre	Additional Instructions
Alfalfa Barn: Clover Field Corn Grass (Bermuda, Blue, Fescue) Jojoba Millet Rice Sorghum Wheat	Aphids Armyworms Chinch Bugs Flies Mites Thrips	1-2	Apply in 25-100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment. Apply in intervals of 7 days or less. Repeat as necessary to maintain control.
Cotton	Aphids Fleahoppers Loopers Mites Pink Boll Worms Thrips	1-2	Apply in 25-100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment. Apply in intervals of 7 days or less. Repeat as necessary to maintain control.
Dry Beans Peanuts Soybeans Sugarbeets Sunflowers	Aphids Bean Beetles Beet Webworms Caterpillars Loopers Mexican Bean Beetles Mites Thrips	1-2	Apply as necessary to maintain control.





MISCELLANEOUS:

Crops	Insect	Rate Pints/Acre	Additional Instructions
Ornamentals	Aphids	1-2	Apply in 25-100 gallons of water by ground sprayer or in 1-2 gallons of water by Low-Volume apparatus, or in 1-2 pints of vegetable oil if applied with ULV spray equipment. Apply in intervals of 7 days or less. Repeat as necessary to maintain control.
	Armyworms		
	Caterpillars		
	Chinch Bugs		
	Flea-hoppers		
	Fruit Flies		
	Japanese Beetles		
	Leafhoppers		
	Leaf-miners		
	Leaf-rollers		
	Loopers		
	Lygus Bugs		
	Mealy Bugs		
	Mites		
	Plant Bugs		
Thrips			
White Flies			

STORAGE AND DISPOSAL

STORAGE: Keep out of heat and cold. Store only in the original container. Keep away from fertilizer, food, other pesticides & feed. Keep out reach of children & animals.

DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

WEBB WRIGHT CORPORATION warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of WEBB WRIGHT CORP. In no case shall WEBB WRIGHT CORP. be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. WEBB WRIGHT CORP. MAKES NO WARRANTY, EXPRESS OR IMPLIED, INCLUDING THE MERCHANTABILITY AND/OR FITNESS FOR ANY PARTICULAR PURPOSE, CONCERNING THIS MATERIAL, EXCEPT THOSE WHICH ARE CONTAINED ON THE LABEL.



P.O. BOX 1572
FT. MYERS FL 33902
EPA REG. NO 30573-2

