

LC 1231 Disinfectant - Cleaner

**FUNGICIDAL
TUBERCULOCIDAL
CONDUCTIVE**

DESCRIPTION

A one-step cleaner and disinfectant-deodorant of proven effectiveness. The combination of multiple germicides and high active ingredients provides an effective one-step method of disinfecting and cleaning at a dilution of one ounce to one gallon water (1:128).

LC 1231 Disinfectant-Cleaner removes mildew stains from hard surfaces.

LC 1231 Disinfectant-Cleaner is conductive - - may be used on conductive floorings as it does not interfere with electrical conductivity, as specified in the procedures in accordance with Paragraph 252, "Conductive Flooring of the National Board of Fire Underwriters", Specification No. 56.

In the AOAC Use-Dilution Method, 11th Edition, LC 1231 Disinfectant-Cleaner kills the following organisms at 1 oz. per gallon of water (1:128) dilution: Staphylococcus aureus, Salmonella choleraesuis, Salmonella pullorum, Salmonella typhosa, Pseudomonas aeruginosa, Proteus vulgaris, Streptococcus mitis, and Escherichia coli.

EPA Reg. no. 30943-2 EPA Est. 30943-FL-1

AOAC Tuberculocidal Test, 11th Edition
Mycobacterium tuberculosis at 1:128

AOAC Official Fungicidal Test, 11th Edition
Trichophyton interdigitale at 1:128

DOT: Corrosive Liquid, NOS

ICC: Disinfectant in bulk, o/t medicinal

Lea Chemicals, Inc.

P. O. Box 868
Marianna, Florida 32446

ACTIVE INGREDIENTS

Isopropyl alcohol, 10.82%; Tetrasodium ethylenediaminetetraacetate, 4.00%; Sodium xylene sulfonate, 3.20%; Sodium lauryl sulfate, 3.00%; Sodium o-benzyl-p-chlorophenate, 2.61%; Sodium p-tertiary-butylphenate, 1.98%; Sodium o-phenylphenate, 1.54%; Sodium 4-chloro-2-cyclopentylphenate, 0.50%.

INERT INGREDIENTS 72.35%

**DANGER
KEEP OUT OF REACH OF CHILDREN**

Corrosive. Causes eye damage and severe skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

For use in sanitation in meat and poultry production - food processing - hospitals - animal and poultry care - veterinary.

Net Contents _____ Gallon

J-73

DIS-2

1. For disinfecting and cleaning hard surfaces in meat and poultry production, such as in processing of meat, poultry, turkeys, and equipment, dilute LC 1231 one ounce to one gallon of water (1:128). Wipe, mop or swab all surfaces with solution. Leave in contact 10 minutes. Surface coming in contact with food must be thoroughly scrubbed with detergent and rinsed with potable water before reuse.

**STAPHYLOCIDAL
PSEUDOMONACIDAL
SALMONELLACIDAL**

2. For general cleanup in food processing plants, patient rooms, surgery, nursing homes, all animal facilities, on walls, floors, hard surfaces, equipment. Use one ounce per gallon of water (1:128). Cover surfaces thoroughly with solution and allow to remain in contact 10 minutes. For porous surfaces, increase concentration to two ounces per gallon. Surfaces coming in contact with food must be thoroughly scrubbed with detergent and rinsed with potable water before reuse.
3. For non-food disinfection of laying and broiler houses and hatcheries; DO NOT use in milking stalls, milking parlors, or milk houses. Remove all animals, poultry and feeds from the premises. Remove all litter and manure from floors and other surfaces. Empty all feeding and watering troughs and appliances. Saturate all surfaces with a 1:128 solution. Ventilate building and other closed spaces. Do not house poultry, livestock or employ equipment until treatment has been absorbed or dried. All treated food racks, waterers and fountains must be thoroughly scrubbed with detergent and rinsed with potable water prior to reuse.
4. Remove gross filth and heavy soil deposits from surfaces before disinfecting and deodorizing garbage and trash receptacles, carts and sick room equipment. Use one ounce of LC 1231 to one gallon of water (one part to 128 water.)
5. For fungicidal activity to aid in preventing the spread of athlete's foot fungi on environmental surfaces, use one ounce per gallon of water; mop or swab this solution onto floors and other similar surfaces normally contacted by the feet in locker rooms, shower stalls, bathrooms and around swimming pools.

ACCEPTED

JUN 28 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 30943-2