

ACCEPTED

OCT 11 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 30943-5

LEA PINE SCENTED DISINFECTANT

DESCRIPTION

Pine Scented Disinfectant and deodorant kills Salmonella, Staphylococci and Pseudomonas organisms. Use in bathrooms, showers, on floor and walls in schools, municipal buildings and offices.

AOAC USE-DILUTION, 11th EDITION, 1970

- 1. Staphylococcus aureus 1:64
- 2. Salmonella choleraesuis 1:64
- 3. Pseudomonas aeruginosa (PRD-10) 1:64

AOAC PHENOL COEFFICIENT, 11th EDITION

Salmonella typhosa 5

Lea Chemicals

P. O. Box 868
Marianna, Florida 32446

ACTIVE INGREDIENTS:

Isopropanol, 10.75%; Pine oil, 6.00%; Soap, 5.90%; Potassium 4-and 6-Chloro-2-phenylphenate, 3.90%; Potassium ortho-benzyl-para-chlorophenate, 2.45%; Tetrasodium ethylene diamine tetraacetate, 0.20%.

INERT INGREDIENTS 70.80%

EPA Reg. No. 30943-5

WARNING

KEEP OUT OF REACH OF CHILDREN

May cause eye damage. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

Rinse empty container thoroughly with water and discard it.

Net Contents _____ Gallon

H 73

Dis 3

DIRECTIONS

Clean surfaces of gross dirt and filth with soap or detergent solution.

For General Disinfecting: Floors, tile and walls—mop or swab with a solution of 2 ounces Pine Scented Disinfectant per gallon of water. Allow contact time of 10 minutes, then rinse with water.

Trucks for handling animals and stock holding pens: DO NOT use in milking stalls. Remove all animals and feeds from premises, cars, boats, trucks. Remove all litter and manure from floors, walls, and surfaces occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Saturate all surfaces with disinfecting solution. Ventilate closed spaces, and DO NOT house livestock or employ equipment until treatment has been absorbed, set, or dried. All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be thoroughly scrubbed with detergents, and rinsed with potable water prior to reuse.

As a Deodorant: Clean surfaces and use 1¼ ounce per gallon of water. Mop or swab on surface.