P11-13

1

ţ

30950-7 UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

30950-7

Until Reregistrat.on

Diazinon 25 % Emulsifiable Concentrate

10 5425227

SEPZYI

10+6

R. Maldonado & Co., Inc. 8810 Tenth Avenue North GPO Box 3231 San Juan, PR 00936

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFR, sec. 3(c)(5) or FIFRA sec. 4 when the Agency requires all registrants of similar products to submit such data.

2. Add the phrase"EPA Registration No. 30950-7" to your label before you release the product for shipment.

3. Submit five (5) copies of your final printed label ng before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed label ng.

If these conditions are not complied with, the registation will be subject to cancellation in accordance with FIFRA section 6 (e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy is enclosed for your records.



George T. LaFocca Product Manager (13) Insecticide-Rodenticide Branch Registration Division (117505C)

Enclosures

| CONCURRENCES | | | | | | | | |
|-------------------------|-------|--|---------------------------|--|--|--------------------|--|--|
| TYMBOL | | | | | | | | |
| | Moals | | | | | | | |
| DATE | | | | | | | | |
| EPA Form 1320-1A (1/90) | | | Printed on Resysted Paper | | | OFFICIAL FILE COPY | | |

"U.S. Government Printing Office: 1992 -- 620 656/40672

Diazinon

25% EMULSIFIABLE CONCENTRATE

Lawn and Garden Insect Control

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

| ACTIVE INGREDIENT: Diazinon: 0,0-Diethyl 0-(2-isopropyl-6-methyl | |
|---|--------|
| -4-pyrimidinyl)phosphorothioate* | |
| TOTAL: | 100.0% |

*Contains aromatic petroleum distillates.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional precautionary statements

PROTECTS

ĺ

Í

Lawns Vegetables Fruits and Nuts Roses, Flowers Trees, Shrubs

See full label in booklet under cap for complete directions for mixing and applying this product and a complete list of pests controlled.

EPA Reg. No. 30950-

NET CONTENTS: 8 oz.

Formulated By:

R. Maldonado & Co., Inc. Calle Julia Num. 2 Urb. Pinero Hato Rey, PR 00936 TEL. 753-7230

| CONTROLS | | |
|------------------------|-----------|---------------------------------------|
| Ants, Aphids | · · · · · | : : : |
| Chinch Bugs | • | · · · · · · · · · · · · · · · · · · · |
| Cutworms | • • • | |
| Fire Ants ¹ | • | ••••• |
| Flea Beetles | ••••• | • |
| Grubs, Mites | • | |
| Sod Webworms | | ••••• |
| and many others | • • | |

2046

¹ Aids in control

EPA Est, No. 30950-PR-1 A COMMENTS in EPA Lotter Dated

Under the Federal Insecticide, Fungleide, and Rodenticide Act on amended, for the peatregistered under EFA Reg. <u>30950</u> - 7

29

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

Diazinon 25% Emulsifiable Concentrate is a product which must be diluted with water before application. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables and fruits and insects which annoy pets and people.

SEE THE BOOKLET UNDER CAP FOR COMPLETE DIRECTIONS FOR USE AND PESTS CONTROLLED.

Notes: (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. SEE STORAGE AND DISPOSAL DIRECTIONS. (2) Do not use in the home. (3) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia or orchids since plant injury may occur. (4) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (5) Do no apply to food crops grown in green houses.

After spraying, wait the following number of days before harvesting fruits, nuts, or vegetables:

| <u>1 day</u> | | | |
|--------------|---|--|--|
| | Tomatoes | | |
| 3 days | Melons, Winter Squash | | |
| 5 days | Peppers, Strawberries | | |
| 7 days | Beans, Broccoli, Blackberrice, Boysenberries, Cauliflower, Cucumbers, Dewberries, Loganberries, Peas, Raspberries, Summer Squash, Sweet Corn | | |
| 10 days | Chinese Broccoli, Chinese Cabbage, Collards, Kale | | |
| 14 days | Carrots, Endive (Escarole), Lettuce, Onions (Bulb and Green), Parsnips, Radishes, Spinach, Turnips | | |
| 21 days | Apricots, Cabbage, Cherries, Nectarines, Peaches, Plums, Prunes | | |
| 28 days | Grapes | | |
| 35 days | Potatoes (Irish) | | |
| 45 days | Walnuts | | |
| • | Almonds (one dormant spray only) | | |

<u>الم</u>

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing h^{-2} rewearing again.

*This table is only a partial listing of the pests controlled by this product. See booklet under cap for complete directions for use and pests controlled.

| Problem Area | Pests* |
|------------------------------------|--|
| Nuisance Pests in Outside Areas | Ants, Brown Dog Ticks, Chiggers, Cock oaches, Crickets, Fleas, Fire Ants, Spiders and others* |
| Lawns | Chinch Bugs, Ants, Sod Webworms, Brown Dog Ticks, Fleas, White Grubs, Lawn Billbugs and others* |

| Vegetables: Foliar Application Beans, Peas (except dried varieties) | Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, M Beetles, Spider Mites | lexican Bean | |
|---|---|---------------------------------------|-----------|
| Broccoli, Chinese Broccoli, Cabbage, Chinese Cabbage, C-uliflower, Collards, Kale | Aphids, Flea Beetles, Dipterous Leafminers, Diamondbach Imported Cabbageworms | r Moths, | |
| Endive (Escarole), Lettuce, Spinach | phids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | | |
| Peppers, Tomatoes | Aphids, Colorado Potato Beetles, Fall Armyworms (Toma Beetles, Leafhoppers, Southern Armyworms (Tomatoes), Drosophila spp. (Tomatoes) | • | |
| Onions (Bulb and Green) | Onion Thrips | | |
| Radishes, Turnips, Carrots, Parsnips, Potatoes | Aphide, Flea Beetles, Leafhoppers, Dipterous Leafminers, Potato Beetles (Potatoes), Southern Armyworms (Potatoes | | |
| Sweet Corn | Aphids, Flea Beetles, Grasshoppers, Spider Mites | | |
| Melons, Squash (Summer, Winter) | Aphids, Thrips, Spider Mites | | |
| Vegetables: Preplant Soil Application Beans, Peas (except dried varieties) | Cutworms, Mole Crickets, Wireworms (Snap Beans) | | |
| Beats, Red (Table) | Cutworms, Mole Crickets | | |
| Cabbage | Cutworms, Root Maggots, Wireworms | | |
| Carrots, Sweet Corn | Cutworms, Mole Crickets, Wireworms | · · · · · · · · · · · · · · · · · · · | |
| Lettuce, Tomatoes | Cutworms, Wireworms | • • • | • • • • • |
| Radisbes, Turnips | Cutworms | | •••• |
| Fruits & Nuts Almonds (California only) | Scale Crawiers | · · · · · · · · · · · · · · · · · · · | •••• |
| Walnuts (California only) | Aphids, Scale Crawlers, Spider Mites and others* | • • • | |
| Apricots, Pesches, Nectarines | Aphids, Aprico: Mealybugs (Apricots), Spider Mites, Scal | le Crawlers | |
| Cherries | Black Cherry Aphids, Cherry Fruitflies, Cherry Rust Mite Crawlers and others* | es, Scale | |

e.

ļ

H

ľ

| Plums, Prunes | Aphids, Spider Mites | | |
|---|---|--|--|
| Caneberries (CA, OR, WA only): Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries | Aphids, Leafhoppers, Two-Spotted Spider Mites, Thrips, Raspberry Fruitworms | | |
| Strawberries | Aphids, Two-Spotted Spider Mites, Strawberry Leafrollers | | |
| Grapes | Aphids, Grape Berry Moths, Grape Leaffolders, Leafhoppers | | |
| Roses & Flowers Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Rhododendron, Roses | Aphids, Flea Beetles, Dipterous Leafminers, Leafhoppers, Mealybugs, Two-Spotted Spider Mites, Thrips, Whiteflies and others* | | |
| Ornamental Trees & Shrubs Arborvitae, Azaleas, Birch, Boxwood, Camellias, Douglas Fir, Elm, Hawthorn Holly, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine. Ornamental Plum, Poplar, Rhododendron, Spruce, Willow | Aphids, Mites, Scale Crawlers, Whiteflies, Tent Caterpillars and others* | | |

Storage and Disposal

Storage

Do not store near heat or open flame. Store in original container in an area inaccessible to children or pet... USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.

Disposal

Do not couse empty of, tainer. Thoroughly rinse container, securely wrap it in several layers of newspape, and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

Statement of Practical Treatment

If swallowed: Call a physician immediately. DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where irrigation (watering) is recommended in the directions or if waterfowl, i.e., duck or geese, can be expected in the treated area after treatment (except spot treatment), apply at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.