

Crop Life

DIPEL[®] 150 DUS

BACILLUS THURINGIENSIS

Net Weight 10 lbs.

ACTIVE INGREDIENT:

Bacillus thuringiensis, Berliner,
720 International Units of Potency
per mg. 0.15 billion International
Units per pound.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law
to use this product in a manner
inconsistent with its labeling.

STORAGE AND DISPOSAL DIRECTIONS

Storage: Reclose containers of
unused Dipel. Store in a dry place.
Disposal: Do not reuse container.
Perforate or crush and discard
empty containers according to
local trash disposal regulations.

SMITH COMPANY & WALLE
Waller, Texas - Pearsall, Texas
EPA Est. No. 3114-TX01
EPA Registration No. 31147-1

Dipel 150 Dust is a biological insecticide, intended
for use against chewing larvae (caterpillars) of
certain lepidopterous species.

Dipel 150 Dust is recommended for use by ground
or by air. Apply when pests are first noticed and
as necessary. Thorough plant coverage is necessary
for control.

APPLICATION RATES:

For the control of cabbage looper on Broccoli, C
Cauliflower, Collards, Kale, Mustard & Turnip Gr
Celery, Lettuce, Melons, Potatoes, Spinach and T
apply 25 to 50 lbs. per acre.

For the control of Imported Cabbage Worms on Bro
Cabbage, Cauliflower, Collards, Kale, Mustard &
Greens - apply 15 to 25 lbs. per acre.

For the control of Grape Leaf Folder on Grapes -
25 to 50 lbs. per acre.

SMITH & WALLE makes no warranty express or
implied, including the warranties of merchantability and/
for any particular purpose, concerning this material,
except those which are contained on this label.

Reg. Trad. Mark - Abbott Laboratories
North Chicago, Ill.

U.S.
Pat.
Off.
Reg.
EPA

Libe

DIPEL[®] 150 DUST

FACILEMENT DISPENSIBLE

50 lbs.

INGREDIENT:
Bacillus thuringiensis, Berliner,
(National Units of Potency
0.15 billion International
Units per pound).

DiPel 150 Dust is a biological insecticide intended
for use against chewing larvae (caterpillars) of
certain lepidopterous species.

DiPel 150 Dust is recommended for use by ground or
by air. Apply when pests are first noticed and repeat
as necessary. Thorough plant coverage is necessary
for control.

APPLICATION RATES:

For the control of cabbage looper on Broccoli, Cabbage,
Cauliflower, Collards, Kale, Mustard & Turnip greens,
Celery, Lettuce, Melons, Potatoes, Spinach and Tomatoes -
apply 25 to 50 lbs. per acre.

For the control of Imported Cabbage Worms on Broccoli,
Cabbage, Cauliflower, Collards, Kale, Mustard & Turnip
greens - apply 15 to 25 lbs. per acre.

For the control of Grape Leaf Folder on Grapes - apply
25 to 50 lbs. per acre.

SWISS ALP VALLE makes no warranty, express or implied,
including the warranties of merchantability and/or fitness
for any particular purpose, concerning this material,
except those which are contained on this label.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

DIRECTIONS FOR USE

CLASSIFICATION

Compliance with Federal law
requires this product in a manner
consistent with its labeling.

DISPOSAL DIRECTIONS

Reclose containers of
this material. Store in a dry place.
Do not reuse container.
Do not crush and discard
containers according to
local disposal regulations.

SWISS ALP VALLE
P.O. Box 100 - Pearsall, Texas
Phone No. 314-2701

Swiss Alp Valle Laboratories
North Chicago, Ill. 60064

