

DIRECTIONS

For maximum insect control, a combination of surface and space treatment is recommended. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

SURFACE OR CONTACT SPRAYING: To control roaches, ants, spiders, crickets, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites, and cadellies: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. Spray insects directly whenever possible. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat treatment as needed.

To control clover mites indoors, spray floors, baseboards, around windows and doors and other possible points of entry. To prevent reinfestation, spray the outside of the foundation and lower walls. Do not spray growing plants with this product. Repeat as needed.

To control silver fish, apply spray thoroughly to cracks and crevices throughout the entire building, including the attic. Treat inside of book cases, dresser drawers, cabinets and closets. Repeat treatment as needed.

To control fleas indoors, spray infested areas thoroughly with particular attention to cracks, crevices, behind baseboards and mouldings and under edges of rugs. Pet bedding should be thoroughly cleaned and sprayed or destroyed and replaced. Pets should be treated using an approved insecticide to prevent reinfestation. Repeat treatment as needed.

To control brown dog ticks indoors, apply spray carefully to all observable crevices, behind base boards, pictures, furniture, and draperies and under furniture and edges of rugs. Dogs should be treated with an approved insecticide to reduce reinfestation. Repeat treatment as needed.

SPACE TREATMENT AS A MIST, VAPOR OR FOG For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths: Close doors and windows and shut off ventilating systems. Apply with conventional thermal, mechanical or compressed air equipment adjusted to deliver a fine mist or fog. Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of one gallon to 50,000-100,000 cubic feet of space. Keep area closed for at least 10 to 15 minutes after treating.

ACCEPTED

NOV 21 1974

INSECTICIDE
FOR ECONOMIC PEST CONTROL ACT
REGISTERED UNDER NO. 111111
SUBJECT TO ATTACHED CONDITIONS.

PYRENONE® NO. 4 Insecticide Industrial and Mill Spray

For Rapid Control of Flying and Crawling Insects

Use in Food Processing Plants, Canneries, Packing Houses, Mills, Elevators, Granaries, Warehouses, Box Cars, Ships' Holds, Breweries, Bottling Plants, Bakeries, Stores, Markets, Theaters, Hospitals, Schools, Hotels, Motels and Restaurants.

The Insecticidal Components (Piperonyl Butoxide and Pyrethrins) are classified as food additives and When This Product is Used as Directed, Allowable Tolerance Levels Will Not Be Exceeded.

ACTIVE INGREDIENTS		100%
Pyrethrins	0.20%	
* Piperonyl Butoxide, Technical	1.00%	
Petroleum Distillate	98.80%	

* Equivalent to 0.80% (butylcarbityl) (6-propylpiperonyl) ether and to 0.20% of related compounds

CAUTION

KEEP OUT OF REACH OF CHILDREN

See rear panel for additional precautions

NET CONTENTS

Manufactured by
JOHNSON-HAYWORTH CHEMICAL CO.
LA GRANGE, ILLINOIS 60525

ACCEPTED

NOV 21 1974

INSECTICIDE
EFFECT
REGISTER-
SUBJECT

PYRENONE® NO. 4

Insecticide

Industrial and Mill Spray

For Rapid Control of Flying and Crawling Insects

Use in Food Processing Plants, Canneries, Packing Houses, Mills, Elevators, Granaries, Warehouses, Box Cars, Ships' Holds, Breweries, Bottling Plants, Bakeries, Stores, Markets, Theaters, Hospitals, Schools, Hotels, Motels and Restaurants.

The Insecticidal Components (Piperonyl Butoxide and Pyrethrins) are classified as food additives and When This Product is Used as Directed, Allowable Tolerance Levels Will Not Be Exceeded

ACTIVE INGREDIENTS		100%
Pyrethrins	0.20%	
* Piperonyl Butoxide, Technical	1.00%	
Petroleum Distillate	98.80%	

* Equivalent to 0.20% (butylcarbityl) (6-propylpiperonyl) ether and to 0.20% of related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See rear panel for additional precautions

NET CONTENTS

Manufactured by
JOHNSON-HAYWORTH CHEMICAL CO.
LA GRANGE, ILLINOIS 60525

Use Precautions

Foods should be removed or covered during treatment. All food surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

Remove any birds, pets or fish aquariums before spraying.

Vacate premises after treatment and ventilate before reoccupying.
Sweep up and remove dead insects.

Repeat treatment when reinfestation occurs.

CAUTION:

Harmful if swallowed. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse empty container. Return to drum conditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

EPA Reg. No. 31268-1

Establishment No. 10000

JOHNSON-HAYWORTH CHEMICAL CO.