

Net Contents 1 quart

## Cut-Heal Brand

Horse, Livestock and Barn Spray.  
For use against houseflies, horn-  
flies, stable flies, mosquitoes, and  
gnats on horses, beef cattle, dairy  
cattle and as a space spray in barns  
and stables.

**Active Ingredients:**

Pyrethrins .....	0.05%
	W/W
Technical Piperonyl Butoxide* .....	0.50%
Petroleum distillate .....	99.45%
	100.00%

\*Equivalent to 0.40% (butylcarbityl) (6-propylpiperonyl) ether and 0.10% related compounds.

### CAUTION:

KEEP OUT OF REACH OF CHILDREN.

(See additional cautionary statements on back panel)

### CAUTION:

DO NOT USE, POUR, SPILL OR STORE  
NEAR HEAT OR OPEN FLAME.

EPA REG. No. 31971-1

Manufactured by:

**Cut-Heal, Inc.**

206 Garvon, Garland, Texas 75040

ACCEPTED

JAN 17 1974

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 31971-1. SUBJECT  
TO ATTACHED COMMENTS.

### DIRECTIONS FOR USE

For control of houseflies, hornflies, stable flies, mosquitoes and gnats. Spray may be applied with hand sprayers or with mist machines adjusted to deliver a fine spray.

**Space Spray** — To attain quick knockdown and kill of flying insects in enclosed spaces, spray toward the ceiling and corners until the space is filled with a fine mist. For best results, close all windows and doors prior to spraying. Keep the room closed for 10 to 15 minutes after spraying. Do not remain in treated areas. Ventillate the area after treatment is completed. Repeat treatment as necessary.

**To Kill Flies on Horse, Beef cattle and dairy cattle** — Spray one or two fluid ounces as a fine mist on each animal or spray until a fine dew appears on the ends of the hair. Avoid wetting coat with any oil spray. Repeat at daily intervals, or as necessary for continued protection.

**Flying Insects in Barns and Stables:** Apply according to directions given above under **Space Spray**. Make sure that no open milk containers are exposed. Remove animals when applying space spray.

### CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Avoid inhalation. Do not wet the skin of animals or apply more than two fluid ounces of spray to an animal at one time. Do not use, pour or spill near an open flame. Keep out of lakes, streams and ponds.

**Food Areas:** Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation.

**Container Disposal** — Do not reuse container. Crush used container and bury in non-cropland soil away from water.