

U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460

EDV	Rea	Numbe
EFA	rcg.	Numbe

Date of Issuance:

32240-2

SEP 1 2 2008

NOTICE OF PESTICIDE:

Registration

XX Reregistration (under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product:

Crop Cure

Name and Address of Registrant (include ZIP Code):

Domain, Inc. 156 High Street

Attn: Henry Jacoby 6709 Ilex Court

New Richmond, WI 54017

New Market, MD 21774

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the thiram Reregistration Eligibility Document(s), EPA has reregistered the product listed above. Enclosed is a copy of your label stamped "Accepted with comments." This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Signature of Approving Official:

Tony Kish

Product Manager, Team 22

Fungicide Branch

Registration Division (7505P)

Date:

- SEP 1 2 2008

- 1. Submit one copy of your final printed labels with the changes herein, before releasing the product for shipment.
- 2. Make the following label changes:
- A. On the label's front panel, delete 'ALL' from the product description beneath the product name.
- B. In the 'Ingredient Statement', add "Total 100%" under 'Inert Ingredient'.
- C. Increase the font size of the signal word "WARNING," so that appears more prominent, and distinct from the surrounding text.
- D. Likewise, throughout the label increase the font size of the text to at least 8 point to ensure guidance is legible.
- E. In the section 'Directions for use on Hay', its final sentence requires 'hay' to be inserted, so that it reads, "Store hay in a well ventilated area."
- F. From the section, 'Directions for use on Silages' delete "or equipment," from the final sentence as it may lend for ambiguity.
- G. Instead of the headings "Below 50%, 50% -70%, Above 70%," used in the section, 'The Following Application Rates should be', append "moisture" to each, so that it reads: "Below 50% moisture, 50%-70% moisture, Above 70% moisture."
- H. The label's subsection 'Container Disposal' must be revised to comply with PR Notice 2007-4.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release of the product for shipment constitutes acceptance of these conditions.

A copy of the label stamped "Accepted with Comments" is enclosed for your records.

Enclosure

CROP CURE Original Formula

registered under EPA Reg. No. 3 2240-2



FOR HAY AND ALL SILAGES

KEEP OUT OF REACH OF CHILDREN

WARNING FIRST AID

if in ayes	Fire eye poon and unse slowly and gently with water for 15 20 minutes.
	 Hemove instantioness, if present, after the first 5 minutes, then continue applied eye
	- Cris a poison to the letter or do for the treatment advice
if swallowed	Can a coson (or tro), enter or do (tor in mediately for treatment advice.
	· riave a person sip a glass of water if abig to swallow.
	Do not indure variating an ess told to do so by a policon control centar or doubte.
	Do not give anything by mouth to an unconstitious person.
(f Inheled	Move detact to trash air.
	 If person is not breathing, sail \$15 or an embalance, then give artifold respiration, preferably mouth-to-mouth it possible.
	Cell a poison control center or doctor for treatment advice.
if on akin or clothing	- Take oil contaminated clottling
	Ricse skip immediately with pier ty of water for 15, 20 giroups
	Cart a polegic - prittor - enter or do , for tor treatment agrice,

HOT LIKE NUMBE

Have the product container or laber with you when calling a policer control center or contour going for realment. For emergency medical attention, lear 1, 800, 822-1925.

NOTE TO PHYSIC

Sociam one state is non-consistent the spirition. However, within primarie of water or takes all, converted into specification where it often specification of the specification

EPA EST. NO. 32240-WI-1

EPA REGISTRATION HO. 32240-2

Net Wt. 50 lbs.



DOMAIN, INC.

New Richmond, WI 54017

CROP CURE - A MOLD INHIBITING HAY PRESERVATIVE For the preservation of baled hay (Alfalfa, Brome Grass, Clover, Orchard Gress and Timothy) to be stored for the feeding season.

DIRECTIONS FOR USEIt is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE ON HAY:

Apply CROP CURE uniformly onto the hay during the bailing process with a Gandy applicator. Store in a wellventilated facility.

THE FOLLOWING APPLICATION RATE SHOULD BE

USED: 15-20% Molsture.......3-4 lbs per ton of forage 20-25% Moisture.4-5 lbs per ton of forege Baled hay should be stacked to allow proper ventilation. To ensure this square bales should be stored with the cut edge up and alternate layers of bales should be placed at right angles to one another. There must also be a small amount of space left between bales in the same layer. Remove all loose hay from the top of each layer of bales before starting the next laver.

USE PRECAUTIONS FOR HAY AND SILAGE

Treating higher moisture levels than recommended or at an application rate below the recommended amount may result in heating, crop discoloration, and spollage. Hay treated with CROP CURE is to be fed to livestock and poultry only.

DO NOT STORE UNTREATED HAY ON TREATED HAY WHILE IT IS CURING OUT.

CROP CURE AS AN AID IN SILAGE FERMENTATION

For the preservation of processed corn, hay, sorghum, oat and grass sliages stored in conventional upright, ox-ygen limiting, pit and bunker slios. And for high moisture ear or shell corn sliage ground to a 1/4" grind, packed well and stored in upright conventional or oxygen limit-

ORECTIONS FOR USE ON SILAGES
Apply CROP CURE at time of chopping or at the blower when silage is being loaded into the silo using a Gandy applicator or equipment or equivalent.

THE FOLLOWING APPLICATION RATES SHOULD BE USED.

Corn, hey, gress, oal, and sorghum stage	Not Recommended	80-70% 1 · 2 lbs/lon	Above 70% Not Recommended
High moisture ensited core	Below 25% Not Reconstrended	25-35% 3-6 lbs/ton	Above 85% Not Recommended

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eye wear (goggles, face shield, or safety glasses). Harmful if swallowed, absorbed through the skin or inhaled. Avoid contact with skin. Avoid breathing dust. Wash thor-oughly with soap and water often after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to lakes, streams or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning equipment or disposing of wash water.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Seal tightly. Do not store in direct sunlight.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or recon-

ditioning, or puncture and dispose in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

To the extent consistent with applicable laws, Domain, Inc. is not responsible for a loss caused by use inconsistent with directions on this label.

CROP CURE
For the selective inhibition of microbes in livestock and poultry feeds. Also, for the improvement of palatability in livestock feeds.

DIRECTIONS FOR USE IN GROUND FEEDS
MIX WELL
Grinders/Mixers: Add grains (feeds) (corn, soybean, wheat, barley, oats) to other ingredients (premix diet supplements, vitamins, minerals) and the required amount of CROP CURE. Then mix in high moisture ingredients (ilquid vitamins, molasses) inadequate mixing will produce inconsistent mold inhibition.

APPLICATION RATES FOR BEEF, SWINE & POULTRY FEEDS.

Moisture Content of Feed	0 -32 F	33'-55'F	56'-80'F
15%	1 lb/ton	1 lb/ton	2 lb/ton
15-20%	2 lb/ton	3 lb/ton	4 lb/ton
20-25%	3 lb/ton	4 lb/ton	5 lb/ton
25-30%	4 lb/ton	5 lb/ton	6 lb/ton

based on a 10-14 day storage period.

RECTIONS FOR USE IN DAIRY RATIONS

Mbxed Feeds: Add grains (corn, soybean, wheat, bar-ley, oats) with other ingredients (premix diet supplements, vitamins, minerals) and required CROP CURE, and mix well. Then mix in high moisture ingredients (liquid vitamins, molasses). CROP CURE may be used in the grain mix at the rate of 3-4 lb/ton of feed. The objective is to achieve 1/2 oz intake per head per day. Higher rates may be needed should feed be high in moisture.



"The CROP CURE composition and uses thereof are covered by United States patents and patent applications under which Domain, Inc. is exclusively licensed."