

FRONT PANEL

1463

PM 21

32342-4

# SUL - CIDE F - 9 flowable sulfur

**ACCEPTED  
REREGISTERED**  
16281986  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 32342-4

**ACTIVE INGREDIENT**

Sulfur as S ..... 70.0%  
**INERT INGREDIENTS** ..... 30.0%

**TOTAL** 100.0%

(Contains 9.0 pounds Sulfur per gallon)

**NET CONTENTS 5 GALLONS**

**DIRECTIONS FOR MIXING BEFORE USING  
STIR UNTIL SMOOTH**

Use recommended amount in partially filled spray tank. Keep agitator running during filling and spraying operation. **DO NOT ALLOW MIXTURE TO STAND.** Failure to maintain agitation will cause F-9 Flowable Sulfur to settle and may necessitate manual stirring to redispense.

The strong adhesive properties of F-9 Flowable Sulfur act as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulfur in any form is corrosive material. TO REDUCE THE EFFECT, EQUIPMENT SHOULD BE FLUSHED DAILY.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See side panel for Additional  
Precautionary Statements



**CHEMICAL  
ENTERPRISES, INC.**  
8582 Katy Freeway  
Suite 202  
Houston, Texas 77024  
Phone: 713/461-0606

**The Sulfur Company**

EPA Reg No 32342-4

EPA Est No 32342-TX 1

203

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing spray mist. Avoid contact with eyes, skin, and clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling and before eating or smoking. Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Promptly drink a large quantity of milk, egg whites, or gelatin solutions. Get medical attention immediately.  
**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.  
**IF IN EYES:** Flush with plenty of water. Get medical attention.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not apply when weather conditions favor drift from target areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL HAZARDS**

Keep away from heat, sparks, or flame. Do not smoke while applying this product.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**RE-ENTRY STATEMENT**

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written or oral warnings must include the following information:

**CAUTION**

(Insert area or field description) treated with SUL-CIDE W-90 wettable sulfur on (insert date of application). Do not enter treated areas for 24 hours unless appropriate protective clothing is worn. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

**DIRECTIONS FOR MIXING BEFORE USING: STIR UNTIL SMOOTH.**

LEFT PANEL

Pour recommended amount in partially filled spray tank. Keep agitator running during filling and spraying operation. DO NOT ALLOW MIXTURE TO STAND. Failure to maintain agitation will cause F-9 Flowable Sulfur to settle and may necessitate manual stirring to redisperse. The strong adhesive properties of F-9 Flowable Sulfur act as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulfur in any form is a corrosive material. TO REDUCE THE EFFECT, EQUIPMENT SHOULD BE FLUSHED DAILY, taking care to observe pesticide disposal procedures below.

**APPLICATION INSTRUCTIONS**

Fruit tree dosage rates are given as pints of F-9 Flowable Sulfur per 100 gallons of water. Wet all foliage.

- APPLE, PEAR** — Scab  
 Pre-bloom through Calyx sprays 1 to 2½ pints  
 Cover sprays ½ to 1½ pints  
 Powdery Mildew  
 Pre-bloom through Calyx sprays 1½ to 2½ pints  
 Cover sprays 1½ pints
- CHERRY, PLUM, PRUNE** — Brown Rot, Leafspot  
 Pink and bloom sprays 1½ to 2½ pints  
 Petal fall shuck and cover sprays ½ to 1½ pints
- CITRUS** — Rust Mite, Clover Mite  
 Apply sprays November thru May
- THRIPS** (Early spring treatment for the reduction of Thrips population)  
 Apply 1 to 2 gallons per acre in sufficient water for thorough coverage. Make application after spring flush has made 3 to 4 inch growth. Repeat if necessary. Do not apply within 21 days of an oil application.
- GRAPES** — Powdery Mildew  
 Use ½ to 1½ pints per 100 gallons water, maximum of 6 pints F-9 Flowable Sulfur per acre in spring and summer. Make first application when shoots are 5 to 8 inches long, second when shoots are 12 to 15 inches long and third at early bloom stage. The timing, rate and number of applications will vary with local conditions. Consult State Agricultural Service or State Agricultural Experiment Stations in your region for specific information. Follow the USE PRECAUTIONS given on this label.
- PEACH, NECTARINE** — Powdery Mildew, Brown Rot, Leafspot  
 Pink and bloom sprays 1 to 3 pints  
 Petal fall, shuck and cover sprays ½ to 1½ pints
- POTATOES** — Powdery Mildew  
 Use 2 to 3 pints in 8 gallons water per acre by air. Apply at first sign of infection and repeat at 3 week intervals.
- STRAWBERRIES** — Powdery Mildew  
 Use ¾ pint in 100 gallons of water per acre. Apply at first sign of infection and repeat at 3 week intervals. For strawberries that will be canned, discontinue application of sulfur well before harvest in accordance with local canner's recommendations.
- TOMATOES** — Tomato Russet Mite  
 Use ½ to ¾ gallon per acre, depending upon size of plants, in enough water for complete wetting of all foliage.
- COTTON** — Atlantic Mite  
 2 gallons for early, mid and late season control. Being sure to wet entire plant.
- PEANUTS** — Rust  
 Apply 1½ to 3 pints per acre in sufficient water for thorough coverage. Begin application approximately 6 weeks after planting. Repeat at 7 to 10 day intervals as long as rust control is required. Do not apply within 14 days of an oil application.
- SUGAR BEETS** — Powdery Mildew  
 For control of powdery mildew apply the recommended rate in 10 gallons of water per acre by air at first sign of infection. Use 1½ to 3 pints per acre in the Pacific Northwest and 1½ to 6 pints per acre in Arizona, Colorado, Nebraska, Wyoming, New Mexico, Texas and Utah. Repeat applications should be made at 10 to 30 day intervals or as necessary throughout the season. Under most disease conditions only one or two applications will be needed.

**STORAGE AND DISPOSAL**

**STORAGE:** Store in a dry location away from children, animals, foods, feeds, seeds, or other agricultural chemicals. This product freezes at 32°F but usually upon thawing is re-useable. In the event of spillage or leakage, soak up material with absorbent clay, sand, sawdust, or other absorbent material. Scrape up and dispose of in accordance with information given under "DISPOSAL." Repackage and relabel remaining useable product in a sound container. In case of fire or other emergency, report at once by toll-free telephone to 800-424-9300.

**DISPOSAL:**

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.