



ALLIED SERVICES

AL-TROL 0.6-TBA

(Contains 0.05 lb. trichlorobenzoic acid per gallon)

See PRODUCT BULLETIN For Directions

ACTIVE INGREDIENTS:

Dimethylamine salt of 2, 3, 6-trichlorobenzoic acid	0.36%
Dimethylamine salt of other chlorinated benzoic acids	0.24%

INERT INGREDIENTS: 99.40%

EPA Reg. No. 32427-12-AA

CAUTION — KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May irritate eyes, nose, throat, and skin. Avoid breathing spray mist. Avoid contact with skin, eyes, and clothing.

In case of contact — Flush with plenty of water; for eyes, get medical attention.

Injury to or loss of desirable trees or other plants may result from failure to observe the following: Do not apply or clean or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Do not use area actually treated for crop production.

Spray drift, and possibly vapors from treated areas, may injure susceptible crops such as beans of all types, tomatoes, pears, cotton, tobacco, various ornamental plants, orchard and vine crops. Do not apply to areas where nearby fields are to be used for growing susceptible crops during the same season.

Do not use equipment with which trichlorobenzoic acid has been applied for spraying insecticides, fungicides, selective herbicides, and fertilizers as plant injury may result.

Thoroughly clean all traces of this product from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Do not contaminate water by cleaning of equipment or disposal of wastes.

Do Not contaminate water used for irrigation or domestic purposes. Keep out of lakes, streams or ponds.

DO NOT STORE NEAR FEEDS OR FOODSTUFFS.

Do Not allow contact with fertilizers, insecticides, fungicides or seeds.

Buyer assumes all risks of use, storage or handling this material not in strict accordance with directions given herewith.

Net Contents

Gallons

ALLIED SERVICES

Division of Allied Equities Corporation

765 Landess Avenue

Milpitas, Calif. 95035

(408) 263-2020



ALLIED SERVICES
Division of Allied Equities Corporation

765 Landess Ave.
Milpitas, CA 95035
(408) 263-2020

1411 N. Paris Road
Columbia, MO 65201
(314) 443-1461

87 E. Olive St.
Suite 2D
Fresno, CA 93728
(209) 237-4318

PRODUCT BULLETIN

AL-TROL 0.6 – TBA

EPA Reg. No. 32427-12

Apply, store, use, handle only as specified on product labeling. Buyer/User may be subject to criminal and/or civil penalties for failure to follow labeling directions and precautions.

DIRECTIONS FOR USE

For use on non-crop areas (utility, highway, pipeline, railroad right-of-ways, lumber yards, tank farms, storage areas, industrial areas) for control of most annual and perennial grasses and broadleaved weeds. Both foliage and root absorption contribute to plant kill. Effects on some species are slow to appear and may not become apparent until the herbicide has been carried into the root zone of the plants by moisture. The degree of control and duration of effect will vary with the amount of herbicide applied, soil type, rainfall, and other conditions; i.e., adsorptive soils (those of heavy texture or high in organic matter) require the higher dosage levels.

HOW TO APPLY: Application may be made with a boom sprayer, hand spray gun, or for small problem areas a sprinkling can. Before applying, calibrate technique to deliver correct dosage necessary to comply with the rates listed below. Thoroughly spray foliage and stems, as well as ground under plants. Apply as a coarse spray to reduce drift.

WHEN TO APPLY: Best results will be obtained if application is made after active growth begins – but before growth has progressed sufficiently to retain more than half the spray. Earlier and later applications also are effective; however, rates near the upper limit of the recommended dosage range may be required. Apply during periods of moderate rainfall. *Do not apply when ground is frozen or during other weather conditions which would favor excessive run-off.*

USE RATES:

PERENNIAL WEEDS: For field bindweed, leafy spurge, Canada thistle, Russian knapweed, bur ragweed, blueweed and climbing milkweed, apply 200 to 400 gallons per acre; for dandelion, plantain, dock, dog fennel, ragweed and wild lettuce, apply 400 to 600 gallons per acre.

WOODY VINES: For trumpet vine, honeysuckle, kudzu and smilax, apply 200 to 400 gallons per acre.

WOODY BRUSH: For conifers (spruce, balsam fir, cedar and pine), wild roses (Macartney or Cherokee), sumac, persimmon, sassafras, black locust and hackberry, apply 80 gallons per acre.

REFER TO THE PRODUCT LABEL FOR ADDITIONAL DIRECTIONS AND PRECAUTIONS.

– OBSERVE THESE CAREFULLY –