



**ALLIED SERVICES**

765 Landess Ave.  
Milpitas, CA 95035  
(408) 263 2020

1411 N. Paris Road  
Columbia, MO 65201  
(314) 443 1461

87 E. Olive St.  
Suite 2D  
Fresno, CA 93728  
(203) 237 4318

# PRODUCT BULLETIN

AL-TROL 0.6 - TPA

EPA Reg. No. 32427-14

*Apply, store, use, handle only as specified on product labeling. Buyer/User may be subject to criminal and/or civil penalties for failure to follow labeling directions and precautions.*

## DIRECTIONS FOR USE

For use on non-crop areas (utility, highway, pipeline, railroad right-of-ways, lumber yards, tank farms, storage areas, industrial areas) for control of selected grasses and broadleaved weeds including those growing in perennial grasses. Both foliage and root absorption contribute to plant kill. Effects on some species are slow to appear and may not become apparent until the herbicide has been carried into the root zone of the plants by moisture. The degree of control and duration of effect will vary with the amount of herbicide applied, soil type, rainfall, and other conditions; i.e., adsorptive soils (those of heavy texture or high in organic matter) require the higher dosage levels.

**HOW TO APPLY:** Application may be made with a boom sprayer, hand spray gun, or for small problem areas a sprinkling can. Before applying, calibrate technique to deliver correct dosage necessary to comply with the rates listed below. Thoroughly spray foliage and stems, as well as ground under plants. Apply as a coarse spray to reduce drift.

**WHEN TO APPLY:** Best result will be obtained if applied just before or during the period when active growth begins and moderate rainfall can be expected. Removal of dense weed growth prior to treatment will improve effectiveness. *Do not apply when ground is frozen or during other weather conditions which would favor excessive run-off.*

## USE RATES:

**ANNUAL WEEDS AND GRASSES:** For foxtail, crabgrass, barnyard grass, smartweed, lambquarters, pigweed, ragweed, puncturevine, Russian thistle, plantain, Koch and turkey mullein apply 300 gallons per acre.

**PERENNIAL WEEDS:** For bur ragweed, Indian rush pea, Texas blueweed, mouse ear prostrate weed, alkali sida, chickweed, plantain, dandelion, bouncingbet apply 300 gallons per acre. For foxtail and wild oat, leafy spurge, Russian knapweed, Canadian thistle apply 350 to 400 gallons per acre.

REFER TO THE PRODUCT LABEL FOR ADDITIONAL DIRECTIONS AND PRECAUTIONS

- OBSERVE THESE CAREFULLY

Feeds  
5,000 sq. ft.  
(50' x 100')

# HOWARD JOHNSON'S

EPA Reg. No. 32802-1

**ACTIVE INGREDIENTS**


Isomyl ester of Silvex 2, 4, 5-trichlorophenoxy Propionic Acid 71  
Isomyl ester of 2, 4-Dichlorophenoxyacetic Acid 25

**INERT INGREDIENTS** 98.51  
100.00

Equivalent to 0.5% Silvex 2, 4, 5-trichlorophenoxy Propionic Acid  
Equivalent to 0.5% 2, 4-Dichlorophenoxyacetic Acid

## weed & feed

NET WT. 18 LBS.



**CAUTION**  
KEEP OUT OF THE REACH  
OF CHILDREN  
SEE THE HOWARD JOHNSON'S BACK LABEL