

## FIRST AID

**EXTERNAL:** Flood with water for 15 minutes.

**INTERNAL:** Drink large amounts of milk. Follow with Milk of Magnesia, vegetable oil, or beaten eggs.

**CALL PHYSICIAN IMMEDIATELY.**

**EYES:** Wash with water for 15 minutes and get prompt medical attention.

## WARNING

Harmful or fatal if swallowed. May produce severe chemical burns. Do not allow contact with eyes, skin, food, mucous membranes, or clothing.

## WARNING

Dispose of spills by flushing thoroughly. Do not mix Step 1 Superchlorinator with any other chemicals or contaminate in any way. Do not reuse empty container. Rinse empty container thoroughly with water and discard.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.



# SUPERCHLORINATOR FOR SWIMMING POOLS

DESTROYS VISIBLE ALGAE

CONCENTRATED SOLUBLE  
ORGANIC CHLORINE

Active Ingredient:

1.3 dichloro-5.5-dimethyl-  
hydantoin 100%

Available  
Free Chlorine 66% minimum

NET WEIGHT: 16 ounces (1 pound)

**KEEP OUT OF REACH OF CHILDREN.  
Strong oxidizer: Contact with heat or  
with other materials may cause fire.  
Store in a cool, clean, dry place.**

Sold by  
HYDROLOGY LABORATORIES, INC.  
SMITHTOWN, N.Y. 11787

EPA Reg. No. 32460-5

## DIRECTIONS

Consult dosage chart to determine the correct amount of Step 1 Superchlorinator to add to the pool water. **Slowly** add the correct dosage of Step 1 Superchlorinator **to the skimmer basket**, with the pump and filter operating during the addition.

POOL SIZE (PACKAGES OF STEP 1  
IN GALLONS) | SUPERCHLORINATOR

1,000-10,999	1
11,000-20,999	2
21,000-30,999	3

Larger Pools: Use one package (1 lb) of Step 1 Superchlorinator for each 10,000 gallons of pool water.

Keep the filter operating and do not resume swimming until the chlorine level drops below 5 P.P.M. Repeat procedure if visible algae persists and chlorine level has dropped below 5 P.P.M. Step 1 Superchlorinator should be used at the first signs of visible algae or anytime the chlorine level falls to zero due to high bather load, high temperatures, or contamination.

## NOTICE

Seller guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling, and accepts the product on these conditions.

