

RL 2 RABON® LIVESTOCK DUST

A READY-TO-USE INSECTICIDE DUST

To Control HORN FLIES and LICE and aid in the control of FACE FLIES on BEEF CATTLE, DAIRY CATTLE and to control LICE on SWINE.

ACTIVE INGREDIENTS BY WEIGHT

Permethrin 2.4%
phosphate 8.00%
INERT INGREDIENTS 89.60%

EPA Reg. No. 32695-1
EPA Est. No. 10543-MO-1

CAUTION:
KEEP OUT OF REACH OF CHILDREN.
See other cautions on side panel.
TAKE TIME — OBSERVE LABEL DIRECTIONS

NET WEIGHT 12½ POUNDS

GENERAL DIRECTIONS

RABON® Livestock Dust is a ready-to-use insecticide dust which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

USE DIRECTIONS

ANIMAL	INSECT	REMARKS
Beef Cattle Dairy Cattle	Horn Flies Face Flies Lice	HAND DUSTING —Apply approximately 2 oz. of dust by shake can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary. SELF-TREATING DUST BAG FORCED USE —Put RABON® Livestock Dust in cotton cloth or double burlap bags or use pre-packed weather-proof cattle dust bags and hang in barn door exits or alleyways leading from animal buildings, stalls or mineral blocks, or watering holes. Protect cloth or burlap bags from weather. FREE CHOICE USE —Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice aids in the control of lice.
Swine	Lice	HAND DUSTING —Apply 3-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% dust should be applied per 150 sq. ft. of bedding.

PRECAUTIONS IN USING

CAUTION: Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs.

Keep away from heat and open flames. Do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty containers. Destroy it by perforating or crushing. Bury or discard in a safe place.

NOTICE OF WARRANTY

DALE ALLEY CO. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

U. S. Patent No. 3777716

DEC 2 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 32695-1

RL 2 RABON LIVESTOCK DUST

GENERAL DIRECTIONS

USE DIRECTIONS

REMARKS

A READY-TO-USE INSECTICIDE DUST

FOR THE CONTROL OF LICE AND MITES ON CATTLE, HORSES, AND SWINE.
- FARMER - - - - - SEE CAUTION DIRECTION AND TO
control LICE on SWINE

ACTIVE INGREDIENTS	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl vinyl dimethyl phosphate	3.00%
INERT INGREDIENTS	97.00%
EPA Reg. No. 32695-1	TOTAL
EPA Est. No. 10543-MQ-1	100.00%

CAUTION:
KEEP OUT OF REACH OF CHILDREN.
See other cautions on side panel.
TAKE TIME — OBSERVE LABEL DIRECTIONS

NET WEIGHT 12½ POUNDS

MANUFACTURED BY DALE ALLEY CO. ST. JOSEPH, MISSOURI 64502

ACCEPTED

NOTE: FOR FARMER ONLY