DIRECTIONS FOR USE

IMPORTANT: Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using CO-RAL 14 Bulk Dust.

conditions of SALE: These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. Dale-Alley Co. and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

CO-RAL 1% Bulk Dust may be used to fill cattle dust bags or it may be applied directly to cattle, swine, or swine bedding by use of a shaker can or a suitable mechanical dust applicator. When used as recommended, CO-RAL 1% Dust controls horn flies and lice on beef and dairy cattle and lice on swine. CO-RAL 1% Dust will also reduce face fly populations on cattle, however, as face flies are able to travel over an area of several miles, best reduction of face fly populations is achieved when CO-RAL 1% Dust is used on an area-wide basis.

RECOMMENDED APPLICATIONS

ANIMAL	PARASITES	REMARKS		
Beef and Dairy Cattle	Horn Flies Lice Face Flies	DUST BAG: Place contents of this package in any commercially available dust bag. Suspend bags in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed, or minerals. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows, bags may be suspended in the exit through which cows leave the milking barn. In all cases, the bags should be adjusted so that the bottom of the bag will hang four to six inches below the topline of the cattle. For reduction of face flies, the bags must be located so the animals will be forced to use them daily and hung at a height to insure that the faces of the cattle will be dusted.		
	Horn Flies Lice	DIRECT APPLICATION: CO-RAL 1% Dust may be applied by use of a shaker can or by using a suitable mechanical dust applicator. Animals should be thoroughly dusted using not more than 2 oz. *(6 level tablespoonfuls) of dust per animal per treatment Repeat as necessary.		
		DIRECT APPLICATION: Apply not more than 1 oz. *(3 level tablespoonfuls) per head as a uniform coat to the head, shoulders, and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary but not more often than once every ten days.		
Swine Lice uniformly over each 30 square use of a shaker can or a suita		BEDDING TREATMENT: Apply 2 oz. *(6 level tablespoonfuls) uniformly over each 30 square feet of fresh, dry bedding by use of a shaker can or a suitable mechanical dust applicator. Repeat as necessary but not more often than once every tendays.		



Brand of coumaphos ANIMAL INSECTICIDE

1% BULK DUST

CONTROL OF HORN FLIES & LICE ON BEEF & DAIRY CATTLE — REDUCTION OF FACE FLIES ON BEEF AND DAIRY CATTLE — CONTROL OF LICE ON SWINE

USE IN DUST BAG, SHAKER CAN, OR MECHANICAL DUST APPLICATOR

BATCH NO.

CAUTION
KEEP OUT OF REACH OF
CHILDREN

(See side panel for warning s -ments)

ACTIVE INGREDIENTS

() () Diethyl () .3 chlora 4 methyl 2
oxo 2// 1 benzapyran 7 yl;
phosphorathicate 1.0%

INERT INGREDIENTS 99 0%
100 0%

EPA Reg No. 32695-2 EPA I t No. 10543-MO-01


USE ONLY AS DIRECTED



MANUFACTURED BY

DALE - ALLEY CO.

St Joseph, Missouri 64502



Fungicide, and Rodentiside Act,

RECOMMENDED APPLICATIONS (COR

ANIMAL	PARASITES	REMARKS	
Swine	Lice	DIRECT APPLICATION and BEDDING infestations both individual animals treated as directed above.	

No interval is required between treatment and slaughter or between milk as food.

*NOTE 1 oz. of this material equals approximately three level tail

RESTRICTIONS

For external, insecticidal use on above specified animals only.

Avoid contamination of feed, troughs, water and water utensils.

Provide thorough ventilation while dusting. Persons treating animal direct contact with skin or inhalation of dust.

CAUTION

Harmful if inhaled or absorbed through the skin. Do not get in breathe dust. Wash thoroughly with soap and warm water after ha

Do not contaminate feed or food. Keep out of reach of children.

If poisoning occurs, obtain prompt medical aid. Prolonged exposur cholinesterase depression.

To Physician — Atropine sulfate is antidotal, 2-PAM is also antido administered in conjunction with atropine.

This product is toxic to fish, birds and other wildlife. Keep out of Do not apply where runoff is likely to occur. Do not apply when drift from areas treated. Do not contaminate water by cleaning of wastes. Apply this product only as specified in this label.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy it by burying with waste or Stay away from smoke or fumes.

WARRANTY AND LIMITATION OF DAM

Dale-Alley Co. warrants that this material conforms to the chemical and is reasonably fit for the purposes referred to in the Directions of Sale subject to the risks referred to therein DALE-ALLEY CO. MA. OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OF FITNESS OR OF MERCHANTABILITY, and no agent of Dale to do so except in writing, with a specific reference to this warranteem a breach of this warranty shall be limited to direct damage consequential commercial damages, such as loss of profits or value

CO-RAL is a Reg. TM of the Parent Company of Farben Fabriken.

including the Warranty

Oust.

ion of experts based on dues on or in food and vide range of conditions tions are followed, it is normal conditions beyond. Buyer uses, this product

ay be applied directly to e mechanical dust applirn flies and lice on beef fuce face fly populations eral miles, best reduction in an area-wide basis.

IS

age in any commercially eas frequented by cattle the animals pass daily by also be suspended in admineral feeders. For suspended in the exit barn. In all cases, the bottom of the bag will pline of the cattle. For se located so the animals displayed hung at a height to ll be dusted.

Dust may be applied a suitable mechanical horoughly dusted using espoonfuls) of dust persary

ore than 1 oz *(3 level to the head, shouldtor suitable mechanical out not more often than

(6 level tablespoonfuls) of fresh, dry bedding by chanical dust applicator. Then than once every ten



Brand of coumaphos ANIMAL INSECTICIDE 1% BULK DUST

CONTROL OF HORN FLIES & LICE ON BEEF & DAIRY CATTLE — REDUCTION OF FACE FLIES ON BEEF AND DAIRY CATTLE — CONTROL OF LICE ON SWINE

USE IN DUST BAG, SHAKER CAN, OR MECHANICAL DUST APPLICATOR

NE	T WEIG	SHT	5 POUNDS
	8/	ATCH I	10.
			1
	CAI	JTI	ON
(EEP	OUT	OF	REACH

KEEP OUT OF REACH OF CHILDREN

(See side panel for warning statements)

> EPA Reg. No. 32695-2 EPA Est. No. 10543-MO-01

USE ONLY AS DIRECTED



MANUFACTURED BY

DALE - ALLEY CO.

St Joseph, Missouri 64502

AUG 1 1975
Under the Federal Insecticide.

Fungicide, and Rodenticide Act, as amended, for the pesticide

RECOMMENDED APPLICATIONS (Cont.)

ANIMAL	PARASITES	REMARKS		
Swine Lice infestations both ind		DIRECT APPLICATION and SeDDING TREATMENT: In severe infestations both individual animals and the bedding may be treated as directed above.		

No interval is required between treatment and slaughter or between treatment and use of milk as food.

*NOTE 1 oz. of this material equals approximately three level tablespoonfuls.

RESTRICTIONS

For external, insecticidal use on above specified animals only.

Avoid contamination of feed, troughs, water and water utensils.

Provide thorough ventilation while dusting. Persons treating animals should avoid repeated direct contact with skin or inhalation of dust.

CAUTION

Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate feed or food. Keep out of reach of children.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

To Physician — Atropine sulfate is antidotal, 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy it by burying with waste or burning Stay away from smoke or fumes.

WARRANTY AND LIMITATION OF DAMAGES

Dale-Alley Co. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the risks referred to therein DALE-ALLEY CO. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, and no agent of Dale-Alley Co. is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty. Fall be limited to direct damages, and that in timelude consequential commercial damages, such as 1 so of profits or values, etc.

CO-RAL is a Reg. IM of the Parent Company of Farben Fabriken Bayer Gold Leverkusen.