

SEW LINE

SEW LINE



Phly-N-Worm 44

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle, when used as directed.

Aids in the control of stomachworms (*Haemonchus*, *Ostertagia*, and *Trichostrongylus* spp.); nodular worms (*Oesophagostomum* spp.); large-mouth bowel-worms (*Chabertia* spp.); and hookworms (*Bunostomum* spp.) in cattle when fed in accordance with the directions for use.

FIVE INGREDIENT

Phenothiazine.....44%
INERT*INGREDIENTS.....56%

TOTAL 100.00%

*Refers only to ingredients which are not pesticides.

Each lb. DALE ALLEY Phly-N-Worm 44 contains 200 g. Phenothiazine.

FEED INGREDIENTS

Processed Grain By-Products, Calcium Carbonate and Mineral Oil

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

HAZARDS TO DOMESTIC ANIMALS

Do not feed to lactating dairy animals. Occasional sensitivity to phenothiazine has been reported in cattle. Do not administer to animals showing symptoms of acute respiratory conditions.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

See Reverse Side For Directions For Use and Storage and Disposal



NET WEIGHT 10 LBS.

Manufactured by

DALE ALLEY CO.

St. Joseph, Missouri 64502

EPA Reg. No. 32695-10

EPA Est. No. 10643-MO-01

LOT NO.

15

DIRECTIONS FOR USE

It is in violation of federal law to use this product in a manner inconsistent with its labeling.

Mix 1 pound of ALLEY Phly-N-Worm 44 in the daily feed, minerals or salt** of 100 adult cattle to provide the 2 grams Phenothiazine intended.

**Self-feed continuously as the sole source of salt.

Younger cattle: Follow the feeding chart below to provide 1/4 gram Phenothiazine per 100 pounds body weight per head daily.

AVERAGE BODY WEIGHT OF YOUNGER CATTLE	Lbs. ALLEY Phly-N-Worm 44		
	per 100	1000	2000 feedings
200 lbs.	1/4	2 1/4	5
300 lbs.	3/8	3 3/8	7 1/2
400 lbs.	1/2	5	10
500 lbs.	5/8	6 1/4	12 1/2
600 lbs.	3/4	7 1/2	15
700 lbs.	7/8	8 3/4	17 1/2

Continuous year round low-level phenothiazine treatment, in conjunction with good herd management practices, will control the specified parasites and will help prevent reinfection.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

- Cattle should be given access to feed, minerals or salt containing ALLEY Phly-N-Worm 44 for several weeks prior to the appearance of adult horn flies in the spring, and should be fed a ALLEY Phly-N-Worm 44 supplement continuously until flies disappear in the fall. This helps prevent the breeding of horn and face flies but has no effect upon existing adult fly populations. Since horn flies over-winter in pupal state, all cattle should be sprayed when the flies first appear. Spraying kills the flies emerging from last year's pupae and will protect cattle for about two weeks until the phenothiazine can take effect.

- Since face flies migrate from farm to farm it is essential that the flyicide program be carried out on an area-wide basis to assist in the reduction of the adult face fly population.

STORAGE AND DISPOSAL

Prohibitions:

Do not contaminate water or food by storage and disposal.

Pesticide Disposal:

Feed supplements containing Phly-N-Worm 44 which cannot be used should be buried in a safe place away from water supplies.

Container Disposal:

Do not reuse empty container. Wrap container and put in trash collection.