

SEW LINE

SEW LINE



Phly-N-Worm 11

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle, when used as directed.

Aids in the control of stomachworms (*Haemonchus*, *Ostertagia*, and *Trichostrongylus* spp.); nodular worms (*Oesophagostomum* spp.); large-mouth bowel-worms (*Chabertia* spp.); and hookworms (*Bunostomum* spp.) in cattle when fed in accordance with the directions for use.

ACTIVE INGREDIENT

Phenothiazine 11%

INERT*INGREDIENTS 89%

TOTAL

100.00%

*Refers only to ingredients which are not pesticides.

Each lb. DALE ALLEY Phly-N-Worm 11 contains 50 g. Phenothiazine.

FEED INGREDIENTS

Processed Grain By-Products, Calcium Carbonate and Mineral Oil

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

HAZARDS TO DOMESTIC ANIMALS

Do not feed to lactating dairy animals. Occasional sensitivity to phenothiazine has been reported in cattle. Do not administer to animals showing symptoms of acute respiratory conditions.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

See Reverse Side For Directions For Use and Storage and Disposal



NET WEIGHT 10 LBS.

Manufactured by

DALE ALLEY CO.

St. Joseph, Missouri 64502

EPA Reg. No. 32695-11

EPA Est. No. 10543-MO-01

LOT NO.

DIRECTIONS FOR USE

It is in violation of federal law to use this product in a manner inconsistent with its labeling.

Mix 1 pound of **ALLEY Phly-N-Worm 11** in the daily feed, minerals or salt** of 25 adult cattle to provide the 2 grams of Phenothiazine intended.

**Self-feed continuously as the sole source of salt.

Younger cattle: Follow the feeding chart below to provide $\frac{1}{2}$ gram Phenothiazine per 100 pounds body weight per head daily.

AVERAGE BODY WEIGHT OF YOUNGER CATTLE	Lbs. ALLEY Phly-N-Worm 11		
	per 100	1000	2000 feedings
200 lbs.	1	10	20
300 lbs.	1½	15	30
400 lbs.	2	20	40
500 lbs.	2½	25	50
600 lbs.	3	30	60
700 lbs.	3½	35	70

Continuous year round low-level phenothiazine treatment, in conjunction with good herd management practices, will control the specified parasites and will help prevent reinfection.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Cattle should be given access to feed, minerals or salt containing **ALLEY Phly-N-Worm 11** for several weeks prior to the appearance of adult horn flies in the spring, and should be fed a **ALLEY Phly-N-Worm 11** supplement continuously until flies disappear in the fall. This helps prevent the breeding of horn fly and face flies but has no effect upon existing adult fly populations. Since horn flies over-winter in pupal state, all cattle should be sprayed when the flies first appear. Spraying kills the flies emerging from last year's pupae and will protect cattle for about two weeks until the phenothiazine can take effect.

Since face flies migrate from farm to farm it is essential that the larvicide program be carried out on an area-wide basis to assist in the reduction of the adult face fly population.

STORAGE AND DISPOSAL

Prohibitions:

Do not contaminate water or food by storage and disposal.

Pesticide Disposal:

Feed supplements containing **Phly-N-Worm 11** which cannot be used should be buried in a safe place away from water supplies.

Container Disposal:

Do not reuse empty container. Wrap container and put in trash collection.

**HAZARDS TO HUMANS AND
DOMESTIC ANIMALS
WARNING**

Hazardous if swallowed, inhaled or absorbed through skin. Do not breathe vapors or spray mist. Do not get into eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking or smoking. Wear clean clothing after use. Contaminated clothing should be removed immediately and washed before reuse. Protective clothing, including impermeable gloves and a pesticide respirator, jointly approved by the Mining Enforcement and Safety Administration (formerly U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR 11, should be worn during spray mixing and loading operations in greenhouse applications and periods of prolonged use. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

ENVIRONMENTAL HAZARDS

ENVIRONMENTAL WARNINGS:
This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes, and estuaries (except for effluence from treated wastewater beds). Do not apply to water that will be used for recreational purposes and human and livestock consumption. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. This product must not be used in areas where impact on threatened/endangered species is likely. Contact your State Fish and Game Agency before applying this product. Apply this product only as specified on this label.

This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.
Do not store at temperatures below 30°F.

DIRECTIONS FOR USE

It is a violation of Federal Law to use "i" s product in a manner inconsistent with its labeling.

Re-entry Statement

Workers entering treated areas within 24 hours of application should wear protective clothing.

APPLICATION DIRECTIONS:

Apply the listed amount per acre when insects first appear and repeat as required to maintain effective control. For ground application use a minimum of 10 gallons of water per acre insuring thorough coverage of the plants. Consult Extension service recommendations for concentrate spraying with this formulation. Observe use limitations if insect control is required beyond and the following use patterns supplement the control program with other suitable pesticides. When applying this material by aircraft a minimum recommended amount is sufficient water to provide a minimum of 1 gallon finished spray per acre on deciduous trees and vines. Where more than 2 1/2 ft. of this material is recommended mix with sufficient water to provide a minimum of finished spray equal to twice the amount of Endosulfan SE C used. Do not plant root crops other than carrots, potatoes, sweet potatoes and sugar beets as follow up crops. The number of days between last application and harvest is indicated by the number in parentheses after the crop.

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal.

2. PESTICIDE DISPOSAL

Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

3. CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

4. GENERAL

Consult federal, state or local disposal authorities for approved alternate procedures.

NOTICE: Because Platte Chemical has no control over storage, handling, and conditions of use, which are of critical importance, Platte Chemical makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies due to misuse, improper handling or storage of this material. Nor does Platte Chemical assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

DISCLAIMER—It is impossible to eliminate all risks inherently associated with use of this product. Occupancy, maintenance, or other intended or unintended use may result in failure of such products under conditions not intended by the manufacturer or the original user of the product. The manufacturer and the original user warrant that the material is used in accordance with the use then intended when used in accordance with the instructions or subject to the risks referred to above. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential damages.

[illegible]

1950

**Insecticide Liquid NOS
RQ Endosulfan NA1993**

P
c/e

FORMULATED FOR **PL.**
14480.