

20 Pound Bag will Need & Feed 4,000 Sq. Ft.
40 Pound Bag will Need & Feed 8,000 Sq. Ft.



ACCEPTED
NOV 18 1986
The Federal Insecticide, Fungicide, and Rodenticide Act
EPA REG. NO. 32802-3

WEED & FEED
WITH 10-6-4 LAWN FOOD

FOR LAWNS AND ORNAMENTAL TURF

KILLS: DANDELIONS - PLANTAIN - BUCKHORN
AND SIMILAR BROADLEAF WEEDS

K I L L S W E E D S - F E E D S L A W N

FOR COOL SEASON TURF SUCH AS BLUEGRASS, RYEGRASS, FESCUE

ACTIVE INGREDIENT:	
Dimethylamine Salt of	
2,4-Dichlorophenoxyacetic acid	1.20%
INERT INGREDIENTS:	98.80%
	Total 100.00%

*Equivalent to 1.0% 2,4-Dichlorophenoxyacetic acid.

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

-----STATEMENTS OF PRACTICAL TREATMENT-----

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water immediately. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. If irritation persists get medical attention.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 32802-3

EPA EST. NO. 32802-WI-1 (A)
32802-MO-1 (B)

Net Contents 20 Pounds (40 Pounds)

HOWARD JOHNSON'S ENTERPRISES, INC.
VIROQUA, WI 54665

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Do not breathe dust. Avoid contact with skin, or clothing. Do not get in eyes. Wear glasses or goggles to protect eyes from dust. Thoroughly wash after handling and before eating or smoking.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to any body of water. Do not apply when weather conditions favor wind drift of product from target area since this product will injure beans, cucumbers, grapes, roses, tomatoes, and most vegetables, flowers and ornamental plants. Do not apply where run-off is likely to occur. Do not contaminate water used for irrigation or domestic purposes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original container in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Pesticides wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. HOUSEHOLD: Do not re-use empty bag. Wrap and put in trash.

GENERAL INFORMATION

This product may be used for control of many broadleaved weeds in Lawns and Ornamental Turf. It is most effective when weeds are small and actively growing and should be applied when the temperatures range from 65°F to 90°F. Do not apply on a windy day! If applied during extreme temperatures (below 65°F or above 95°F) weed control will be erratic. Do not apply within 2 hours of expected rain. If Weed & Feed is washed off within 2 hours after application retreat in 7 to 10 days. Reduced weed control and fertilizer "Streaking" will result if product is poorly applied.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	10%
2.40% Ammoniacal Nitrogen	
7.60% Urea Nitrogen	
Available Phosphoric Acid (P2O5)	6%
Soluble Potash (K2O)	4%
Derived from: Urea, Ammoniated Phosphate and Muriate of Potash.		
Calcium Carbonate equivalent 500 lbs. per ton.		

Supplies the following Nitrogen per 1,000 Sq. Ft. when used as Directed
20 lbs. 4,000 Sq. Ft. supplies lawn with 0.50 lbs. (N) nitrogen 1,000 Sq. Ft.

BEST AVAILABLE COPY

DIRECTIONS FOR USE (cont.)

Kills Weeds - Feeds Lawns

Controls Dandelions, Plantain, Buckhorn and Similar Broadleaf Weeds.

WHERE TO APPLY: Use on Bluegrass, Ryegrass, Fescue, Bermuda, or Zoysiagrass. Do not use on: Newly seeded lawns; Newly sprouted lawns until after the second mowing; Centipedegrass; Dichondra; Bentgrass; St. Augustinegrass or where desirable clovers are present.

WHEN TO APPLY: For best results apply to moist established lawns when temperatures range from 65°F to 90°F in the spring or early summer when weeds are young and actively growing. A Late summer or early fall application is also recommended to control broadleaf weeds that have developed from seeds during the summer. The lawn should be mowed two days prior to treatment. Wait 2 to 3 weeks to reseed areas when dead weeds can be removed by raking or tilling.

HOW TO APPLY: Select a suitable Lawn Fertilizer Spreader and apply to turf by "marking your spreader path" to avoid double applications or total missing of turf area. Apply the contents of this bag (20 lbs.) to cover 4,000 Sq.Ft. of turf area. Do not water in. Thoroughly water the lawn the 3rd or 4th day to move the fertilizer into the root zone to stimulate new lawn growth as weeds die.

SUGGESTED SPREADER SETTINGS


The following spreader settings are for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreaders condition with regard to age and use, your walking speed and the evenness of the terrain, your spreader may require a slight adjustment. If you have reason to believe that your spreader is not calibrated correctly test apply by placing the amount for 500 Sq.Ft. into your spreader and apply to your pre-measured 500 Sq.Ft. (5 ft. wide by 100 ft long)

HOW TO MAKE SURE YOUR SPREADER IS OPERATING CORRECTLY

For the 20 lb. rate to cover 4,000 Sq.Ft. Using a Scott Spreader place 2.50 lbs. Weed & Feed in your spreader using No.5 setting and apply to 500 Sq.Ft. With a 20 inch spreader this will equal 3 spreader widths wide (3 passes) by 100 feet long. If you run out before you cover the pre-measure area close the preader setting by 1/2 the number. If you have granules left over, open (increase) the spreader setting by 1/2 the number. For further help refer to your Spreader Guide Book.

this copy differs only in settings from basic label copy; as approved

SPREADER SETTINGS*	
Spreader Model	**20 lbs. per 4,000 Sq.Ft. =220#/Acre
Scott (20 inch)	5
Central	7
Cyclone Model B	2 1/2
Jackson	5
Modern	6
Lawn Beauty	6



**Equivalent to 40 lbs. per 8,000 Sq.Ft.
 NOTE: Broadcast spreaders such as the Cyclone are dependent on rate of walking and turning speed of the broadcast unit. Carefully calculate the area to be fertilized and make certain the proper number of pounds are applied to the area. *For settings when using a new spreader.