



the use of such systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the plants in the equipment on an approved site to avoid spills will help prevent undesirable contamination. A stopped copy is enclosed for your records.

Sincerely,

Jcarne J. Miller  
Product Manager (207)  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Enclosure:

BEST AVAILABLE COPY

385

HOWARD JOHNSON'S WEED & FEED

Active Ingredient:

Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid...	1.2 %
Inert Ingredients:.....	98.8 %
Total.....	100.0 %

Equivalent to by isomer specific assay AOAC method D01-5:  
100% of Dichlorophenoxyacetic Acid

EPA Reg. No. 32802-3

EPA Est. No. 32802-WI-1 (A)  
32802-MO-1 (B)

The first letter of code on end of package indicates location of manufacturing plant.

KEEP OUT OF REACH OF CHILDREN  
CAUTION

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED:

Get medical attention.

IF IN EYES:

Flush eyes with water for 15 minutes.  
Get medical attention.

IF ON SKIN:

Wash skin with soap and water.

ACCEPTED  
with COMMENTS  
by EPA Label Desk

OCT 04 1991

For the Medical Professional  
Add: ...  
32802-3

(Product is labeled for the following package weights and coverages.  
Only one weight/coverage sequence is on any one package.)

- 18 pounds covers 4,000 square feet
- 18 pounds covers 5,000 square feet
- 20 pounds covers 4,000 square feet
- 20 pounds covers 5,000 square feet

Manufactured by: Howard Johnson's Enterprises, Inc.  
Milwaukee, WI 53201

GUARANTEED FERTILIZER ANALYSIS

(Guaranteed Fertilizer Analysis will vary as to per cent of customer requirements.)

Total Nitrogen (N).....	10.0 %
2.3 % Ammoniacal Nitrogen	
2.7 % Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ).....	1.0 %
Soluble Potash (K <sub>2</sub> O).....	0.7 %

Plant foods derived from: Urea, ammoniated phosphate and muriate of potash.

BEST AVAILABLE COPY

**PRECAUTIONARY STATEMENTS**

**CAUTION:** Avoid inhalation of dust and contact with eyes, skin or clothing. Wear long pants, long-sleeved shirt and shoes when applying this product. Wash thoroughly after handling. Do not allow persons or pets on treated areas until dusts have settled.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water or wetlands (swamps, bays, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**STORAGE:** Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

**PRODUCT DISPOSAL:** To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**DIRECTIONS FOR USE, CONT.****GENERAL INFORMATION**

This product may be used for control of many broadleaf weeds in lawns and ornamental turf. It is most effective when weeds are small and actively growing and should be applied when the temperatures range from 65°F to 90°F. Do not apply on a windy day. If applied during extreme temperatures (below 65°F to above 90°F) weed control will be erratic. Do not apply within 2 hours of expected rain. If Weed & Feed is washed off within 2 hours after application retreat in 7 to 10 days. Reduced weed control and fertilizer "streaking" will result if product is poorly applied.

**DIRECTIONS**

Controls dandelions, plantain, buckhorn, and several broadleaf weeds.

**WHERE TO APPLY:** Use on bluegrass, ryegrass, fescue, bermuda or zoysia grass. Do not use on newly seeded lawns, newly mowed lawns until after the second mowing. Control weeds on dry hardy or best mix of St. Augustine grass or where desirable clovers are present.

**BEST AVAILABLE COPY**

WHEN TO APPLY: For best results apply to moist established lawns when temperatures range from 65°F to 90°F in the spring or early summer when weeds are young and actively growing. A late summer or early fall application is also recommended to control broadleaf weeds that have developed from seeds during the summer. The lawn should be mowed two days prior to treatment. Wait 2 to 3 weeks to reseed areas after application of Weed & Feed.

HOW TO APPLY: Select a suitable lawn fertilizer spreader and apply to lawn by "marking your spreader path" to avoid double applications or total missing of lawn area. Apply contents of this bag to cover area indicated on front of package. Do not water in. Thoroughly water the lawn the 3<sup>rd</sup> or 4<sup>th</sup> day to move the fertilizer into the root zone to stimulate new lawn growth as weeds die.

#### SUGGESTED SPREADER SETTINGS

The following spreader settings are for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreader's condition with regard to age and use, your walking speed and the evenness of the terrain, your spreader may require slight adjustment. If you have reason to believe that your spreader is not calibrated correctly test apply by placing the amount for 500 square feet into your spreader and apply to your pre-measured 500 square foot area.

#### SPREADER SETTINGS

(Note: These will be updated as needed to reflect market changes.)

Scott's 4.50  
Carousel 2.00  
Jackson 2.00  
Cyclone 3.50  
Modern 5.50  
Central 6.50

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

9-91

BEST AVAILABLE COPY