

PM 23 - 32802 - 3

1072

HOWARD JOHNSON'S WEED & FEED

ACTIVE INGREDIENT:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid - - - 1.0%

INERT INGREDIENTS: - - - - - 99.0%

Total 100.00%

*Equivalent to 0.83% 2,4-Dichlorophenoxyacetic acid Isomer Specific by AOAC Method No. 6.275;6.279(13th Edition).

20 Pound Bag will Weed & Feed 5,000 Sq. Ft.

Do not use on turf grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention. **IF IN EYES:** Flush eyes with water for 15 minutes. Get medical attention. **IF ON SKIN:** Wash skin with soap and water.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 32802-3 - EPA EST. NO. 32802-WI-1 (A) 32802-MO-1 (B) The first letter of code on end of package indicates location of manufacturing plant.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation of dust and contact with eyes, skin or clothing. When using this product, wear long pants, long-sleeved shirt, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor wind drift of product from target area since this product will injure beans, cucumbers, grapes, roses, tomatoes, and most vegetables, flowers and ornamental plants. Do not apply where run-off is likely to occur. Do not contaminate water used for irrigation or domestic purposes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. Prevent cross-contamination, do not store near other fertilizers or near herbicides, insecticides, fungicides or seeds. **PESTICIDE DISPOSAL:** To dispose of bag which contains unused product, securely wrap original container several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Thoroughly wrap original container in several layers of newspaper and discard in trash.

GENERAL INFORMATION

This product may be used for control of many broad-leaved weeds in Lawns and Ornamental Turf. It is most effective when weeds are small and actively growing and should be applied when the temperatures range from 65°F to 90°F. Do not apply on a windy day! If applied during extreme temperatures (below 65°F or above 95°F) weed control will be erratic. Do not apply within 2 hours of expected rain. If Weed & Feed is washed off within 2 hours after application retreat in 7 to 10 days. Reduced weed control and fertilizer "Streaking" will result if product is poorly applied.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - -	10%
2.40% Ammoniacal Nitrogen	
7.60% Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅) - - - - -	6%
Soluble Potash (K ₂ O) - - - - -	4%
Plant Food Derived from Urea, Ammoniated Phosphate and Muriate of Potash	

Controls Dandelions, Plantain, Buckhorn and Similar Broadleaf Weeds.

WHERE TO APPLY

Use on Bluegrass, Ryegrass, Fescue, Bermuda, or Zoysiagrass. Do not use on: Newly seeded lawns; Newly sprouted lawns until after the second mowing; Centipedegrass; Dichondra; Bentgrass; St. Augustinegrass or where desirable clovers are present.

WHEN TO APPLY

For best results apply to moist established lawns when temperatures range from 65°F to 90°F, in the spring or early summer when weeds are young and actively growing. A Late summer or early fall application is also recommended to control broadleaf weeds that have developed from seeds during the summer. The lawn should be mowed two days prior to treatment. Wait 2 to 3 weeks to reseed areas after application of Weed & Feed.

HOW TO APPLY

The maximum rate to turf is 1.45 pounds of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year. Select a suitable lawn fertilizer spreader and apply to turf by "marking your spreader path" to avoid double applications or total missing of turf area. Apply the contents of this bag to cover area indicated on front of package. Do not water in. Thoroughly water the lawn the 3rd or 4th day to move the fertilizer into the root zone to stimulate new lawn growth as weeds die.

SUGGESTED SPREADER SETTINGS

The following spreader settings are for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreaders condition with regard to age and use, your walking speed and the evenness of the terrain, your spreader may require a slight adjustment. If you have reason to believe that your spreader is not calibrated correctly test apply by placing the amount for 500 sq. ft. into your spreader and apply to your pre-measured 500 sq ft area.

SPREADER SETTINGS*

Scott (20 inch)	5
Central	7
Cyclone Model B	2 1/2
Jackson	5
Modern	6
Lawn Beauty	6

NOTICE Buyer and user assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label (32802-3 PM5 \ Subm'd 05-17-94 \ Acct'd 12-07-94)

Manufactured by



Net Contents 20 Pounds

Milwaukee, WI 5320

Manufactured By:

HOWARD JOHNSON'S ENTERPRISES, INC.

MILWAUKEE, WI 53102

1072

2 of 2

ACCEPTED
MAR 17 1995
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 32802-3

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

BEST COPY AVAILABLE

2