

# WEED & FEED

## WITH 2,4-D

### CONTROLS DANDELIONS, PLANTAINS, BUCKHORN & SIMILAR BROADLEAF WEEDS

**ACTIVE INGREDIENT:**

Dimethylamine salt of  
 2,4-dichlorophenoxyacetic acid ..... 1.0%  
**INERT INGREDIENTS:** ..... 99.0%  
**TOTAL:** ..... 100.0%

Equivalent to 0.83% 2,4-dichlorophenoxyacetic acid  
 Isomer Specific by AOAC Method No. 6.275;6.279 (13<sup>th</sup> Ed.)

EPA Reg. No. 32802-3

EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

First letter of bag code shows establishment where manufactured. See tag attached to bag for fertilizer guaranteed analysis.

Manufactured by:

**HOWARD JOHNSON'S ENTERPRISES, INC.**  
 700 W. Virginia Street, Milwaukee, WI 53204-1548

**KEEP OUT OF REACH OF CHILDREN  
 PRECAUTIONARY STATEMENTS  
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Avoid inhalation of dust and contact with eyes, skin or clothing. When using this product, wear long pants, long-sleeved shirt, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

**ENVIRONMENTAL HAZARDS:** Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor wind drift of product from target area since this product will injure beans, cucumbers, grapes, roses, tomatoes, and most vegetables, flowers and ornamental plants. Do not apply where runoff is likely to occur. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes.

**First Aid**

<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or physician immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or physician.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

**Hot line number:**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product may be used for control of many broad-leaved weeds in lawns and ornamental turf. It controls Dandelions, Plantain, Buckhorn and similar broadleaf weeds.

**WHERE TO APPLY:** Use on Bluegrass, Ryegrass, Fescue, Bermuda or Zoysiagrass. Do not use on: Newly seeded lawns; newly sprouted lawns until after the second mowing; Centipedegrass; Dichondra; Bentgrass; St. Augustine-grass or where desirable clovers are present.

**WHEN TO APPLY:** For best results apply to moist established lawns when temperatures range from 65°F to 90°F. in the spring or early summer when weeds are young and actively growing. A late summer or early fall application is also recommended to control broadleaf weeds that have developed from seeds during the summer. The lawn should be mowed two days prior to treatment. Wait two to three weeks to reseed areas after application of this product.

This product is most effective when weeds are small and actively growing and should be applied when the temperatures range from 65°F to 90°F. Do not apply on a windy day! If applied during extreme temperatures (below 65°F or above 95°F) weed control will be erratic. Do not apply within 2 hours of expected rain. If the product is washed off within 2 hours after application, retreat in 7 to 10 days. Reduced weed control and fertilizer "streaking" will result if product is poorly applied.

**HOW TO APPLY:** Apply this product at the rate of 4 pounds per 1000 square feet of turf. Do not water in. Thoroughly water the lawn the 3rd or 4th day to move the fertilizer into the root zone to stimulate new lawn growth as weeds die. The maximum rate to turf is 1.45 pounds of 2,4-D acid equivalent per acre, per application per site. The maximum number of broadcast applications per treatment site is two per year. Select a suitable lawn fertilizer spreader and apply to turf by "marking your spreader path" to avoid double applications or total missing of turf area.

The following spreader settings are for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreader's condition with regard to age and use, your walking speed and the evenness of the terrain, your spreader may require a slight adjustment. If you have reason to believe that your spreader is not calibrated correctly test apply by placing the amount for 500 square feet into your spreader and apply to your pre-measured 500 square foot area (5 ft. wide by 100 ft. long).

**SPREADER SETTINGS:**

The settings shown here are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed, and equipment condition. Calibrate spreader to ensure correct application rate. Clean and lubricate your spreader after each use.

Spreader setting to apply at listed rate (pounds per 1000 sq. ft.)	4 lbs.
Spyker 24-22 Rotary	4.1
Lesco Rotary	12(F)
Prizelawn CB Rotary	4.5
EarthWay 2400 Rotary	18
Cydome Pro	4.0
Scotts R8A	1
Lely Broadcast	3.0
Bigfoot	H-I

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. Prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

**PRODUCT DISPOSAL:** To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. (Alternate wording for household use:)

**IF EMPTY:** Do not reuse this container. Place in trash or offer for recycling if available.

**IF PARTLY FILLED:** Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain or toilet.

**NOTICE:** Buyer and user assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

**ACCEPTED**  
JAN 14 2003  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 32802-3

