

DIAZINON 5 G

**ALL
SEASON**

**ALL
SEASON**

ACCEPTED
with COMMENTS
in EPA Letter Dated:

OCT 08 1982

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
this product is registered under
EPA Reg. No. 106-100-01

DIAZINON 5 G INSECT CONTROL

**Controls Sod Web Worms, Chinch
Bugs, Cut Worms And Many Other
Lawn Insects**

(See Back Panel For Other Insects Controlled)

INSECT CONTROL

Controls Sod Web Worms, Chinch Bugs, Cut Worms And Many Other Lawn Insects

(See Back Panel For Other Insects Controlled)

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of food and feed products. Keep children and pets off treated areas until this material is washed into the soil and the area is dry.

NET WT. 10 LBS. (4.5 Kg)

TREATS UP TO 5,000 Sq. Ft.

ALL SEASON DIAZINON 5G INSECT CONTROL

32802-5

32802-WI-1

**ALL
SEASON**

**DIAZINON 5 G
INSECT CONTROL**

BEST DOCUMENT AVAILABLE



NET WT. 10 LBS. (4.5 Kg)

TREATS UP TO 5,000 Sq. Ft.



DIAZINON 5 G INSECT CONTROL

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

ALL SEASON Diazinon 5G controls the following lawn insects on grass and Dichondra lawns:

Ants, Armyworms, Bermuda Mites, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Hyperodes Weevil (New York only), Lawnmoths (Sod Webworms), Leafhoppers, Millipedes, Sowbugs, Springtails (Collembola).

Do not hand spread. Apply **All Season Diazinon Insect Control** to your entire lawn (or in a 5-foot band application around or near the foundation to control those insects which may enter a house) at a rate of 2 LBS. per 1,000 SQ. FT. of area (an area 20' by 50') or 3.2 OZ. per 100 SQ. FT. (an area 10' by 10'). For heavy-textured lawns such as St. Augustine, use 3 LBS. per 1,000 SQ. FT. (or 4.8 OZ. per 100 SQ. FT.). Apply when grass is dry. Water lawn lightly after application. Repeat as necessary.

All Season Diazinon Insect Control also controls Billbugs and White Grubs of Japanese Beetle, European Chafer, Southern Chafer and *Ategenius spretulus* (Dung Beetle). Apply at 2.5 LBS. per 1,000 SQ. FT. For Billbugs, apply when activity is first observed, or when chewed or brown grass indicates damage. For Grub control, excluding Dung Beetle, treat any time between late July and early October. For Dung Beetle Grubs, apply in spring or early summer when damage is first noticed. Thoroughly water grass with $\frac{1}{2}$ - $\frac{3}{4}$ inch of water after application. Repeat treatment in late summer or early fall if a second generation is in evidence. For Hyperodes Weevil (New York only), apply 2 LBS. per 1,000 SQ. FT. in mid-April and again in mid-May, watering thoroughly after each application. See back panel for recommended spreader settings.

All Season Diazinon Insect Control can be used for localized control of ants and Digger Wasps. Apply 1 1/2 teaspoons over and around each anthill, or in and around each wasp ground nest opening. Apply to wasp nests around dusk, when activity is low.

For garden insect control, apply **All Season Diazinon Insect Control** as listed for crops below. Apply just before planting and immediately mix into soil to the specified depth.

Cutworms, above and below ground: Broccoli, Cabbage, Cantaloupe, Cauliflower, Celery, Collards, Cucumbers, Endive (Escarole), Kale, Lettuce, Lima Beans, Muskmelons, Parsley, Peas, Peppers, Potatoes, Snap Beans, Spinach, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash. Use 7 to 14 OZ. per 500 SQ. FT. and mix into soil to a depth of 2-3 inches for control of surface cutworms and 3-6 inches for control of cutworms below the surface. Mix into soil 4-8 inches deep for potatoes.

Mole Crickets: Broccoli, Cabbage, Cauliflower, Collards, Endive (Escarole), Kale, Lettuce, Mustard, Peppers, Radishes, Strawberries, Tomatoes, Turnips. Use 4 OZ. per 500 SQ. FT. and mix into soil to a depth of 1-2 inches.

Wireworms: Broccoli, Cabbage, Cantaloupe, Carrots, Cauliflower, Celery, Collards, Corn, Cucumbers, Endive (Escarole), Kale, Lettuce, Lima Beans, Muskmelons, Onions, Parsley, Peas, Peppers, Snap Beans, Spinach, Summer Squash, Sweet Potatoes, Swiss Chard, Tomatoes, Watermelons, Winter Squash. Use 7 to 14 OZ. per 500 SQ. FT. and mix into soil to a depth of 4-8 inches.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Keep container closed and away from food, feedstuffs and domestic water supplies. Do not reuse empty container. Wrap container and put in trash collection.

Note to Physician: "If signs of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine."

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife feeding on treated areas may be killed. Keep out of lakes, streams, and ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STATEMENT OF PRACTICAL TREATMENT

Call a physician or Poison Control Center immediately

If swallowed — Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled — Remove victim to fresh air. Apply respiration if indicated.

If on skin — Remove contaminated clothing and wash affected areas with soap and water.

If in eyes — Flush eyes for at least 15 minutes with water. Call a physician if irritation persists.

WARRANTY STATEMENT

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

All Season Dinitro 5G controls the following lawn insects on grass and Dichondra lawns:

Ants, Armyworms, Bermuda Mites, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Hyperodes Weevil (New York only), Lawnmoths (Sod Webworms), Leafhoppers, Millipedes, Sowbugs, Springtails (Collembola).

Do not hand spread. Apply **All Season Dinitro Insect Control** to your entire lawn (or in a 5-foot band application around or near the foundation to control those insects which may enter a house) at a rate of 2 LBS. per 1,000 SQ. FT. of area (an area 20' by 50') or 3.2 OZ. per 100 SQ. FT. (an area 10' by 10'). For heavy-textured lawns such as St. Augustine, use 3 LBS. per 1,000 SQ. FT. (or 4.8 OZ. per 100 SQ. FT.). Apply when grass is dry. Water lawn lightly after application. Repeat as necessary.

All Season Dinitro Insect Control also controls Billbugs and White Grubs of Japanese Beetle, European Chafer, Southern Chafer and *Ategenus spretus* (Dung Beetle). Apply at 2.5 LBS. per 1,000 SQ. FT. For Billbugs, apply when activity is first observed, or when chewed or brown grass indicates damage. For Grub control, excluding Dung Beetle, treat any time between late July and early October. For Dung Beetle Grubs, apply in spring or early summer when damage is first noticed. Thoroughly water grass with $\frac{1}{2}$ - $\frac{3}{4}$ inch of water after application. Repeat treatment in late summer or early fall if a second generation is in evidence. For Hyperodes Weevil (New York only), apply 2 LBS. per 1,000 SQ. FT. in mid-April and again in mid-May, watering thoroughly after each application. See back panel for recommended spreader settings.

All Season Dinitro Insect Control can be used for localized control of ants and Digger Wasps. Apply 1½ teaspoons over and around each anthill, or in and around each wasp ground nest opening. Apply to wasp nests around dusk, when activity is low.

For garden insect control, apply **All Season Dinitro Insect Control** as listed for crops below. Apply just before planting and immediately mix into soil to the specified depth.

Cutworms, above and below ground: Broccoli, Cabbage, Cantaloupe, Cauliflower, Celery, Collards, Cucumbers, Endive (Escarole), Kale, Lettuce, Lima Beans, Muskmelons, Parsley, Peas, Peppers, Potatoes, Snap Beans, Spinach, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash.

Use 7 to 14 OZ. per 500 SQ. FT. and mix into soil to a depth of 2-3 inches for control of surface cutworms and 3-6 inches for control of cutworms below the surface. Mix into soil 4-6 inches deep for potatoes.

Mole Crickets: Broccoli, Cabbage, Cauliflower, Collards, Endive (Escarole), Kale, Lettuce, Mustard, Peppers, Radishes, Strawberries, Tomatoes, Turnips. Use 4 OZ. per 500 SQ. FT. and mix into soil to a depth of 1-2 inches.

Wireworms: Broccoli, Cabbage, Cantaloupe, Carrots, Cauliflower, Celery, Collards, Corn, Cucumbers, Endive (Escarole), Kale, Lettuce, Lima Beans, Muskmelons, Onions, Parsley, Peas, Peppers, Snap Beans, Spinach, Summer Squash, Sweet Potatoes, Swiss Chard, Tomatoes, Watermelons, Winter Squash. Use 7 to 14 OZ. per 500 SQ. FT. and mix into soil to a depth of 4-6 inches.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Keep container closed and away from food, feedstuffs and domestic water supplies. Do not reuse empty container. Wrap container and put in trash collection.

Note to Physician: "If signs of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine."

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife feeding on treated areas may be killed. Keep out of lakes, streams, and ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STATEMENT OF PRACTICAL TREATMENT

Call a physician or Poison Control Center immediately.

If swallowed — Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled — Remove victim to fresh air. Apply respiration if indicated.

If on skin — Remove contaminated clothing and wash affected areas with soap and water.

If in eyes — Flush eyes for at least 15 minutes with water. Call a physician if irritation persists.

WARRANTY STATEMENT

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with any directions provided, with proper equipment in good working order and under normal conditions of use. Buyer assumes the risk of any contrary use. Sellers make no other express or implied warranty including the warranty of fitness for a particular purpose, and no agent of seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Distributed by: Howard Johnson Enterprises Inc., Viroqua, Wisconsin 54655

NOTE: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary, in order to assure correct rate of application.

*SPREADER SETTING NUMBER (per 1000 sq. ft.)

INSECTS (see label on bag)

SPREADER - Make & Model	Grass or Dichondra Lawns	St. Augustine Lawn	WHITE GRUBS, BILLBUGS
Central Imperial Mark No. 1622-C	6½	7½	7
Central Suburban Lawn Spreader - No. 1622-B	6½	7½	7
Cushman Model B1 Battery	6½	8	6½
Crownfield Model 300	6½	7½	6½
Orlita Drive Spreader Model 3000	4½	6½	5
Orlita Broadcast Spreader Model 6000	4½	7	6
J. C. Penney - Model 6000	7½	8	7½
J. C. Penney Battery Spreader	5	7	6
Scot - Model 20	6	8	6½
Spay-Craftman Model 671-19100	6	7	6½

161 / 76152
18 / 1

32802-5

OCT 8 1982

All Season Dioxinon 5G

Howard Johnson's Enterprises, Inc.
P. O. Box 67, Railroad Avenue
Viroqua, WI 54655

Done
10/8/82

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment:

- a. Add the phrase "EPA Registration No. 32802-5."
- b. Insert the word "Dioxinon" immediately in front of its chemical formula in the statement of active ingredients.
- c. Insert the signal word "Caution" below "Keep out of reach of children." In our letter of June 7, 1982, we asked you to only delete the signal word after Hazards to Humans and Domestic Animals.
- d. In the State of Practical Treatment, revise the "If in eyes" statement to "Flush eyes with plenty of water. Call a physician if irritation persists."

BEST DOCUMENT AVAILABLE

2. The note to physicians in the Storage and Disposal Statement would be better located at the bottom of the Statement of Practical Treatment.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(a). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (13)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure