

**CRABGRASS AND MOST OTHER
ANNUAL WEED GRASSES
ON LAWNS AND GOLF COURSES**

IN ESTABLISHED BLUEGRASS (PERENNIAL), RYEGRASS (PERENNIAL), FESCUE, BAHIAGRASS, BERMUDAGRASS, CENTIPEDEGRASS, ST. AUGUSTINEGRASS AND ZOYSIAGRASS

ACTIVE INGREDIENT:
N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine 2.5%
INERT INGREDIENTS: 97.5%
TOTAL: 100.0%

NOTIFICATION

EPA Reg. No. 32802-7
EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B) **DEC 6 1996**
First letter of bag code shows establishment where manufactured.

Sold by:
HOWARD JOHNSON'S ENTERPRISES, INC.
700 W. Virginia Street, Milwaukee, WI 53204-1548

*BALAN and the BALAN logo are trademarks of DowElanco

In areas of the United States where germination of these annual weed grass species continues for extended periods of time, a second application of this product at the recommended rate may be made in 8 to 10 weeks after initial treatment for continued control.

ANNUAL BLUEGRASS (*Poa annua*): In areas of heavy infestation of Annual Bluegrass, its elimination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by Annual Bluegrass.

NORTHERN (COOL SEASON) TURFGRASS: For pre-emergence control, apply 80 pounds of this product (2 pounds active ingredient) per acre in late summer or early fall, and again in late winter or early spring 1 to 2 weeks prior to the expected germination of Annual Bluegrass.

SOUTHERN (WARM SEASON) TURFGRASS: For pre-emergence control, apply 120 pounds of this product (3 pounds active ingredient) per acre in late summer or early fall 1 to 2 weeks prior to the expected germination of Annual Bluegrass. In the southern areas of the United States where germination of Annual Bluegrass continues for extended periods of time, a second application of this product at 120 pounds per acre can be made in 10 to 12 weeks after the initial treatment.

SPREADER SETTINGS:

Approximate settings as a guide for calibration for applying this product in one pass over the turf. Spreader may need calibration, due to differences in the terrain, walking speed and equipment condition. Calibrate before use to ensure correct application rate. Clean and lubricate spreader after each use.

Spreader setting to apply at listed rate (lbs per acre)	60 lbs	80 lbs	120 lbs
Spyker 24-22 Rotary	3.5	3.6	3.9
Lesco Rotary	9(D-E)	10(E)	11(E-F)
Prizelawn CB Rotary	3.7	3.8	4.2
EarthWay 2400 Rotary	13	14	15
Cyclone Pro	3.2	3.4	3.5
Scotts R8A	F	F-G	G
Lely Broadcast	1.5	2.3	2.5
Bigfoot	E-F	F	G

**KEEP OUT OF REACH OF CHILDREN
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling and before eating or smoking.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor run-off or drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its label.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a pre-emergence herbicide which is applied directly to established turf for the control of Smooth Crabgrass, Hairy Crabgrass, Goosegrass (Silver Crabgrass), Annual Bluegrass (*Poa annua*), Barnyardgrass (Watergrass), and Green and Yellow Foxtail. It controls annual grass weeds by killing their seeds as they germinate. It does not control established weeds. When it is applied as directed the following established grasses are tolerant to this product: Perennial Bluegrasses, Perennial Ryegrass, Fescue, Bahiagrass, Bermudagrass, Centipedegrass, Zoysiagrass and St. Augustinegrass. Apply to turf before germination of annual grass weeds. Do not reseed within 42 days after application.

FOR APPLICATION TO SMALLER AREAS: Apply the following amount of this product:

Amount per acre	Amount per 1000 square feet
60 pounds / acre	1.5 pounds / 1000 sq ft
80 pounds / acre	2 pounds / 1000 sq ft
120 pounds / acre	3 pounds / 1000 sq ft

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store product in original container in a cool, dry, locked place out of reach of children.

Pesticide disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE: Buyer assumes all risk and liability for use, storage and/or handling of this material not in accordance with the terms of this label.

Net weight:

BEST AVAILABLE COPY