

JAN 30 1997

Mr. Thomas J. Schmit
Howard Johnson's Enterprises, Inc.
700 W. Virginia St., #222
Milwaukee, WI 53204-1548

Dear Mr. Schmit:

Subject: Worker Protection Standards-Exclusion Statement
All Season Crabgrass Preventer Plus
EPA Reg. No. 32802-8
Your submission dated Nov. 13, 1996

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable and a stamped copy is enclosed for your records.

Sincerely yours,

Philip V. Errico
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

2013

ALL SEASON CRABGRASS PREVENTER PLUS

25-3-3 FERTILIZER WITH 0.7% IRON

FOR USE ON HOME LAWNS AND ORNAMENTAL TURF FOR CONTROL OF CRABGRASS, GOOSEGRASS, FOXTAIL, WATERGRASS AND ANNUAL BLUEGRASS (*Poa annua*)

CONTAINS BALAN* A SELECTIVE PREEMERGENCE HERBICIDE FOR USE ON THE FOLLOWING ESTABLISHED NORTHERN AND SOUTHERN LAWNS AND TURF: BLUEGRASS (Perennial), RYEGRASS (Perennial), FESCUE, BAHIA GRASS, BERMUDAGRASS, CENTIPEDEGRASS, ST. AUGUSTINEGRASS & ZOYSIAGRASS

ACTIVE INGREDIENT:

N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine 0.92%

INERT INGREDIENTS: 99.08%

TOTAL: 100.00%

EPA Reg. No. 32802-8

EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

First letter of bag code shows establishment where manufactured.

Manufactured by:

HOWARD JOHNSON'S ENTERPRISES, INC.
 100 W. Virginia Street, Milwaukee, WI 53204-1548

*BALAN - Registered trademarks of DowElanco

NET WEIGHT: 50 POUNDS

a spring application to Bermuda lawns that have been overseeded with winter grasses since thinning may occur. Do not use on Dichondra, nor to newly sprigged areas of Bermudagrass, St. Augustinegrass, or Centipedegrass. Do not use on bentgrass putting greens. Use only on well established turfgrasses since injury may occur to young tender growth. This product may thin Annual Bluegrass or fine-leaved Fescue lawns when used at higher rates than 50 lb/12,500 sq. ft. Do not overdose. Follow directions carefully.

APPROPRIATE TIMES TO APPLY THIS PRODUCT FOR CONTROL OF GERMINATING WEED SEEDS											
For Northern Cool-Season Grasses:											
Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec
						1					
2											
For Southern Warm-Season Grasses:											
Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec
									1		
2											
1. Annual Bluegrass (<i>Poa annua</i>) Germination Periods											
2. Crabgrass, Foxtail, Goosegrass, and Watergrass Germination Periods											

The above chart gives suggested timing for application since the successful pre-emergence control of the annual grass weeds listed on this label requires proper timing of application. Apply approximately 1 to 2 weeks ahead of the germination of the annual grass weeds. Any cultural practices that disturb the soil such as aerification or verticutting, should be done prior to this application. Optimum weed control performance will be obtained if the treated areas are irrigated soon after application.

NORTHERN (COOL SEASON) TURFGRASS: For control of Crabgrass, Foxtail, and Barnyardgrass apply the contents of this bag to cover 12,500 sq. ft. For control of Annual Bluegrass (*Poa annua*), apply 50 pounds to cover 10,000 sq. ft of troublesome areas.

SOUTHERN (WARM SEASON) TURFGRASS: For control of Crabgrass, Foxtail, and Barnyardgrass apply the contents of this bag to cover 10,000 sq. ft. For weed seeds that germinate over an extended period of time a second treatment may be made in 8 to 10 weeks. For control of Annual Bluegrass (*Poa annua*), apply 50 pounds to cover 6750 sq. ft. When seed germination continues for an extended period of time a second application can be made at the same rate 10 to 12 weeks after the initial treatment.

RESEEDING THE AREA: Reseeding of any desirable turf grass seed should be delayed for at least 6 weeks after using this product at the rate of 50 pounds/10,000 - 12,500 sq. ft. Best results will be obtained when the amount of seed per 1000 sq. ft. is increased 25%.

LAWN APPLICATORS AND RATES: Make sure of your spreader calibration. Use a "Drop Type" or "Cyclone Type".

SPREADER MAKE	NORTHERN GRASSES		SOUTHERN GRASSES	
	Crabgrass	Poa annua	Crabgrass	Poa annua
Scott #35	5-1/2	6	6	7
Cyclone B-1	4	3-2/3	3-2/3	3

STORAGE AND DISPOSAL
 Do not contaminate water, food, or feed by storage or disposal.
Storage: Store product in original container in a cool, dry, locked place out of reach of children.
Disposal: Do not re-use empty bag. Wrap and put in trash.

NOTICE: Buyer assumes all risk and liability for use, storage and/or handling of this material not in accordance with the terms of this label.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	96	25%
1.3% Ammoniacal Nitrogen		
20.0% Urea Nitrogen		
3.7% Sulfur Coated Urea		
Available Phosphate (P2O5)	NOV 25	3%
Soluble Potash (K2O)		3%
Sulfur (S)	25	1.0%
Iron (Fe)		0.7%
Primary Plant Food Derived From: Diammonium Phosphate, Urea, Sulfur Coated Urea, and Muriate of Potash. Sulfur Derived From Sulfur Coated Urea. Iron Derived From Ferric Oxide. Potential Acidity: 1012 lbs. Calcium Carbonate per ton.		

KEEP OUT OF REACH OF CHILDREN CAUTION
 See back panel for additional precautionary statements and statement of practical treatment.

PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling and before eating or smoking.
STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
IF IN EYES: Flush eyes with plenty of water. Get medical attention.
IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water.
ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor run-off or drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its label.
 Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
 This product contains a pre-emergence herbicide plus a complete lawn food which should be applied directly to established turf for the control of Smooth and Hairy Crabgrass, Goosegrass (Silver Crabgrass or Clawfoot), Annual Bluegrass (*Poa annua*), Barnyardgrass (Watergrass), and Green and Yellow Foxtail.
WHEN, WHERE AND HOW TO USE: Apply this product to the turf prior to the germination of the seeds of weedy grasses since established weeds already present will not be controlled. Do not make

ACCEPTED
with COMMENTS
In EPA Letter Dated

JAN 30 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

32802-1