


"Your Money's worth and a Little Bit More"
TESTED AND GUARANTEED



26%	4%	5%
Total Nitrogen	Available Phosphoric Acid	Soluble Potash

Treats up to 5,000 Sq.Ft.

SLOW RELEASE NITROGEN FOR LONG-TERM RESULTS - - -

WEED KILLER-Prevents Crabgrass From Getting Started!

FERTILIZER-To Promote a Healthier, Thicker Lawn.
"For Greener Grass Leaves"
"For Healthier Root Systems"
"For More Vigorous Plant Growth"

CRABGRASS PREVENTER plus TURF FOOD

ACCEPTED
JUN 10 1985
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 32802-9

FOR USE ON HOME LAWNS AND ORNAMENTAL TURF -
FOR CONTROL OF CRABGRASS, FOXTAIL, ANNUAL BLUEGRASS(Poa Annua), GOOSEGRASS AND WATERGRASS.

Contains BALAN[®] a Selective Pre-Emergence Herbicide
For Use On The Following Established Northern and Southern Lawns and Turf: BLUEGRASS(Perennial), RYGRASS (Perennial), FESCUE, BAHIAGRASS, BERMUDAGRASS, CENTIPEDEGRASS, ST.AUGUSTINEGRASS AND ZOYSIAGRASS

BALAN[®]- Reg. Trademark for BENEFIN; Elanco Products Company

ACTIVE INGREDIENT:
N-butyl-N-ethyl,alpha,alpha,alpha,trifluoro-2,6-dinitro-p-toluidine 1.02%

INERT INGREDIENTS: 98.98%

Total 100.00%

EPA Reg. No. 32802-

EPA Est. No. 32802-WI-1(a);MO-1(B)

KEEP OUT OF REACH OF CHILDREN - 12 PT.
CAUTION - 18 PT SIZE TYPE

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. IF IN EYES: Flush eyes with plenty of water. Get medical attention. IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water.

See Back Panel For Additional Precautionary Statements

NET WEIGHT 18 POUNDS (8.20 Kg.)


Howard Johnson Enterprises Inc., Viroqua, Wisconsin 54665

OPTIONAL FRONT PANEL COPY - TO BE INCLUDED IN SUBMITTED LABELING FOR ACCEPTANCE

FOR BEST RESULTS, APPLY THIS PRODUCT DURING THE FOLLOWING SEASON(S):

ALL SEASON -[[EARLY SPRING]]- LATE SPRING - SUMMER - FALL

- # Apply Crabgrass Preventer Plus only once - in the spring - and stop crabgrass before it has a chance to start. One application prevents crabgrass all year long. Also stops Foxtail, Barnyardgrass, but won't harm lawn grasses.
- # Contains Benefin, a highly effective crabgrass controller. Apply in the spring before temperatures reach 70-75°F to prevent seed germination.
- # The "Plus" in (This Product) means your lawn will get a generous feeding of the essential nutrients necessary to get your lawn off to a healthy start when applied in the early spring.
- # In addition to a full-feeding, an early application of (This Product) will work to keep weed seed from germinating, thus providing your lawn with full-scale protection for an entire growing season.



EPA FILE SYMBOL: 32802-

PRECAUTIONARY STATEMENTS

Hazards To Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling and before eating or smoking.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor run-off or drift from treated areas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original container in a cool, dry, locked place out of reach of children. DISPOSAL: Do not re-use empty bag. Wrap and put in trash.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product contains a pre-emergence herbicide plus a complete lawn food which should be applied directly to established turf for the control of Annual Bluegrass (*Poa annua*), Smooth and Hairy Crabgrass, Goosegrass (Silver Crabgrass or Crowfoot), Barnyardgrass (Watergrass) and Green and Yellow Foxtail. When properly applied it supplies Northern Grasses with 0.94 lb. of nitrogen per 1,000 Sq. Ft. and 1.17 lb. Nitrogen / 1,000 Sq. Ft. to Southern Grasses.

FOR THE ALTERNATE FORMULATION OF INERTS THAT CONTAINS A FERTILIZER FORMULA THE LAST SENTENCE MUST BE CHANGED TO THE FOLLOWING
When properly applied it supplies Northern Grasses with _____ lb. of nitrogen per 1,000 Sq. Ft. and _____ lbs. Nitrogen / 1,000 Sq. Ft. to Southern Grasses.

WHERE, WHEN AND HOW TO USE: Apply this product to the turf prior to the germination of the seeds of weedy grasses since established weeds, already present, will not be controlled. Do not make a spring application to Bermuda lawns that have been overseeded with winter grasses since thinning may occur. Do not use (Bentgrass putting greens. Do not use on Dichondra, nor to newly sprigged areas of Bermudagrass, St. Augustinegrass or Centipedegrass. Use only on well established turfgrasses since injury may occur to young tender growth. This product may thin Bluegrass or Fescue lawns when used at higher rates than 18 lbs./5,000 Sq. Ft. - Do not over-dose. Follow Directions carefully.

APPROPRIATE TIMES TO APPLY THIS PRODUCT FOR CONTROL OF GERMINATING WEED SEEDS

	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
1												
2												
3												
4												

- 1. *Poa Annua* (Annual Blue Grass) Germination periods.
- 2. Crabgrass, Foxtail, Goosegrass and Watergrass Germination periods.

NOTICE: Buyer assumes all risk and liability of use, storage, and/or handling of this material not in accordance with the terms of this label.
(This to appear at the end of all Label Copy)

WHERE, WHEN AND HOW TO USE: (Cont.)

The above chart gives suggested timing for application since the successful pre-emergence control of the annual grass weeds listed on this label require proper timing of application. Apply approximately one to two weeks ahead of the germination of the annual grass weeds. Any cultural practices that disturb the soil, such as aerification or verti-cutting, should be done prior to this application. Optimum weed control performance will be obtained if the treated areas are irrigated soon after application

NORTHERN (cool season) TURFGRASS: For control of Crabgrass, Foxtail and Barnyardgrass apply the contents of this bag to cover 5,000 Sq.Ft. Apply evenly over the turf using a properly calibrated fertilizer spreader. Refer to Spreader Settings for NORTHERN GRASSES. - For control of Poa Annua apply 18 lbs. to cover 4,000 Sq.Ft. of troublesome areas. (2 lbs. active Ingredient per acre). Refer to the special Spreader setting for Poa Annua to apply this rate. If Poa Annua is growing in patches treat only those areas to avoid thinning of established Bluegrass or Fine Leaved Fescues.

SOUTHERN (warm season) TURFGRASS: For control of Crabgrass, Foxtail and Barnyardgrass apply the contents of this bag to cover 4,000 Sq.Ft. (50'x 80'). Apply evenly over the turf by using the Special Spreader Setting for SOUTHERN GRASSES. For weed seeds that germinate over an extended period a second treatment may be made in 8 to 10 weeks. For control of Poa Annua apply 18 lbs. to cover 2,700 Sq.Ft. (50 ft. x 55 ft.). Refer to the Special Spreader Setting for this application rate. When seed germination continues for an extended period of time a second application can be made, at the same rate 10 to 12 weeks after the initial treatment.


RESEEDING THE AREA

Reseeding of any desirable turf grass seed should be delayed for at least 6 weeks after using this product at the rate of 18 lbs./4,000 to 5,000 Sq.Ft. Wait 12 to 16 weeks when applying at the rate of 18 lbs./2,700 Sq.Ft. Best results will be obtained when the amount of seed per 1,000 Sq.Ft. is increased by 25%.

LAWN APPLICATORS AND RATES

Make sure of your spreader calibration. Use a "Drop Type" or "Cyclone Type".

SPREADER MAKE	NORTHERN GRASSES		SOUTHERN GRASSES	
		Poa Annua		Poa Annua
Scott #35	4-3/4 to 5-1/2	5-6	5-6	7 to 7-1/2
Cyclone B-1	3-3/4 to 4	4-1/2	4-1/2	4-3/4 to 5.0



Always calibrate your own spreader according to the manufacturer's recommendations.

COMPARISON APPLICATION RATES

AREA	POUNDS/PRODUCT/SQUARE FEET/LBS.		BALAN/ACRE	
Northern	18 lbs.	to treat	5,000 Sq.Ft.	
	159 lbs.	" "	1 Acre = 1.62 lbs.	Actual BALAN/Acre
Northern	18 lbs.	" "	4,000 Sq.Ft.	
(Poa)	198 lbs.	" "	1 Acre = 2.02 lbs.	" " "
Southern	18 lbs.	" "	4,000 Sq.Ft.	
	198 lbs.	" "	1 Acre = 2.02 lbs.	" " "
Southern	18 lbs.	" "	2,700 Sq.Ft.	
(Poa)	293 lbs.	" "	1 Acre = 2.98 lbs.	" " "

OPTIONAL WARRANTY

WARRANTY STATEMENT: Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with any directions provided, with proper equipment in good working order and under normal conditions of use. Buyer assumes the risk of any contrary use. Seller makes no other express or implied warranty including () warranty of fitness for a particular purpose, and no agent of seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen(N)	26.0%
Ammoniacal	1.6%
Urea Nitrogen	17.4%
Sulfur Coated Urea	7.0%
Available Phosphoric Acid (P ₂ O ₅)	4.0%
Soluble Potash (K ₂ O)	6.0%
Sulfur (S)	5.2%

Primary Plant Food derived from: Diammonium Phosphate, Urea, Sulfur Coated Urea, and Muriate of Potash. Sulfur derived from Sulfur Coated Urea & Sulfur.

Potential Acidity: 1116 lbs. Calcium Carbonate per ton.

OPTIONAL COPY

GUARANTEE: This product is formulated to [Company name] rigid specifications and is 100% guaranteed to perform to your satisfaction, if used as directed, or the purchase price will be refunded. (Limited to replacement or refund only, excluding any labor costs).

OPTIONAL WARRANTY

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

HOWARD JOHNSON ENTERPRISES, INC.
VIROQUA, WISCONSIN 54665

